

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 1 199 362 A2**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:

**24.04.2002 Bulletin 2002/17**

(51) Int Cl.<sup>7</sup>: **C12N 15/12**, C12N 15/62,  
C07K 14/705, C07K 19/00,  
C07K 16/28

(21) Application number: **01130678.4**

(22) Date of filing: **12.05.2000**

(84) Designated Contracting States:

**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE**

Designated Extension States:

**AL LT LV MK RO SI**

(30) Priority: **12.05.1999 US 310463**

(62) Document number(s) of the earlier application(s) in  
accordance with Art. 76 EPC:

**00928985.1 / 1 177 292**

(71) Applicant: **IMMUNEX CORPORATION**

**Seattle, WA 98101-2936 (US)**

(72) Inventors:

- **Cosman, David, J.**  
**Bainbridge Island, Washington 98110 (US)**

- **Anderson, Dirk, M.**  
**Seattle, Washington 98107 (US)**
- **Borges, Luis**  
**Seattle, Washington 98107 (US)**

(74) Representative: **Dörries, Hans Ulrich, Dr.**  
**Dörries, Frank-Molnia & Pohlman,**  
**Triftstrasse 13**  
**80538 München (DE)**

Remarks:

This application was filed on 21 - 12 - 2001 as a  
divisional application to the application mentioned  
under INID code 62.

(54) **Immunoregulator from the family designated 'leukocyte immunoglobulin-like receptors'  
(LIR)**

(57) A family of immunoreceptor molecules of the  
immunoglobulin superfamily, (LIR) polypeptides is de-  
scribed. Disclosed are sequences encoding LIR family  
members and their deduced amino acid sequences,  
polypeptides encoded by DNA that hybridizes to defined  
nucleotide sequences, processes for producing

polypeptides of the LIR family, and specific antibodies  
directed against LIR polypeptides. LIR family members  
can be used to treat autoimmune diseases and disease  
states associated with suppressed immune function.

**EP 1 199 362 A2**

## Description

**BACKGROUND OF THE INVENTION**

**[0001]** Immune system cellular activity is controlled by a complex network of cell surface interactions and associated signaling processes. When a cell surface receptor is activated by its ligand a signal is sent to the cell, depending upon the signal transduction pathway that is engaged, the signal can be inhibitory or activatory. For many receptor systems cellular activity is regulated by a balance between activatory signals and inhibitory signals. In some of these it is known that positive signals associated with the engagement of a cell surface receptor by its ligand are downmodulated or inhibited by negative signals sent by the engagement of a different cell surface receptor by its ligand.

**[0002]** The biochemical mechanisms of these positive and negative signaling pathways have been studied for a number of known immune system receptor and ligand interactions. Many receptors that mediate positive signaling have cytoplasmic tails containing sites of tyrosine phosphatase phosphorylation known as immunoreceptor tyrosine-based activation motifs (ITAM). A common mechanistic pathway for positive signaling involves the activation of tyrosine kinases which phosphorylate sites on the cytoplasmic domains of the receptors and on other signaling molecules. Once the receptors are phosphorylated, binding sites for signal transduction molecules are created which initiate the signaling pathways and activate the cell. The inhibitory pathways involve receptors having immunoreceptor tyrosine based inhibitory motifs (ITIM) which, like the ITAMs, are phosphorylated by tyrosine kinases. Receptors having these motifs are involved in inhibitory signaling because these motifs provide binding sites for tyrosine phosphatases which block signaling by removing tyrosine from activated receptors or signal transduction molecules. While many of the details of the activation and inhibitory mechanisms are unknown, it is clear that functional balance in the immune system depends upon opposing activatory and inhibitory signals.

**[0003]** One example of immune system activity that is regulated by a balance of positive and negative signaling is B cell proliferation. The B cell antigen receptor is a B cell surface immunoglobulin which, when bound to antigen, mediates a positive signal leading to B cell proliferation. However, B cells also express FcγRIIb1, a low affinity IgG receptor. When an antigen is part of an immune complex with soluble immunoglobulin, the immune complex can bind B cells by engaging both the B cell antigen receptor via the antigen and FcγRIIb1 via the soluble immunoglobulin. Co-engagement of the FcγRIIb1 with the B cell receptor complex downmodulates the activation signal and prevents B cell proliferation. FcγRIIb1 receptors contain ITIM motifs which are thought to deliver inhibitory signals to B cells via interaction of the ITIMs with tyrosine phosphatases upon co-engagement with B cell receptors.

**[0004]** The cytolytic activity of Natural Killer (NK) cells is another example of immune system activity which is regulated by a balance between positive signals that initiate cell function and inhibitory signals which prevent the activity. The receptors that activate NK cytotoxic activity are not fully understood. However, if the target cells express cell-surface MHC class I antigens for which the NK cell has a specific receptor, the target cell is protected from NK killing. These specific receptors, known as Killer Inhibitory Receptors (KIRs) send a negative signal when engaged by their MHC ligand, downregulating NK cell cytotoxic activity.

**[0005]** KIRs belong to the immunoglobulin superfamily or the C-type lectin family (see Lanier et al., *Immunology Today* 17:86-91, 1996). Known human NK KIRs are members of the immunoglobulin superfamily and display differences and similarities in their extracellular, transmembrane and cytoplasmic regions. A cytoplasmic domain amino acid sequence common to many of the KIRs is an ITIM motif having the sequence YxxL/V. In some cases, it has been shown that phosphorylated ITIMs recruit tyrosine phosphatases which dephosphorylate molecules in the signal transduction pathway and prevent cell activation (see Burshtyn et al., *Immunity* 4:77-85, 1996). The KIRs commonly have two of these motifs spaced apart by 26 amino acids [YxxL/V(x)<sub>26</sub>YxxL/V]. At least two NK cell receptors, each specific for a human leukocyte antigen (HLA) C allele (an MHC class I molecule), exist as an inhibitory and an activatory receptor. These receptors are highly homologous in the extracellular portions, but have major differences in their transmembrane and cytoplasmic portions. One of the differences is the appearance of the ITIM motif in the inhibitory receptor and the lack of the ITIM motif in the activating receptor (see Biassoni et al., *Journal. Exp. Med.*, 183:645-650, 1996).

**[0006]** An immunoreceptor expressed by mouse mast cells, gp49B1, also a member of the immunoglobulin superfamily, is known to downregulate cell activation signals and contains a pair of ITIM motifs. gp49B1 shares a high degree of homology with human KIRs (Katz et al., *Cell Biology*, 93: 10809-10814, 1996). Mouse NK cells also express a family of immunoreceptors, the Ly49 family, which contain the ITIM motif and function in a manner similar to human KIRs. However, the Ly49 immunoreceptors have no structural homology with human KIRs and contain an extracellular C-type lectin domain, making them a member of the lectin superfamily of molecules (see Lanier et al., *Immunology Today* 17:86-91, 1996).

**[0007]** Clearly, the immune system activatory and inhibitory signals mediated by opposing kinases and phosphatases are very important for maintaining balance in the immune system. Systems with a predominance of activatory signals will lead to autoimmunity and inflammation. Immune systems with a predominance of inhibitory signals are less able to challenge infected cells or cancer cells. Isolating new activatory or inhibitory receptors is highly desirable for studying

the biological signal(s) transduced *via* the receptor. Additionally, identifying such molecules provides a means of regulating and treating diseased states associated with autoimmunity, inflammation and infection.

[0008] For example engaging a newly discovered cell surface receptor having ITIM motifs with an agonistic antibody or ligand can be used to downregulate a cell function in disease states in which the immune system is overactive and excessive inflammation or immunopathology is present. On the other hand, using an antagonistic antibody specific to the receptor or a soluble form of the receptor can be used to block the interaction of the cell surface receptor with the receptor's ligand to activate the specific immune function in disease states associated with suppressed immune function. Conversely, since receptors lacking the ITIM motif send activatory signals once engaged as described above, the effect of antibodies and soluble receptors is the opposite of that just described.

## **SUMMARY OF THE INVENTION**

[0009] The present invention provides a new family of immunoreceptor molecules of the immunoglobulin superfamily, designated herein as the Leukocyte Immunoglobulin-Like Receptor (LIR) polypeptides. Within the scope of the present invention are DNA sequences encoding LIR family members and their deduced amino acid sequences disclosed herein. Further included in the present invention are polypeptides encoded by DNA that hybridize to oligonucleotide probes having defined sequences or to DNA or RNA complementary to the probes. The present invention also includes recombinant expression vectors comprising DNA encoding LIR family members. Also within the scope of the present invention are nucleotide sequences which, due to the degeneracy of the genetic code, encode polypeptides that are identical to polypeptides encoded by the nucleic acid sequences described above, and sequences complementary to those nucleotide sequences.

[0010] Further, the present invention includes processes for producing polypeptides of the LIR family by culturing host cells transformed with a recombinant expression vector that contains an LIR family member encoding DNA sequence under conditions appropriate for expressing an LIR polypeptide family member, then recovering the expressed LIR polypeptide from the culture.

[0011] The invention also provides agonistic and antagonistic antibodies to LIR family proteins.

[0012] Further still within the present invention are fusion proteins that include a soluble portion of an LIR family member and the Fc portion of Ig.

[0013] Certain autoimmune disorders are associated with the failure of a negative signaling LIR to downregulate cell function. Such disorders may be treated by administering a therapeutically effective amount of an agonistic antibody or ligand of one or more a LIR family member to a patient afflicted with such a disorder. Disorders mediated by disease states associated with suppressed immune function can be treated by administering a soluble form of the negative signaling LIR. Conversely, disorders mediated by diseases associated with failure of a activatory signaling LIR can be treated by administering an agonistic antibody of the activatory receptor. Disorders mediated by states associated with autoimmune function can be treated by administering a soluble form of the activatory receptor.

## **DETAILED DESCRIPTION OF THE INVENTION**

[0014] A viral glycoprotein having a sequence similarity to MHC class I antigens has been used to isolate and identify a new polypeptide, designated LIR-P3G2, and several members of a new family of cell surface polypeptides that has been designated the LIR polypeptide family. The present invention encompasses isolated nucleic acid molecules encoding LIR polypeptides, and further encompasses isolated LIR polypeptides. Exemplary nucleic acids encoding LIR polypeptides according to the present invention include those nucleotide sequences shown in SEQ ID NOS:1, 3, 7, 9, 11, 13, 15, 17, 19, 21, 29, 31, 33, 35 and 37, and exemplary LIR polypeptide sequences are shown in SEQ ID NOS: 2, 4, 8, 10, 12, 14, 16, 18, 20, 22, 30, 32, 34, 36 and 38.

[0015] The LIR polypeptide family members possess extracellular regions having immunoglobulin-like domains, placing the members in a new subfamily of the immunoglobulin superfamily. While the LIR family members are characterized as having very similar extracellular portions, the family includes three groups of polypeptides that are distinguishable by their transmembrane regions and their cytoplasmic regions. One group of the LIR polypeptides has a transmembrane region that includes a positively charged residue and a short cytoplasmic tail and a second group has a nonpolar transmembrane region and a long cytoplasmic tail. A third group includes polypeptides expressed as soluble proteins having no transmembrane region or cytoplasmic tail. One of the LIR proteins has characteristics of both groups one and two, and may represent a fourth group. A number of recent reports have described nucleic acid molecules having sequences related to the LIR family of proteins (Hillier et al., GenBank Accession Number N95687, April 9, 1996; Colonna, M., GenBank Accession Nos. AF041261 and AF041262, January 7, 1999; Lamerdin et al., GenBank Accession No. AC006293, January 6, 1999; Steffans et al., GenBank Accession Nos. AH007466 and AH007465, March 4, 1999; Cosman et al., *Immunity* 7:273-282 (1997); Borges et al., *J. Immunol.* 159:5192-96 (1997); Samaridis and Colonna, *Eur. J. Immunol* 27:660-665 (1997); Colonna et al., *J. Exp. Med.* 186:1809-1818 (1997); Wagtman et al., *Curr. Biol.*

7:615-618 (1997); Rojo et al., *J. Immunol.* 158:9-12 (1997); Arm et al., *J. Immunol.* 159:2342-2349 (1997); Cella et al., *J. Exp. Med.* 185:1743-51 (1997); Torkar et al., *Eur. J. Immunol.* 28:3959-67 (1998); Yamashita et al., *J. Biochem.* 123:358-68 (1998); WO 98/31806; WO 98/24906; WO 98/09638).

**[0016]** The LIR polypeptides encompassed by the subject invention contain at least one Ig-like domain in the extra-cellular region of the protein, preferably contain either two or four Ig-like domains in the extracellular region. Some LIR polypeptides may contain more than four Ig-like domains. An Ig-like domain is a structural unit that has been identified in a wide variety of cellular proteins. Ig-like domains contain a common fold that forms a sandwich of two  $\beta$  sheets that is stabilized by a characteristic intrachain disulfide bond. Ig-like domains are readily recognizable by reference to a large body of knowledge concerning this structural entity (see, e.g., Williams and Barclay, *Ann. Rev. Immunol.* 6:381-405 (1988)). Typically, Ig-like domains contain about 100 amino acids, although the number of amino acids may vary, e.g., from about 85 to 105 amino acids. Molecules that exhibit Ig-like domains generally play a recognition role at the cell surface, often mediating cell-cell interactions in a variety of biological systems.

**[0017]** LIR-P3G2 (SEQ ID NO:2) is expressed by a variety of cells and recognizes HLA-B44 molecules, HLA-A2 MHC molecules and the alleles described in Example 14. Another LIR family member, designated LIR-pbm8 (SEQ ID NO:9) is expressed by a variety of cells and also recognizes a number of MHC class I molecules. By analogy with known molecules, LIR-P3G2, LIR-pbm8 and LIR members have a role in immune recognition and self/nonself discrimination.

**[0018]** Examples 1-3 below describe isolating cDNA encoding P3G2 (LIR-P3G2) and a substantially identical polypeptide designated 18A3 (LIR-18A3). Briefly, the LIR-P3G2 family member was isolated by first expressing UL18, a Class I MHC-like molecule and using UL18 to isolate and identify P3G2 and 18A3, which are closely related and probably are variants of the same gene, which is designated "LIR-1." The nucleotide sequences of the isolated P3G2 cDNA and 18A3 cDNA are presented in SEQ ID NO:1 and SEQ ID NO:3, respectively. The amino acid sequences encoded by the cDNA presented in SEQ ID NO:1 and SEQ ID NO:3 are presented in SEQ ID NO:2 and SEQ ID NO:4, respectively. The P3G2 amino acid sequence (SEQ ID NO:2) has a predicted extracellular domain of 458 amino acids (1-458) including a signal peptide of 16 amino acids (amino acids 1-16); a transmembrane domain of 25 amino acids (amino acids 459-483) and, a cytoplasmic domain of 167 amino acids (amino acids 484-650). The extracellular domain includes four immunoglobulin-like domains. Ig-like domain I includes approximately amino acids 17-118; Ig-like domain II includes approximately amino acids 119-220; Ig-like domain III includes approximately amino acids 221-318; and Ig-like domain IV includes approximately amino acids 319-419. Significantly, the cytoplasmic domain of this polypeptide includes four ITIM motifs, each having the consensus sequence of YxxL/V. The first ITIM motif pair is found at amino acids 533-536 and 562-565 and the second pair is found at amino acids 614-617 and 644-647. This feature is identical to the ITIM motifs found in KIRs except that KIRs contain only one pair of ITIM motifs.

**[0019]** The 18A3 amino acid sequence (SEQ ID NO:4) has a predicted extracellular region of 459 amino acids (1-459) including a signal peptide of 16 amino acids (amino acids 1-16); a transmembrane domain of 25 amino acids (amino acids 460-484) and a cytoplasmic domain of 168 amino acids (485-652). The 18A3 amino acids sequence (SEQ ID NO:4) is substantially identical to that of P3G2 (SEQ ID NO:2) except that 18A3 has two additional amino acids (at amino acid 438 and 552) and 18A3 possesses an isoleucine residue at amino acid 142 in contrast to a threonine residue for P3G2. Additionally, 18A3 has a serine residue at amino acid 155 and P3G2 has an isoleucine at 155. Finally, the 18A3 polypeptide has a glutamic acid at amino acid 627 and P3G2 has a lysine at 625 which is aligned with the 627 residue of the 18A3 polypeptide. The four ITIM motifs in the 18A3 cytoplasmic domain are at amino acids 534-537 and 564-567 and at 616-619 and 646-649. Glycosylation sites occur at the amino acid triplet Asn-X-Y, where X is any amino acid except Pro and Y is Ser or Thr. Thus, potential glycosylation sites on LIR-P3G2 occur at amino acids 140-142; 281-283; 302-304; and 341-343. Sites on LIR-18A3 are at 281-283; 302-304; and 341-343. The features of these encoded polypeptides are consistent with type I transmembrane glycoproteins.

**[0020]** Examples 8-10 describe isolating and identifying eight additional LIR polypeptide family members by probing cDNA libraries for plasmids that hybridize to a probe obtained from DNA encoding the extracellular region of LIR-P3G2. The nucleotide sequences (cDNA) of the isolated LIR family members are presented in SEQ ID NO:7 (designated pbm25, or LIR-4), SEQ ID NO:9 (designated pbm8, or LIR-2), SEQ ID NO:11 (designated pbm36-2, or LIR-6b), SEQ ID NO:13 (designated pbm36-4, or LIR-6a); SEQ ID NO:15 (designated pbmhh, or LIR-7); SEQ ID NO:17 (designated pbm2, or LIR-5), SEQ ID NO:19 (designated pbm17, or LIR-3) and SEQ ID NO:21 (designated pbmnew, or LIR-8). The amino acid sequences encoded thereby are presented in SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO: 12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO: 18, SEQ ID NO: 20 and SEQ ID NO:22, respectively.

**[0021]** Example 15 describes the isolation of LIR-9m1 (SEQ ID NOS:29, 30), LIR-9m2 (SEQ ID NO:31, 32), LIR-9s1 (SEQ ID NO:33, 34), and LIR-9s2 (SEQ ID NO:35, 36), which are four alternatively spliced variants of LIR-9, another new member of the LIR family. The first step in identifying these LIR-9 group of clones was the isolation of a short cDNA clone that was obtained from a human dendritic cell library and whose sequence analysis indicated that it had significant homology with the LIR family, particularly with the sequences shown in SEQ ID NOS:11, 13 and 15. Using PCR primers based on this clone, further cloning efforts yielded four full-length cDNAs corresponding to LIR-9m1,

-9m2, -9s1 and -9s2. LIR-9m1 and LIR-9m2 are transmembrane proteins that differ by 12 amino acids that are found in the extracellular region of LIR-9m1, but that are absent from LIR9m2. These 12 amino acids correspond to amino acids 29-40 of SEQ ID NO:30. LIRs-9s1 and -9s2 do not contain a transmembrane domain, thus encode soluble versions of LIR-9. The LIR-9s1 polypeptide (SEQ ID NO:34) includes the 12 amino acid insert that is present in LIR-9m1. Amino acids 1-238 of LIR-9s1 (SEQ ID NO:34) and LIR-9m1 (SEQ ID NO:30) are identical, but the remainder of the LIR-9s1 sequence is not identical to the corresponding region of LIR-9m1. Amino acids 1-226 of LIR-9s2 (SEQ ID NO:36) are identical to the first 226 amino acids of LIR-9m2 (SEQ ID NO:32), but the remaining amino acid sequence of LIR-9s2 diverges from that of LIR-9m2.

**[0022]** The same PCR primers that were used to isolate the LIR-9 clones yielded an additional cloned LIR cDNA that has been designated LIR-10 (SEQ ID NOS:37 and 38). By comparing the nucleotide sequence of LIR-10 with the most closely related LIRs that were previously identified, i.e. with SEQ ID NOS:13 and 15, it has been determined that the LIR-10 cDNA is an incomplete clone that lacks sequences located at the 5' end of the corresponding mRNA, including the 5' untranslated region, and nucleotides encoding the first 26 amino acids of the LIR-10 protein.

**[0023]** The identified extracellular, transmembrane and cytoplasmic regions for the polypeptides of LIR family members shown in SEQ ID NOS:10, 12, 14, 16, 18, 20, 22, 30, 32, 34, 36 and 38 are presented below. The polypeptides presented in SEQ ID NOS:8, 34 and 36 are soluble proteins having no transmembrane or cytoplasmic regions. As will be understood by the skilled artisan, the transmembrane region of P3G2 and 18A3 described above and those of LIR polypeptide family members presented below are identified in accordance with conventional criteria for identifying hydrophobic domains associated with such regions. Accordingly, the precise boundaries of any selected transmembrane region may vary from those presented herein. Typically, the transmembrane domain does not vary by more than five amino acids on either end of the domain as described herein. Computer programs known in the art and useful for identifying such hydrophobic regions in proteins are available.

**[0024]** The polypeptide presented in SEQ ID NO:8 (LIR-pbm25) has an extracellular domain that includes the entire amino acid sequence of amino acids 1-439 and a signal peptide of amino acids 1-16. The amino acid sequence presented in SEQ ID NO:10 (LIR-pbm8) has a predicted extracellular region of 458 amino acids (1-458) including a 16 amino acid signal peptide (amino acids 1-16); a transmembrane domain that includes amino acids 459-483; and a cytoplasmic domain that includes amino acids 484-598. The extracellular domain includes four immunoglobulin-like domains and the cytoplasmic domain includes an ITIM motif at amino acids 533-536 and 562-565.

**[0025]** The amino acid sequence presented in SEQ ID NO:12 (LIR-pbm36-2) has a predicted extracellular domain of amino acids including a 16 amino acid signal peptide of from amino acids 1-16; a transmembrane domain which includes amino acids 262-280 and a cytoplasmic domain of from amino acids 281-289. The transmembrane domain includes a charged arginine residue at 264 and the cytoplasmic domain is short, having only a length of only 9 amino acids.

**[0026]** The amino acid sequence presented in SEQ ID NO:14 (LIR-pbm36-4) has a predicted extracellular domain of amino acids 1-461 including a signal peptide from amino acids 1-16; a transmembrane domain that includes amino acids 462-480 and possesses a charged arginine residue at amino acid 464; and a cytoplasmic domain that includes amino acids 481-489. SEQ ID NO:14 is nearly identical to that of SEQ ID NO:12 except that it possesses four immunoglobulin domains in contrast to the two domains found in the extracellular region of SEQ ID NO:12. The amino acid sequences presented in SEQ ID NO:12 and SEQ ID NO:14 are likely proteins encoded by alternatively spliced transcripts from the same gene.

**[0027]** The amino acid sequence presented in SEQ ID NO:16 (LIR-pbmhh) has a predicted extracellular domain that includes amino acids 1-449 and a signal peptide from amino acids 1-16; a transmembrane domain that includes amino acids 450-468 with a charged arginine residue at amino acid 452; and a cytoplasmic domain that includes amino acids 469-483. The cytoplasmic domain is short with a length of 15 amino acids. The extracellular domain includes four immunoglobulin-like domains.

**[0028]** The amino acid sequence presented in SEQ ID NO:18 (LIR-pbm2) has a predicted extracellular region that includes amino acids 1-259 and a signal peptide of amino acids 1-16; a transmembrane domain that includes amino acids 260-280; and a cytoplasmic domain that includes amino acids 281-448. This LIR family member has cytoplasmic domain which includes an ITIM motif at amino acids 412-415 and 442-445. The extracellular domain includes two immunoglobulin-like domains.

**[0029]** The amino acid sequence presented in SEQ ID NO:20 (LIR-pbm17) has a predicted extracellular domain of amino acids 1-443 that includes a signal peptide of amino acids 1-16; a transmembrane domain which includes amino acids 444-464; and a cytoplasmic domain of amino acids 465-631. The extracellular domain has four immunoglobulin-like domains. SEQ ID NO:20 has two pairs of ITIM YxxLV motifs in the cytoplasmic domain. A first pair is at amino acids 514-517 and 543-546, and a second pair is at amino acids 595-598 and 625-628.

**[0030]** The amino acid sequence presented in SEQ ID NO:22 (LIR-pbmnew) has a predicted extracellular domain of amino acids 1-456 including a signal peptide of amino acids 1-16; a transmembrane domain which includes amino acids 457-579; and a cytoplasmic domain of amino acids 580-590. The extracellular includes four immunoglobulin-like

domains. SEQ ID NO:22 has an ITIM motif at amino acids 554-557 and 584-587.

**[0031]** The LIR-9m1 protein has an extracellular domain located at amino acids 1-262 of SEQ ID NO:30, including a signal peptide at amino acids 1-34 of SEQ ID NO:30. Amino acids 263-284 of SEQ ID NO:30 define the transmembrane region of LIR-9m1, and amino acids 285-299 of SEQ ID NO:30 form the cytoplasmic region. For LIR-9m2, the extracellular region corresponds to amino acids 1-250 of SEQ ID NO:32, including a signal sequence at amino acids 1-35 of SEQ ID NO:32, a transmembrane region at residues 251-272 of SEQ ID NO:32, and a cytoplasmic region at amino acids 273-287 of SEQ ID NO:32. LIR-9s1 (SEQ ID NO:34) and LIR-9s2 (SEQ ID NO:36) consist, respectively, of 265 and 253 amino acids, with their signal sequences being found at amino acids 1-34 of SEQ ID NO:34, and amino acids 1-35 of SEQ ID NO:36.

**[0032]** For LIR-10, amino acids 1-393 of SEQ ID NO:38 correspond to most of the extracellular portion of the LIR-10 protein, though the coding sequences for about 26 amino acids at the amino terminus of this protein, including the signal peptide, are believed to be missing from the LIR-10 cDNA clone that is described herein. The transmembrane region of LIR-10 is defined by amino acids 394-417 of SEQ ID NO:38, and the intracellular region by amino acids 418-449. A single ITIM motif is located at amino acids 438-443 of SEQ ID NO:38.

**[0033]** The amino acid sequences presented in SEQ ID NO: 2, 4, 8, 10, 12, 14, 16, 18, 20, 22, 30, 32, 34, 36 and 38 reveal that the LIR family, with the exception of LIR-10, can be categorized into three groups of polypeptides. One group includes the polypeptides of SEQ ID NOS:12, 14, 16, 30 and 32, which are distinguishable by a charged arginine residue in their transmembrane regions and their short cytoplasmic regions. A second group includes SEQ ID NO: 2, 4, 10, 18, 20 and 22 which are distinguishable by their hydrophobic cytoplasmic domains and the presence of one or more ITIM motifs in their cytoplasmic regions. A third group includes the polypeptides of SEQ ID NOS: 8, 34 and 36, which are expressed as soluble polypeptides and have no transmembrane or cytoplasmic regions. These soluble polypeptides may function to block the interactions of cell surface family members with their receptors. Alternatively, the soluble polypeptides may act as an activatory signal when bound to the receptor. Like the members of group one, LIR-10 has a relatively short cytoplasmic domain and a charged residue in its transmembrane domain, though its charged residue is histidine instead of arginine. However, LIR-10 also has an ITIM motif in its cytoplasmic domain, like the members of group two. Thus, LIR-10 has some of the characteristics of both groups one and two, and may represent a fourth group of LIR proteins. The LIR polypeptides are characterized generally by the ability of their encoding DNA to hybridize to DNA encoding the P3G2 extracellular region.

**[0034]** The invention should be understood to encompass isolated nucleic acid molecules encoding LIR polypeptides having the amino acid sequences shown in SEQ ID NOS:2, 4, 8, 10, 12, 14, 16, 18, 20, 22, 30, 32, 34, 36 and 38. In one embodiment of the invention, these nucleic acid molecules have the nucleic acid sequences shown in SEQ ID NOS:1, 3, 7, 9, 11, 13, 15, 17, 19, 21, 29, 31, 33, 35 and 37.

**[0035]** The extracellular regions of the LIR family member proteins presented in SEQ ID NO:2, 4, 8, 10, 12, 14, 16, 18, 20, 22, 30, 32, 34, 36 and 38 have a high degree of homology, which varies from 59%-84. Several of the LIR isolates are closely related, thus must represent allelic variants or splicing variants. For example, the extracellular regions of SEQ ID NO: 12 and SEQ ID NO: 14 share sequence homology which is close to 100%, thus indicating that these polypeptides derive from the same gene. In addition, SEQ ID NOS:2 and 4 share sequence homology that is in excess of 95%, thus probably represent two alleles of the same gene. Moreover, as discussed above, the extracellular regions of SEQ ID NOS:30, 32, 34 and 36 are nearly identical, thus indicating that these four proteins derive from mRNAs that are splicing variants.

**[0036]** While sharing some structural similarities with other members of the immunoglobulin superfamily, the LIR family members have limited homology to other members of the immunoglobulin superfamily. Molecules having the closest structural similarity to the LIRs are the human KIRs and mouse gp49. However, LIR extracellular regions share only a 38-42% identity with the extracellular regions of NKAT3 and p58 C1-39, respectively. The extracellular regions of the LIR family members are only 35-47% homologous with that of mouse gp49. In contrast, KIRs in general are known to share at least a 80% amino acid identity, with NKAT3 and p58 CL-39 being 81% homologous. Additionally, none of the known KIR molecules has four extracellular immunoglobulin domains which is characteristic of all but two of the known LIR family members. In view of the high sequence homology among the LIR related polypeptides disclosed herein and their relatively low homology with KIRs, the LIR polypeptides are members of a new family of immunoregulators.

**[0037]** An analysis of the amino acid sequences of the LIR polypeptides reveals that specific stretches of amino acids of the LIR polypeptides are highly conserved. One conserved region is a sequence of 46 amino acids found at amino acids 5-50 of SEQ ID NO:2. A data base search determined that the LIR family members differ substantially from the most structurally similar prior art polypeptides in this LIR conserved region. The data base search and structural analysis was performed using BLAST NB1, a local alignment search tool for searching data bases and aligning amino acid sequences to determine identities and variations in a given sequence. The BLAST NB1 software is accessible on the internet at <http://www3.ncbi.nlm.nih.gov/entrez/blast>. The BLAST NB1 search for sequences having homology to the sequence of amino acids 5 to 50 of SEQ ID NO:2 found that the most structurally similar proteins are FcγIIR, gp49B

form 2, and gp49B form 1 having identities with amino acids 5 to 50 of SEQ ID NO:2 of 63%, 67%, and 67% respectively. This contrasts with an LIR family identity with amino acids 5 to 50 of SEQ ID NO:2 which ranges from about 71% to 100%. Specifically, LIR family members of the present invention contain conserved regions near their amino termini having the following identities with amino acids 5-50 of SEQ ID NO:2: SEQ ID NO:8 has a 96% identity; SEQ ID NO:10 has a 90% identity; SEQ ID NO:12 has a 96% identity; SEQ ID NO:14 has a 91% identity; SEQ ID NO:16 has a 97% identity; SEQ ID NO:18 has a 77% identity; SEQ ID NO:20 has an 80% identity; SEQ ID NO:22 has an 80% identity; SEQ ID NO:30 has a 78% identity; SEQ ID NO:32 has a 71% identity; SEQ ID NO:34 has a 78% identity; SEQ ID NO:36 has a 71% identity. This conserved region appears to be present also in LIR-10 (SEQ ID NO:38), but is incomplete due to the LIR-10 cDNA clone disclosed herein being truncated at its 5' end.

**[0038]** Sequence identity as used herein is the number of aligned amino acids which are identical, divided by the total number of amino acids in the shorter of the two sequences being compared. A number of computer programs are available commercially for aligning sequences and determining sequence identities and variations. These programs provide identity information based upon the above stated definition of identity. One suitable computer program is the GAP program, version 6.0, described by Devereux et al. (*Nucl. Acids Res.* 12:387, 1984) and available from the University of Wisconsin Genetics Computer Group (UWCGC). The GAP program utilizes the alignment method of Needleman and Wunsch (*J. Mol. Biol.* 48:443, 1970), as revised by Smith and Waterman (*Adv. Appl. Math* 2:482, 1981). The preferred default parameters for the GAP program include: (1) a unary comparison matrix (containing a value of 1 for identities and 0 for non-identities) for nucleotides or amino acids, and the weighted comparison matrix of Gribskov and Burgess, *Nucl. Acids Res.* 14:6745, 1986, as described by Schwartz and Dayhoff, eds., *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, pp. 353-358, 1979; (2) a penalty of 3.0 for each gap and an additional 0.10 penalty for each symbol in each gap; and (3) no penalty for end gaps. Another similar program, also available from the University of Wisconsin as part of the GCG computer package for sequence manipulation is the BESTFIT program.

**[0039]** In another aspect, the polypeptides of the present invention have conserved regions which are uniquely characterized as having the amino acid sequence (SEQ ID NO:28):

Leu Xaa<sub>a</sub> Leu Ser Xaa<sub>b</sub> Xaa<sub>c</sub> Pro Arg Thr Xaa<sub>d</sub> Xaa<sub>e</sub> Gln Xaa<sub>f</sub> Gly Xaa<sub>g</sub> Xaa<sub>h</sub> Pro  
Xaa<sub>i</sub> Pro Thr Leu Trp Ala Glu Pro Xaa<sub>j</sub> Ser Phe Ile Xaa<sub>j</sub> Xaa<sub>70</sub> Ser Asp Pro Lys  
Leu Xaa<sub>k</sub> Leu Val Xaa<sub>m</sub> Thr Gly,

where Xaa<sub>a</sub> is Gly or Arg; Xaa<sub>b</sub> is Leu or Val; Xaa<sub>c</sub> is Gly or Asp; Xaa<sub>d</sub> is His Arg or Cys; Xaa<sub>e</sub> is Val or Met; Xaa<sub>f</sub> is Ala or Thr; Xaa<sub>g</sub> is His Pro or Thr; Xaa<sub>h</sub> Leu Ile or Phe; Xaa<sub>i</sub> is Gly Asp or Ala; Xaa<sub>j</sub> is Thr Ile Ser or Ala; Xaa<sub>k</sub> is Gly or Val; Xaa<sub>m</sub> is Met or Ala; and Xaa<sub>70</sub> is a sequence of 70 amino acids.

**[0040]** As mentioned above, certain LIR family members have ITIM motifs (YxxL/V<sub>25-26</sub>YxxL/V) in their cytoplasmic domains. It is known that many immune regulating receptors such as KIRs, CD22, FcγRIIb1 also have ITIMs in their cytoplasmic domain and function to send inhibitory signals which down regulate or inhibit cell function. It has been shown that these receptors associate with SHP-1 phosphatase via binding to the ITIM motifs. Recruitment of the SHP-1 phosphatase by the receptor appears to be required for intracellular signaling pathways that regulate the inhibitory function of the receptors. The experiment described in Example 11 demonstrates that LIR-P3G2 and LIR-pbm8 polypeptides associate with SHP-1 phosphatase upon phosphorylation and generate inhibitory signals through monocyte activation pathways. It is known that many immune regulating receptors such as KIRs, CD22, FcγRIIb1 have ITIMs in their cytoplasmic domain and function to send inhibitory signals which down regulate or inhibit cell function. Thus, by analogy with KIRs, CD22 and FcγRIIb1, LIR family members presented in SEQ ID NO:2, 4, 10, 18, 20, 22 and 38 that have ITIM motifs deliver an inhibitory signal via the interaction of its ITIM with SHP-1 tyrosine phosphatase, or other tyrosine phosphatases, when the LIR is coligated with an appropriate receptor. Also by analogy with immunoregulatory receptors possessing ITIMs, LIR family members have a regulatory influence on humoral, inflammatory and allergic responses.

**[0041]** The LIR family members presented in SEQ ID NO:12, 14, 16, 30 and 32 have relatively short cytoplasmic domains, have transmembrane regions possessing at least one charged residue, and do not possess the ITIM motif. By analogy with membrane proteins that lack ITIM motifs and have charged transmembrane regions, these family members mediate stimulatory or activatory signals to cells. For example, membrane bound proteins containing a charged residue in the transmembrane regions are known to associate with other membrane-bound proteins that possess cytoplasmic tails having motifs known as immunoreceptor tyrosine-based activation motifs (ITAM). Upon association, the ITAMs become phosphorylated and propagate an activation signal.

**[0042]** The LIR polypeptide designated LIR-P3G2 is expressed on the surface of transfected or normal cells. This is evidenced by the results of the experiments described in Example 3 and Example 5 in which flow cytometry and precipitation techniques demonstrate that LIR-P3G2 is found on monocytes, a subpopulation of NK cells, and B cells. P3G2 was detected on small subset of T cells. P3G2 is expressed as a 110-120 kDa glycoprotein. Since P3G2 has



four potential glycosylation sites, the molecular size of this protein will vary with the degree of its glycosylation. Glycosylation sites occur at the amino acid triplet Asn-X-Y, where X is any amino acid except Pro and Y is Ser or Thr. Potential glycosylation sites on P3G2 occur at amino acids 139-141; 280-282; 302-304; and 340-342.

**[0043]** P3G2-LIR isolated as described in Example 3 was tested for its ability to bind to cell surface ligands distinct from UL18. As demonstrated by the experimental results detailed in Example 7, P3G2 binds HLA-B 44 and HLA-A2, class I MHC antigens. Similarly, as demonstrated in Example 14, LIR-P3G2 and LIR-pbm8 bind to a variety of HLA-A, -B, and -C alleles and recognize a broad spectrum of MHC class I specificities. Since Class I MHC molecules play a central role in immune surveillance, self/non-self discrimination, the immune response to infection etc., the LIR-P3G2 and LIR-pbm8 polypeptides have a role in regulation of immune responses. It is known that NK cytolytic activity for killing tumor cells and cells infected with a virus is regulated by a delicate modulation of activatory and inhibitory signals. It has been shown that receptors specific for the same HLA class I molecules to which LIR-P3G2 and LIR-pbm8 bind may be activatory or inhibitory in their triggering mechanism. By analogy, LIR-P3G2 and LIR-pbm8, which bind MHC class I molecules, play a role in balancing immune system cell activity and are useful in treating disease states in which the immune system balance is disrupted.

**[0044]** Within the scope of the present invention are polypeptides which include amino acid sequences encoded by DNA that hybridizes to LIR-P3G2 extracellular DNA probes under moderate to highly stringent conditions as taught herein. Probes that hybridize to DNA that encode polypeptides of the present invention include probes which encompass nucleotides 310-1684 of SEQ ID NO: 1 or fragments thereof. Fragments of SEQ ID NO:1 utilized as hybridization probes are preferably greater than 17 nucleotides in length, and more typically are greater than 20 nucleotides in length, and may include nucleotides 358-1684; nucleotides 322-459 (encoding LIR conserved sequence); or DNA or RNA sequences complementary to SEQ ID NOS:5, 6, 23, 24, 27 and 1 or fragments thereof. Fragments of SEQ ID NOS:5, 6, 23, 24 and 27 include these sequences without the restriction sites. The nucleotide sequences described herein also can be used to design PCR primers, for which a convenient length is about 17-30 nucleotides.

**[0045]** Conditions for hybridization may be moderately stringent conditions described in, for example, in Sambrook et al, *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press, 1989, which is hereby incorporated by reference (see, e.g., Vol. 1, pp 1.101-104). Conditions of moderate stringency, as defined by Sambrook et al., include, for example, the use of a prewashing solution containing 5x SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0) and hybridization conditions of about 55°C in 5x SSC, incubated overnight. Highly stringent conditions include higher temperatures of hybridization and washing. The skilled artisan will recognize that a given degree of stringency may be maintained while varying the hybridization or wash temperature or composition of the hybridization buffer in accord with formulae known to those in the art (e.g., see Sambrook et al., 9.50-9.51 and 11.45-11.47). Such formulae take into account factors such as the length of the probe, the G+C content of the probe, salt concentration of the hybridization buffer. If desired, formamide may be added to the hybridization buffer, which permits the use of lower hybridization temperatures (e.g., see Sambrook et al., 9.50-9.51).

**[0046]** Preferred embodiments include amino acid sequences encoded by DNA that hybridizes to probes of the extracellular region of LIR-P3G2 having at least 17 nucleotides. Preferred hybridizing conditions include an incubation temperature of 63°C for 16 hours in a solution of Denhart's solution, 0.05 M TRIS at pH 7.5, 0.9 M NaCl, 0.1% sodium pyrophosphate, 1% SDS and 200 µg/mL salmon sperm DNA, followed by washing with 2x SSC at 63°C for one hour and then washing with 1x SSC at 63°C for one hour. However, as explained above, one skilled in the art can devise other hybridization conditions that produce the same degree of stringency. Generally, stringent hybridization conditions involve a combination of buffer and incubation temperature that supports the formation of specific, i.e., well-matched duplexes while still allowing the formation of stable duplexes at an acceptable rate. Conditions of reduced stringency permit the formation of stable duplexes containing a higher degree of mismatched base pairs than can form under more stringent conditions.

**[0047]** Stringent hybridization conditions for PCR primers can be achieved, for example, by hybridizing labeled probes to filter-bound target nucleic acid overnight at 50-55° C in aqueous buffer containing 5 x SSC or 6 x SSC (1 x SSC=0.15 M NaCl, 0.015 M sodium citrate), followed by washes in 6 x SSC at 50-55° C. However, the skilled artisan will recognize that stringent hybridization conditions for oligonucleotide probes will vary, depending on the length, base composition and sequence of the probe (e.g., see Sambrook et al., 11.45-11.49).

**[0048]** The present invention includes polypeptides having amino acid sequences that differ from, but are highly homologous to, those presented in SEQ ID NOS:2, 4, 8, 10, 12, 14, 16, 18, 20, 22, 30, 32, 34, 36 and 38. Examples include, but are not limited to, homologs derived from other mammalian species, variants (both naturally occurring variants and those generated by recombinant DNA technology), and LIR P3G2 and LIR family member fragments that retain a desired biological activity. Preferably, such polypeptides exhibit a biological activity associated with the LIR polypeptides described in SEQ ID NOS:2, 4, 8, 10, 12, 14, 16, 18, 20, 22, 30, 32, 34, 36 and 38, and comprise an amino acid sequence that is at least 80% identical to any of the amino acid sequences of the signal peptide and extracellular domains of the polypeptides presented in SEQ ID NOS:2, 4, 8, 10, 12, 14, 16, 18, 20, 22, 30, 32, 34, 36 and 38. Preferably such polypeptides are at least 90% identical to any of the amino acid sequences of the signal peptide



and extracellular domains of the polypeptides presented in SEQ ID NOS: 2, 4, 8, 10, 12, 14, 16, 18, 20, 22, 30, 32, 34, 36 and 38. Determining the degree of identity between polypeptides can be achieved using any algorithms or computer programs designed for analyzing protein sequences. The commercially available GAP program described below is one such program. Other programs include the BESTFIT and GCG programs which are also commercially available.

**[0049]** Within the scope of the present invention are LIR polypeptide fragments that retain a desired biological property of an LIR polypeptide family member such as binding to MHC class I or other ligand. In one such embodiment, LIR polypeptide fragments are soluble LIR polypeptides comprising all or part of the extracellular domain, but lacking the transmembrane region that would cause retention of the polypeptide on a cell membrane. Soluble LIR polypeptides are capable of being secreted from the cells in which they are expressed. Advantageously, a heterologous signal peptide is fused to the N-terminus such that the soluble LIR is secreted upon expression. Soluble LIR polypeptides include extracellular domains incorporating the signal peptide and those in which the signal peptide is cleaved signal peptide.

**[0050]** The use of soluble forms of a LIR family member is advantageous for certain applications. One such advantage is the ease of purifying soluble forms from recombinant host cells. Since the soluble proteins are secreted from the cells, the protein need not be extracted from cells during the recovery process. Additionally, soluble proteins are generally more suitable for intravenous administration and can be used to block the interaction of cell surface LIR family members with their ligands in order to mediate a desirable immune function.

**[0051]** Further encompassed within the present invention are soluble LIR polypeptides, which may include the entire extracellular domain or any desirable fragment thereof, including extracellular domains that exclude signal peptides. Thus, for example, soluble LIR polypeptides include amino acids  $x_1$ -458 of SEQ ID NO:2, where  $x_1$  is amino acids 1 or 17; amino acids  $x_2$ -459 of SEQ ID NO:4, where  $x_2$  is amino acid 1 or 17; amino acids  $x_3$ -439 of SEQ ID NO:8, where  $x_3$  is amino acid 1 or 17; amino acids  $x_4$ -458 of SEQ ID NO:10, where  $x_4$  is amino acid 1 or 17; amino acids  $x_5$ -241 of SEQ ID NO:12, where amino acid  $x_5$  is amino acid 1 or 17; amino acids  $x_6$ -461 of SEQ ID NO:14, where  $x_6$  is amino acid 1 or 17; amino acids  $x_7$ -449 of SEQ ID NO:16, where  $x_7$  is amino acid 1 or 17; amino acids  $x_8$ -259 of SEQ ID NO:18, where  $x_8$  is amino acid 1 or 17; amino acids  $x_9$ -443 of SEQ ID NO:20, where  $x_9$  is amino acid 1 or 17; amino acids  $x_{10}$ -456 of SEQ ID NO:22, where  $x_{10}$  is amino acid 1 or 17; amino acids  $x_{11}$ -262 of SEQ ID NO:30, where  $x_{11}$  is amino acid 1 or 35; amino acids  $x_{12}$ -250 of SEQ ID NO:32, where  $x_{12}$  is amino acid 1 or 36; amino acids  $x_{13}$  of SEQ ID NO:34, where  $x_{13}$  is amino acid 1 or 35; amino acids  $x_{14}$  of SEQ ID NO:36, where  $x_{14}$  is amino acid 1 or 36; and amino acids 1-393 of SEQ ID NO:38. The above identified soluble LIR polypeptides include LIR extracellular regions that include and exclude signal peptides. Also encompassed herein are LIRs that lack a transmembrane and cytoplasmic region, such as SEQ ID NOS:8, 34 and 36. Additional soluble LIR polypeptides include fragments of the extracellular domains of family members that retain a desired biological activity, such as binding to ligands that include MHC class I molecules.

**[0052]** LIR family member fragments, including soluble polypeptides, may be prepared by any of a number of conventional techniques. A DNA sequence encoding a desired LIR polypeptide encoding fragment may be subcloned into an expression vector for production of the LIR polypeptide fragment. The selected encoding DNA sequence advantageously is fused to a sequence encoding a suitable leader or signal peptide. The desired LIR member encoding DNA fragment may be chemically synthesized using known DNA synthesis techniques. DNA fragments also may be produced by restriction endonuclease digestion of a full length cloned DNA sequence, and isolated by electrophoresis on an appropriate gel. If necessary, oligonucleotides that reconstruct the 5' or 3' terminus to a desired point may be ligated to a DNA fragment generated by restriction enzyme digestion. Such oligonucleotides may additionally contain a restriction endonuclease cleavage site upstream of the desired coding sequence, and position an initiation codon (ATG) at the N-terminus of the coding sequence.

**[0053]** Another technique useful for obtaining a DNA sequence encoding a desired protein fragment is the well-known polymerase chain reaction (PCR) procedure. Oligonucleotides which define the termini of the desired DNA are used as primers to synthesize additional DNA from a desired DNA template. The oligonucleotides may also contain recognition sites for restriction endonucleases, to facilitate inserting the amplified DNA fragment into an expression vector. PCR techniques are described, for example, in Saiki et al., *Science* 239:487(1988); *Recombinant DNA Methodology*, Wu et al., eds., Academic Press, Inc., San Diego (1989), pp. 189-196; and *PCR Protocols: A Guide to Methods and Applications*, Innis et al., eds., Academic Press, Inc. (1990).

**[0054]** The LIR nucleic acid molecules of the present invention include isolated cDNA, chemically synthesized DNA, DNA isolated by PCR, cloned genomic DNA, and combinations thereof. Genomic LIR family DNA may be isolated by hybridization to the LIR family cDNA disclosed herein using standard techniques. Isolated RNA transcribed from LIR family DNA molecules is also encompassed by the present invention.

**[0055]** Within the scope of the present invention are DNA fragments such as LIR polypeptide coding regions and DNA fragments that encode soluble polypeptides. Examples of DNA fragments that encode soluble polypeptides include DNA that encodes entire extracellular regions of LIR family members and DNA that encodes extracellular region

fragments such as regions lacking the signal peptide. More specifically, the present invention includes nucleotides 310-2262 of SEQ ID NO:1 (P3G2 coding region); nucleotides  $x_1$ -1683 of SEQ ID NO:1, where  $x_1$  is 310 or 358 (encoding the P3G2 extracellular domain); nucleotides 168-2126 of SEQ ID NO:3 (the 18A3 coding region) and nucleotides  $x_2$ -1544 of SEQ ID NO:3, where  $x_2$  is 168 or 216 (the 18A3 extracellular domain coding region); nucleotides  $x_3$ -1412 of SEQ ID NO:7, where  $x_3$  is 93 or 141 (the pbm25 coding region and extracellular region); nucleotides 184-1980 of SEQ ID NO:9, (the pbm8 coding region) and nucleotides  $x_4$ -1557 of SEQ ID NO:9, where  $x_4$  is 184 or 232 (the pbm8 extracellular domain coding region); nucleotides 171-1040 of SEQ ID NO:11 (pbm36-2 coding region) and nucleotides  $x_5$ -878 of SEQ ID NO:11, where  $x_5$  is 171 or 219 (encoding the pbm36-2 extracellular domain); nucleotides 183-1652 of SEQ ID NO:13 (coding region for pbm36-4) and nucleotides  $x_6$ -1565 of SEQ ID NO:13, where  $x_6$  is 183 or 231 (encoding the pbm36-4 extracellular domain); nucleotides 40-1491 of SEQ ID NO:15 (the pbmhh coding region) and nucleotides  $x_7$ -1386 of SEQ ID NO:15, where  $x_7$  is 40 or 88 (encoding the pbmhh extracellular domain); nucleotides 30-1376 of SEQ ID NO:17 (the pbm2 coding region) and nucleotides  $x_8$ -806 of SEQ ID NO:17, where  $x_8$  is 30 or 78 (encoding the pbm2 extracellular region); nucleotides 66-1961 of SEQ ID NO:19 (the pbm17 coding region) and nucleotides  $x_9$ -1394 of SEQ ID NO:19, where  $x_9$  is 66 or 114 (encoding the pbm17 extracellular domain); nucleotides 67-1839 of SEQ ID NO:21 (the pbmnew coding region) and nucleotides  $x_{10}$ -1434 of SEQ ID NO:21, where  $x_{10}$  is 67 or 115 (encoding the pbmnew extracellular domain); nucleotides 69-968 of SEQ ID NO:29 (the coding region of LIR-9m1) and nucleotides  $x_{11}$ -854 of SEQ ID NO:29, where  $x_{11}$  is 69 or 170 (encoding the LIR-9m1 extracellular domain); nucleotides 95-958 of SEQ ID NO:31 (the LIR-9m2 coding region) and nucleotides  $x_{12}$ -844 of SEQ ID NO:31, where  $x_{12}$  is 95 or 200 (encoding the LIR-9m2 extracellular domain); nucleotides  $x_{13}$ -912 of SEQ ID NO:33, where  $x_{13}$  is 115 or 216 (the LIR-9s1 coding region and extracellular region); nucleotides  $x_{14}$ -834 of SEQ ID NO:35, where  $x_{14}$  is 73 or 178 (the LIR-9s2 coding region and extracellular region); nucleotides 1-1350 of SEQ ID NO:37 (the LIR-10 coding region) and nucleotides 1-1179 of SEQ ID NO:37 (encoding all but a few amino-terminal amino acids of the LIR-10 extracellular domain).

**[0056]** Included in the present invention are DNAs encoding biologically active fragments of the LIR proteins whose amino acid sequences are presented in SEQ ID NOS:2, 4, 8, 10, 12, 14, 16, 18, 20, 22, 30, 32, 34, 36 and 38.

**[0057]** The present invention encompasses nucleotide sequences which, due to the degeneracy of the genetic code, encode polypeptides that are identical to polypeptides encoded by the nucleic acid sequences described above, and sequences complementary to them. Accordingly, within the present invention are DNA encoding biologically active LIR family members that include the coding region of a native human LIR family member cDNA, or fragments thereof, and DNA that is degenerate as a result of the genetic code to the native LIR polypeptide DNA sequence or the DNA of native LIR family members described herein.

**[0058]** In another aspect, the present invention includes LIR variants and derivatives as well as variants and derivatives of LIR family polypeptides, both recombinant and non-recombinant, that retain a desired biological activity. An LIR variant, as referred to herein, is a polypeptide substantially homologous to a native LIR polypeptide, as described herein, except the variant amino acid sequence differs from that of the native polypeptide because of one or more deletions, insertions or substitutions.

**[0059]** LIR family variants may be obtained from mutations of native LIR nucleotide sequences. Within the present invention are such DNA mutations or variants that include nucleotide sequences having one or more nucleotide additions, nucleotide deletions, or nucleotide substitutions compared to native DNA of LIR family members and that encode variant LIR polypeptides or variant LIR family members having a desired biological activity. Preferably the biological activity is substantially the same as that of the native LIR polypeptide.

**[0060]** Variant amino acid sequences and variant nucleotide sequences of the present invention preferably are at least 80% identical to that of a native LIR family member sequence. One method for determining the degree of homology or identity between a native amino acid or nucleotide sequence and a variant amino acid or nucleotide sequence is to compare the sequences using computer programs available for such purposes. One suitable computer program is the GAP program, version 6.0, described by Devereux et al. (*Nucl. Acids Res.* 12:387, 1984) and available from the University of Wisconsin Genetics Computer Group (UWGCG). The GAP program utilizes the alignment method of Needleman and Wunsch (*J. Mol. Biol.* 48:443, 1970), as revised by Smith and Waterman (*Adv. Appl. Math* 2:482, 1981). Briefly, the GAP program defines identity as the number of aligned symbols (i.e., nucleotides or amino acids) which are identical, divided by the total number of symbols in the shorter of the two sequences being compared. The preferred default parameters for the GAP program include: (1) a unary comparison matrix (containing a value of 1 for identities and 0 for non-identities) for nucleotides, and the weighted comparison matrix of Gribskov and Burgess, *Nucl. Acids Res.* 14:6745, 1986, as described by Schwartz and Dayhoff, eds., *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, pp. 353-358, 1979; (2) a penalty of 3.0 for each gap and an additional 0.10 penalty for each symbol in each gap; and (3) no penalty for end gaps.

**[0061]** Alterations of native LIR amino acid sequences may be provided by using any of a number of known techniques. As described above, mutations can be introduced at selected sequence sites by synthesizing oligonucleotides containing a mutant coding sequence, flanked by restriction sites enabling its ligation to fragments of the native se-

quence. After ligating the synthesized oligonucleotides to the native sequence fragments, the resulting reconstructed nucleotide sequence will encode an analog or variant polypeptide having the desired amino acid insertion, substitution, or deletion. Another procedure suitable for preparing variant polypeptides is oligonucleotide-directed site-specific mutagenesis procedures which provide genes having specific codons altered in accordance with the desired substitution, deletion, or insertion. Techniques for making such alterations include those disclosed in the following references: Walder et al. *Gene*, 42:133, 1986; Bauer et al., *Gene* 37:73, 1985; Craik, *BioTechniques*, 12-19 January, 1985; Smith et al. *Genetic Engineering: Principles and Methods*, Plenum Press, 1981; and U.S. Patent Nos. 4,518,584 and 4,737,462, all of which are incorporated herein by reference.

**[0062]** Variant polypeptides of the present invention may have amino acid sequences which are conservatively substituted, meaning that one or more amino acid residues of a native LIR polypeptide family member is replaced by different residues, such that the variant polypeptide retains a desired biological activity that is essentially equivalent to that of a native LIR family member. In general, a number of approaches to conservative substitutions are well known in the art and can be applied in preparing variant of the present invention. For example, amino acids of the native polypeptide sequence may be substituted for amino acids which do not alter the secondary and/or tertiary structure of the LIR polypeptide. Other suitable substitutions include those which involve amino acids outside of the ligand-binding domain of interest. One approach to conservative amino acid substitutions involves replacing one or amino acids with those having similar physiochemical characteristics, e.g. substituting one aliphatic residue for another such as Ile, Val, Leu, or Ala for one another); substituting one polar residue for another (such as between Lys and Arg; Glu and Asp; or Gln and Asn); or substituting entire regions having similar hydrophobicity or hydrophilic characteristics.

**[0063]** LIR polypeptide variants can be tested for binding to cells as described in Examples 5 and 6 and for phosphatase binding activity as described in Example 11 to confirm biological activity. Other LIR variants within the present invention include polypeptides which are altered by changing the nucleotide sequence encoding the polypeptide so that selected polypeptide Cys residues are deleted or replaced with one or more alternative amino acids. These LIR variants will not form intramolecular disulfide bridges upon renaturation. Naturally occurring LIR polypeptides selected for alteration by deleting or altering Cys residues preferably do not have biological activities which depend upon disulfide bridges formed by the Cys residue. Other possible variants are prepared by techniques which cause the modification of adjacent dibasic amino acid residues to enhance expression in yeast systems in which KEX2 protease activity is present. EP 212,914 discloses site-specific mutagenesis techniques for inactivating KEX2 protease processing sites in a protein. KEX2 protease processing sites are inactivated by deleting, adding or substituting residues to alter Arg-Arg, Arg-Lys, and Lys-Arg pairs to eliminate the occurrence of these adjacent basic residues. Lys-Lys and pairings are considerably less susceptible to KEX2 cleavage, and conversion of Arg-Lys or Lys-Arg to Lys-Lys represents a conservative and preferred approach to inactivating KEX2 sites.

**[0064]** Naturally occurring LIR variants are also encompassed by the present invention. Examples of such variants are proteins that result from alternative mRNA splicing events or from proteolytic cleavage of an LIR polypeptide. Alternative splicing of mRNA may yield a truncated but biologically active LIR polypeptide such as a naturally occurring soluble form of the protein. Variations attributable to proteolysis include difference in the N- or C- termini upon expression in different types of host cells, due to proteolytic removal of one or more terminal amino acids from the LIR polypeptide. In addition, proteolytic cleavage may release a soluble form of LIR from a membrane-bound form of the polypeptide. Other naturally occurring LIR variations are those in which differences from the amino acid sequence of SEQ ID Nos:2, 4, 8, 10, 12, 14, 16, 18, 20, 22, 30, 32, 34, 36 and 38 are attributable to genetic polymorphism, the allelic variation among individuals.

**[0065]** Within the scope of the present invention are derivative LIR family polypeptides which include native or variant LIR polypeptides modified to form conjugates with selected chemical moieties. The conjugates can be formed by covalently linking another moiety to a native or variant LIR or by non-covalently linking another moiety to a native or variant LIR. Suitable chemical moieties include but are not limited to glycosyl groups, lipids, phosphates, acetyl groups, and other proteins or fragments thereof. Techniques for covalently linking chemical moieties to proteins are well known in the art and are generally suitable for preparing derivative LIR polypeptides. For example, active or activated functional groups on amino acid side chains can be used as reaction sites for covalently linking a chemical moiety to a LIR polypeptide. Similarly, the N-terminus or C-terminus can provide a reaction site for a chemical moiety. LIR polypeptides or fragments conjugated with other proteins or protein fragments can be prepared in recombinant culture as N-terminal or C-terminal fusion products. For example, the conjugate or fusion portions may include a signal or leader sequence attached to an LIR molecule at its N-terminus. The signal or leader peptide co-translationally or post-translationally directs transfer of the conjugate from its site of synthesis to a site inside or outside of the cell membrane.

**[0066]** One useful LIR polypeptide conjugate is one incorporating a poly-His or the antigenic identification peptides described in U.S. Patent No. 5,011,912 and in Hopp et al., *BioTechnology* 6:1124, 1988. For example, the FLAG® peptide, Asp-Tyr-Lys-Asp-Asp-Asp-Lys (SEQ ID NO:39) is highly antigenic and provides an epitope reversibly bound by a specific monoclonal antibody, thus enabling rapid assay and facile purification of expressed recombinant protein. This sequence is specifically cleaved by bovine mucosal enterokinase at the residue immediately following

the Asp-Lys pairing. Fusion proteins capped with this peptide may be resistant to intracellular degradation in *E. coli*. Murine hybridoma designated 4E11 produced a monoclonal antibody that binds the peptide of SEQ ID NO:39 in the presence of certain divalent metal cations, and has been deposited with the American Type Culture Collection under accession no HB 9259. Expression systems useful for producing recombinant proteins fused to the FLAG® peptide, and monoclonal antibodies that bind the peptide and are useful in purifying the recombinant proteins, are available from Eastman Kodak Company, Scientific Imaging Systems, New Haven, Connecticut.

**[0067]** Particularly suitable LIR fusion proteins are those in which an LIR polypeptide is in the form of an oligomer. Oligomers may be formed by disulfide bonds between cysteine residues on more than one LIR polypeptide, or by noncovalent interactions between LIR polypeptide chains. In another approach, LIR oligomers can be formed by joining LIR polypeptides or fragment thereof via covalent or noncovalent interactions between peptide moieties fused to the LIR polypeptide. Suitable peptide moieties include peptide linkers or spacers, or peptides that have the property of promoting oligomerization. Leucine zippers and certain polypeptides derived from antibodies are among the peptides that can promote oligomerization of LIR polypeptides attached thereto.

**[0068]** Other LIR fusion proteins which promote oligomer formation are fusion proteins having heterologous polypeptides fused to various portions of antibody-derived polypeptides (including the Fc domain). Procedures for preparing such fusion proteins are described in Ashkenazi et al. *PNAS USA* 88:10535, 1991; Byrne et al. *Nature* 344:667, 1990, and Hollenbaugh and Aruffo *Current Protocols in Immunology*, Supplement 4, pages 10.19.1-10.19.11, 1992; all of which are incorporated herein by reference. Example 1 and Example 5 below describe methods for preparing UL18:Fc and P3G2:Fc fusion proteins, respectively, by fusing P3G2 and UL18 to an Fc region polypeptide derived from an antibody. This is accomplished by inserting into an expression vector a gene fusion encoding the P3G2:Fc fusion protein and expressing the P3G2:Fc fusion protein. The fusion proteins are allowed to assemble much like antibody molecules, whereupon interchain disulfide bonds form between the Fc polypeptides, yielding divalent P3G2 polypeptide. In a similar approach, P3G2 or any LIR polypeptide may be substituted for the variable portion of an antibody heavy or light chain. If fusion proteins are made with heavy and light chains of an antibody, it is possible to form a LIR oligomer with as many as four LIR regions.

**[0069]** Thus, the invention encompasses nucleic acids that encode fusion proteins that include the Fc region of Ig and an amino acid sequence including the extracellular region of any of the LIR family member proteins. Such extracellular regions include, e.g., amino acids  $x_1$ -458 of SEQ ID NO:2, where  $x_1$  is amino acids 1 or 17; amino acids  $x_2$ -459 of SEQ ID NO:4, where  $x_2$  is amino acid 1 or 17; amino acids  $x_3$ -439 of SEQ ID NO:8, where  $x_3$  is amino acid 1 or 17; amino acids  $x_4$ -458 of SEQ ID NO:10, where  $x_4$  is amino acid 1 or 17; amino acids  $x_5$  to 261 of SEQ ID NO:12, wherein  $x_5$  is amino acid 1 or 17; amino acids  $x_6$  to 461 of SEQ ID NO:14, wherein  $x_6$  is amino acid 1 or 17; amino acids  $x_7$ -449 of SEQ ID NO:16, where  $x_7$  is amino acid 1 or 17; amino acids  $x_8$ -259 of SEQ ID NO:18, where  $x_8$  is amino acid 1 or 17; amino acids  $x_9$ -443 of SEQ ID NO:20, where  $x_9$  is amino acid 1 or 17; amino acids  $x_{10}$  to 456 of SEQ ID NO:22, wherein  $x_{10}$  is amino acid 1 or 17; amino acids  $x_{11}$  to 262 of SEQ ID NO:30, wherein  $x_{11}$  is amino acid 1 or 35; amino acids  $x_{12}$  to 250 of SEQ ID NO:32, wherein  $x_{12}$  is amino acid 1 or 36; amino acids  $x_{13}$  to 265 of SEQ ID NO:34, wherein  $x_{13}$  is amino acid 1 or 35; and amino acids  $x_{14}$  to 253 of SEQ ID NO:36, wherein  $x_{14}$  is amino acid 1 or 36; and amino acids 1-393 of SEQ ID NO:38.

**[0070]** As used herein, a Fc polypeptide includes native and mutein forms, as well as truncated Fc polypeptides containing the hinge region that promotes dimerization. One suitable Fc polypeptide is the native Fc region polypeptide derived from a human IgG1, which is described in PCT application WO 93/10151, hereby incorporated herein by reference. Another useful Fc polypeptide is the Fc mutein described in U.S. Patent 5,457,035. The amino acid sequence of the mutein is identical to that of the native Fc sequence presented in WO 93/10151, except that amino acid 19 has been changed from Leu to Ala, amino acid 20 has been changed from Leu to Glu, and amino acid 22 has been changed from Gly to Ala. This mutein Fc exhibits reduced affinity for immunoglobulin receptors.

**[0071]** Alternatively, oligomeric LIR polypeptide variants may include two or more LIR peptides joined through peptide linkers. Examples include those peptide linkers described in U.S. Patent No. 5,073,627, incorporated herein by reference. Fusion proteins which include multiple LIR polypeptides separated by peptide linkers may be produced conventional recombinant DNA technology.

**[0072]** Another method for preparing oligomeric LIR polypeptide variants involves use of a leucine zipper. Leucine zipper domains are peptides that promote oligomerization of the proteins in which they are found. Leucine zippers were first identified in several DNA-binding proteins (Landschulz et al. *Science* 240:1759, 1988). Among the known leucine zippers are naturally occurring peptides and peptide derivatives that dimerize or trimerize. Examples of leucine zipper domains suitable for producing soluble oligomeric LIR polypeptides or oligomeric polypeptides of the LIR family are those described in PCT application WO 94/10308, incorporated herein by reference. Recombinant fusion proteins having a soluble LIR polypeptide fused to a peptide that dimerizes or trimerizes in solution may be expressed in suitable host cells, and the resulting soluble oligomeric LIR polypeptide recovered from the culture supernatant.

**[0073]** Numerous reagents useful for cross-linking one protein molecule to another are known. Heterobifunctional and homobifunctional linkers are available for this purpose from Pierce Chemical Company, Rockford, Illinois, for ex-

ample. Such linkers contain two functional groups (e.g., esters and/or maleimides) that will react with certain functional groups on amino acid side chains, thus linking one polypeptide to another.

[0074] One type of peptide linker that may be employed in the present invention separates polypeptide domains by a distance sufficient to ensure that each domain properly folds into the secondary and tertiary structures necessary for the desired biological activity. The linker also should allow the extracellular portion to assume the proper spatial orientation to form the binding sites for ligands.

[0075] Suitable peptide linkers are known in the art, and may be employed according to conventional techniques. Among the suitable peptide linkers are those described in U.S. Patents 4,751,180 and 4,935,233, which are hereby incorporated by reference. A peptide linker may be attached to LIR polypeptides by any of the conventional procedures used to attach one polypeptide to another. The cross-linking reagents available from Pierce Chemical Company as described above are among those that may be employed. Amino acids having side chains reactive with such reagents may be included in the peptide linker, e.g., at the termini thereof. Preferably, a fusion proteins formed via a peptide linker are prepared by recombinant DNA technology.

[0076] The fusion proteins of the present invention include constructs in which the C-terminal portion of one protein is fused to the linker which is fused to the N-terminal portion of another protein. Peptides linked in such a manner produce a single protein which retains the desired biological activities. The components of the fusion protein are listed in their order of occurrence (i.e., the N-terminal polypeptide is listed first, followed by the linker and then the C-terminal polypeptide).

[0077] A DNA sequence encoding a fusion protein is constructed using recombinant DNA techniques to insert separate DNA fragments encoding the desired proteins into an appropriate expression vector. The 3' end of a DNA fragment encoding one protein is ligated (*via* the linker) to the 5' end of the DNA fragment encoding another protein with the reading frames of the sequences in phase to permit translation of the mRNA into a single biologically active fusion protein. A DNA sequence encoding an N-terminal signal sequence may be retained on the DNA sequence encoding the N-terminal polypeptide, while stop codons, which would prevent read-through to the second (C-terminal) DNA sequence, are eliminated. Conversely, a stop codon required to end translation is retained on the second DNA sequence. DNA encoding a signal sequence is preferably removed from the DNA sequence encoding the C-terminal polypeptide.

[0078] A DNA sequence encoding a desired polypeptide linker may be inserted between, and in the same reading frame as, the DNA sequences encoding the two proteins using any suitable conventional technique. For example, a chemically synthesized oligonucleotide encoding the linker and containing appropriate restriction endonuclease cleavage sites may be ligated between the sequences encoding Fc and a P3G2 polypeptide.

[0079] Within the scope of the present invention are recombinant expression vectors for expressing polypeptides of the LIR family, and host cells transformed with the expression vectors. Expression vectors of the invention include DNA that encodes a LIR family member operably linked to suitable transcriptional or translational regulatory nucleotide sequences, such as those derived from a mammalian, microbial, viral, or insect gene. Examples of regulatory sequences include transcriptional promoters, operators, or enhancers, an mRNA ribosomal binding site, and appropriate sequences which control transcription and translation initiation and termination. Nucleotide sequences are operably linked when the regulatory sequence functionally relates to the LIR DNA sequence. Thus, a promoter nucleotide sequence is operably linked to a LIR DNA sequence if the promoter nucleotide sequence controls the transcription of the LIR DNA sequence. An origin of replication that confers the ability to replicate in the desired host cells, and a selection gene by which transformants are identified, are generally incorporated in the expression vector.

[0080] In addition, a sequence encoding an appropriate signal peptide can be incorporated into expression vectors. A DNA sequence for a signal peptide (secretory leader) may be fused in frame to the LIR sequence so that the LIR is initially translated as a fusion protein comprising the signal peptide. A signal peptide that is functional in the intended host cells promotes extracellular secretion of the LIR polypeptide. The signal peptide is cleaved from the LIR polypeptide upon secretion of the LIR polypeptide from the cell.

[0081] The recombinant expression vectors of the present invention may include any DNA encoding a LIR polypeptide. Exemplary DNAs for inclusion in such expression vectors include the nucleic acid molecules whose sequences are shown in SEQ ID NOS:1, 3, 7, 9, 11, 13, 15, 17, 19, 21, 29, 31, 33, 35 and 37.

[0082] Suitable host cells for expression of LIR polypeptides include prokaryotes, yeast or higher eukaryotic cells. Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are described, for example, in Pouwels et al. *Cloning Vectors: A Laboratory Manual*, Elsevier, New York, (1985). Cell-free translation systems could also be employed to produce P3G2 polypeptides using RNAs derived from DNA constructs disclosed herein.

[0083] Prokaryote host cells suitable in the practice of the present invention include gram negative or gram positive organisms, for example, *E. coli* or *Bacilli*. Suitable prokaryotic host cells for transformation include, for example, *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium*, and various other species such as *Pseudomonas*, *Streptomyces*, and *Staphylococcus*. In a prokaryotic host cell, such as *E. coli*, a P3G2 polypeptide may include an N-terminal methionine

residue to facilitate expression of the recombinant polypeptide. The N-terminal Met may be cleaved from the expressed recombinant LIR polypeptide.

**[0084]** Expression vectors for use in prokaryotic host cells generally include one or more phenotypic selectable marker genes. A phenotypic selectable marker gene is, for example, a gene encoding a protein that confers antibiotic resistance or that supplies an autotrophic requirement. Examples of useful expression vectors for prokaryotic host cells include those derived from commercially available plasmids such as the cloning vector pBR322 (ATCC 37017). pBR322 contains genes for ampicillin and tetracycline resistance and thus provides simple means for identifying transformed cells. An appropriate promoter and a LIR family DNA may be inserted into the pBR322 vector. Other commercially available vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and pGEM1 (Promega Biotec, Madison, WI, USA).

**[0085]** Promoter sequences commonly used for recombinant prokaryotic host cell expression vectors include  $\beta$ -lactamase (penicillinase), lactose promoter system (Chang et al. *Nature* 75:615, 1978; and Goeddel et al., *Nature* 281:544, 1979), tryptophan (trp) promoter system (Goeddel et al., *Nucl. Acids Res.* 8:4057, 1980); and EP-A-36776) and tac promoter (Maniatis, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, p. 412, 1982). A particularly useful prokaryotic host cell expression system employs a phase  $\lambda$ P<sub>L</sub> promoter and a cl857ts thermolabile repressor sequence. Plasmid vectors available from the American Type Culture Collection which incorporate derivatives of the  $\lambda$ P<sub>L</sub> promoter include plasmid pHUB2 (resident in *E. coli* strain JMB9, ATCC 37092) and pPLc28 (resident in *E. coli* RR1, ATCC 53082).

**[0086]** Alternatively, LIR polypeptides may be expressed in yeast host cells, preferably from the *Saccharomyces* genus (e.g., *S. cerevisiae*). Other genera of yeast, such as *Pichia* or *Kluyveromyces* may also be employed. Yeast vectors will often contain an origin of replication sequence from a 2 $\mu$  yeast plasmid, an autonomously replicating sequence (ARS), a promoter region, sequences for polyadenylation, sequences for transcription termination, and a selectable marker gene. Suitable promoter sequences for yeast vectors include, among others, promoters for metallothionein, 3-phosphoglycerate kinase (Hitzeman et al., *J. Biol. Chem.* 255:2073, 1980) or other glycolytic enzymes (Hess et al., *J. Adv. Enzyme Reg.* 7:149, 1968); and Holland et al., *Biochem.* 17:4900, 1978), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phospho-glucose isomerase, and glucokinase. Other suitable vectors and promoters for use in yeast expression are further described in Hitzeman, EPA-73,675. Another alternative is the glucose-repressible ADH2 promoter described by Russell et al. (*J. Biol. Chem.* 258:2674, 1982) and Beier et al. (*Nature* 300:724, 1982). Shuttle vectors replicable in both yeast and *E. coli* may be constructed by inserting DNA from pBR322 for selection and replication in *E. coli* (Amp<sup>r</sup> gene and origin of replication) into the above-described yeast vectors.

**[0087]** The yeast  $\alpha$ -factor leader sequence may be employed to direct secretion of the LIR polypeptide. The  $\alpha$ -factor leader sequence is often inserted between the promoter sequence and the structural gene sequence. See, e.g., Kurjan et al., *Cell* 30:933, 1982 and Bitter et al., *Proc. Natl. Acad. Sci. USA* 81:5330, 1984. Other leader sequences suitable for facilitating secretion of recombinant polypeptides from yeast hosts are known to those of skill in the art. A leader sequence may be modified near its 3' end to contain one or more restriction sites. This will facilitate fusion of the leader sequence to the structural gene.

**[0088]** Yeast transformation protocols are known to those of skill in the art. One such protocol is described by Hinnen et al., *Proc. Natl. Acad. Sci. USA* 75:1929, 1978. The Hinnen et al. protocol selects for Trp<sup>+</sup> transformants in a selective medium, wherein the selective medium consists of 0.67% yeast nitrogen base, 0.5% casamino acids, 2% glucose, 10  $\mu$ g/mL adenine and 20  $\mu$ g/mL uracil.

**[0089]** Yeast host cells transformed by vectors containing an ADH2 promoter sequence may be grown for inducing expression in a "rich" medium. An example of a rich medium is one having 1% yeast extract, 2% peptone, and 1% glucose supplemented with 80  $\mu$ g/mL uracil. Derepression of the ADH2 promoter occurs when glucose is exhausted from the medium.

**[0090]** Mammalian or insect host cell culture systems may be used to express recombinant LIR polypeptides. Baculovirus systems for production of heterologous proteins in insect cells are reviewed by Luckow and Summers, *Bio/Technology* 6:47 (1988). Established cell lines of mammalian origin also may be employed. Examples of suitable mammalian host cell lines include the COS-7 line of monkey kidney cells (ATCC CRL 1651) (Gluzman et al., *Cell* 23:175, 1981), L cells, C127 cells, 3T3 cells (ATCC CCL 163), Chinese hamster ovary (CHO) cells, HeLa cells, and BHK (ATCC CRL 10) cell lines, and the CV1/EBNA cell line derived from the African green monkey cell line CV1 (ATCC CCL 70) as described by McMahan et al. (*EMBO J.* 10:2821, 1991). COS-1 (ATCC CRL-1650).

**[0091]** Transcriptional and translational control sequences for mammalian host cell expression vectors may be excised from viral genomes. Commonly used promoter sequences and enhancer sequences are derived from Polyoma virus, Adenovirus 2, Simian Virus 40 (SV40), and human cytomegalovirus. DNA derived from the SV40 viral genome, for example, SV40 origin, early and late promoter, enhancer, splice, and polyadenylation sites may be used to provide other genetic elements for expression of a structural gene sequence in a mammalian host cell. Viral early and late



promoters are particularly useful because both are easily obtained from a viral genome as a fragment which may also contain a viral origin of replication (Fiers et al., *Nature* 273:113, 1978). Smaller or larger SV40 fragments may also be used, provided the approximately 250 bp sequence extending from the HIND III site toward the Bgl/I site located in the SV40 viral origin of replication site is included.

5 [0092] Suitable expression vectors for use in mammalian host cells can be constructed as disclosed by Okayama and Berg (*Mol. Cell. Biol.* 3:280, 1983). One useful system for stable high level expression of mammalian receptor cDNAs in C127 murine mammary epithelial cells can be constructed substantially as described by Cosman et al. (*Mol. Immunol.* 23:935, 1986). A high expression vector, PMLSV N1/N4, described by Cosman et al., *Nature* 312:768, 1984 has been deposited as ATCC 39890. Additional mammalian expression vectors are described in EP-A-0367566, and  
10 in WO 91/18982. Still additional expression vectors for use in mammalian host cells include pDC201 (Sims et al., *Science* 241:585, 1988), pDC302 (Mosley et al. *Cell* 59:335, 1989), and pDC406 (McMahan et al., *EMBO J.* 10:2821, 1991). Vectors derived from retroviruses also may be employed. One preferred expression system employs pDC409 as discussed in Example 5 below.

15 [0093] For expression of LIR polypeptides the expression vector may comprise DNA encoding a signal or leader peptide. In place of the native signal sequence, a heterologous signal sequence may be added, such as the signal sequence for interleukin-7 (IL-7) described in United States Patent 4,965,195; the signal sequence for interleukin-2 receptor described in Cosman et al., *Nature* 312:768, 1984; the interleukin-4 signal peptide described in EP 367,566; the type I interleukin-1 receptor signal peptide described in U.S. Patent 4,968,607; and the type II interleukin-1 receptor signal peptide described in EP 460,846.

20 [0094] Further contemplated within the present invention are purified LIR family polypeptides, and processes for their purification. The purified polypeptides of the present invention may be purified from the above-described recombinant expression systems or may be purified from naturally occurring cells. The desired degree of purity may depend on the intended use of the protein with a relatively high degree of purity preferred when the protein is intended for *in vivo* use. Preferably, LIR polypeptide purification processes are such that no protein bands corresponding to proteins  
25 other than the desired LIR protein are detectable by SDS-polyacrylamide gel electrophoresis (SDS-PAGE). It will be recognized by one skilled in the art that multiple bands corresponding to any LIR polypeptide may be detected by SDS-PAGE, due to differential glycosylation, variations in post-translational processing, and the like, as discussed above. Most preferably, any specific LIR polypeptide is purified to substantial homogeneity, as indicated by a single protein band upon analysis by SDS-PAGE. The protein band may be visualized by silver staining, Coomassie blue  
30 staining, or by autoradiography or fluorescence if the protein is appropriately labeled.

[0095] One process for providing purified LIR polypeptides includes first culturing a host cell transformed with an expression vector comprising a DNA sequence that encodes the desired polypeptide under conditions that promote expressing the desired LIR polypeptide and then recovering the LIR polypeptide. As the skilled artisan will recognize, procedures for recovering the polypeptide will vary according to such factors as the type of host cells employed and  
35 whether the polypeptide is secreted in the culture medium is extracted from cells.

[0096] When the expression system secretes the polypeptide into the culture medium, the medium may be first concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. Following the concentration step, the concentrate can be applied to a suitable purification matrix such as a gel filtration medium. Alternatively, an anion exchange resin can be employed, such as a resin matrix or  
40 resin substrate having pendant diethylaminoethyl (DEAE) groups. The matrices can be acrylamide, agarose, dextran, cellulose or other types commonly employed in protein purification. Similarly, a purification matrix having cation exchange groups such as sulfopropyl or carboxymethyl functionalities on an insoluble matrix can be used. Sulfopropyl groups are preferred. Still other purification matrices and methods suitable for providing purified LIR are high performance liquid chromatography using hydrophobic reversed phase media (RP-HPLC). One skilled in the art will recognized  
45 the any or all of the foregoing purification steps, in various combinations, can be employed to provide a purified LIR polypeptide.

[0097] Alternatively, LIR polypeptides can be purified by immunoaffinity chromatography. An affinity column containing an antibody that binds a LIR polypeptide may be prepared by conventional procedures and employed in purifying LIR. Example 5 describes a procedures for generating monoclonal antibodies directed against P3G2 which may be  
50 utilized in immunoaffinity chromatography.

[0098] Recombinant protein produced in bacterial culture may be isolated by first disrupting the host cells by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents and then extracting the polypeptide from cell pellets if the polypeptide is insoluble, or from the supernatant fluid if the polypeptide is soluble. After the initial isolation step, the purification process may include one or more concentrating,  
55 salting out, ion exchange, affinity, or size exclusion chromatography purification steps. For many application a final RP-HPLC purification step is beneficial.

[0099] Additional methods for providing LIR polypeptides and purified LIR polypeptides involves fermenting yeast which express proteins as a secreted protein. Secreted recombinant protein resulting from a large-scale fermentation



can be purified by methods analogous to those disclosed by Urdal et al. (*J. Chromatog.* 296:171, 1984), involving two sequential, reversed-phase HPLC steps for purification of a recombinant protein on a preparative HPLC column.

**[0100]** LIR-P3G2 DNA in pDC406 vector was deposited with the American Type Culture Collection on April 22, 1997 and assigned accession No. 97995. The deposit was made under the terms of the Budapest Treaty.

**[0101]** As described above and shown in Examples 6 and 14, LIR-P3G2 and LIR-pbm8 are MHC class I receptor molecules found on the surface of certain monocytes, B cells, and NK cells. With respect to monocytes, the expression of LIRs that are MHC class I binding proteins suggests that there is some requirement for monocytes to recognize MHC class I molecules. LIR-P3G2, LIR-pbm8 LIR and certain additional LIR family members contain cytoplasmic ITIM motifs. By analogy with the structure and function of known MHC class I receptor molecules, these LIRs are inhibitory receptors mediating negative signaling. Indeed, the results demonstrated in Example 11 reveal that LIRs associate with SHP-1 and inhibit FcR-mediated activation events. Thus, monocytes may express class I receptors in order to suppress cell-mediated lytic mechanisms. Monocytes rapidly phagocytose extracellular pathogens via FcR and, monocyte-FcR engagement induces propagation of immune responses by producing more systemic mediators, particularly TNF- $\alpha$ , IL-6 and IL-8. Thus, the LIRs play a role in monocyte and macrophage regulation of cytolytic and inflammatory responses against self tissues. The interplay between the FcR activatory signals and LIR inhibitory signal may allow low levels of self-reactive IgG to exist in circulation and bind to the monocyte membrane with initiating an immune response. For example, the expression of these inhibitory receptors can protect the developing embryo from maternal antibody-mediated allogeneic recognition.

**[0102]** With respect to LIRs on cells of the DC lineage, as described in Example 13 CD33<sup>+</sup>CD14<sup>+</sup>CD16<sup>+</sup>HLA-DR<sup>+</sup> DC co-express LIR-P3G2 and LIR-pbm8. It is suggested the DC FcR play a role in binding immune complexes and triggering DC activation signal following binding. Thus, LIRs expressed on DC may suppress DC activation through interactions of FcR.

**[0103]** Many LIR family members lack the ITIM motif and by analogy with the structure and function of known MHC class I receptors lacking ITIMs are activatory receptors. Failure of a receptor that mediates negative signaling could result in autoimmune diseases. Thus, engaging an LIR family member having ITIM motifs with an agonistic antibody or ligand can be used to downregulate a cell function in disease states in which the immune system is overactive and excessive inflammation or immunopathology is present. On the other hand, using an antagonistic antibody specific to the ITIM possessing LIR receptor or a soluble form of the receptor can be used to block the interaction of the cell surface receptor with the receptor's ligand to activate the specific immune function in disease states associated with suppressed immune function. Since receptors lacking the ITIM motif send activatory signals once engaged as described above, failure of a receptor that mediates an activatory signal could result in suppressed immune function. Engaging the receptor with its agonistic antibody or ligand can be used to treat diseases associated with the suppressed immune function. Using an antagonistic antibody specific to the activatory LIR receptor or a soluble form of the receptor can be used to block the interaction of the activatory receptor with the receptor's ligand to downregulate the activatory signaling.

**[0104]** Since LIR-P3G2 binds to various cells, LIR-P3G2 may be used to purify or isolate these cells from heterogeneous preparations. Additionally, P3G2 probes can be used to isolate and identify related molecules.

**[0105]** LIR polypeptides of the present invention may be used in developing treatments for any disorder mediated directly or indirectly by defective or insufficient amounts of any of the LIR polypeptides. A therapeutically effective amount of purified LIR protein is administered by a patient afflicted with such a disorder. Alternatively, LIR DNA may be employed in developing a gene therapy approach to treating such disorders. Disclosure herein of native LIR nucleotide sequence permits the detection of defective LIR genes, and the replacement thereof with normal LIR-encoding genes. Defective genes may be detected in *in vitro* diagnostic assays, and by comparison of the native LIR nucleotide sequence disclosed herein with that of an LIR gene derived from a person suspected of harboring a defect in the gene.

**[0106]** The present invention also provides pharmaceutical compositions which may include an LIR polypeptide, or fragments or variants thereof with a physiologically acceptable carrier or diluent. Such carriers and diluents will be nontoxic to recipients at the dosages and concentrations employed. Such compositions may further include buffers, antioxidants such as ascorbic acid, low molecular weight (less than about ten residues) polypeptides, proteins, amino acids, carbohydrates including glucose, sucrose or dextrans, chelating agents such as EDTA, glutathione and other stabilizers and excipients commonly used in pharmaceutical compositions. The pharmaceutical compositions of the present invention may be formulated as a lyophilizate using appropriate excipient solutions as diluents. The pharmaceutical compositions may include an LIR polypeptide in any form described herein, including but not limited to active variants, fragments, and oligomers. LIR polypeptides may be formulated according to known methods that are used to prepare pharmaceutically useful compositions. Components that are commonly employed in pharmaceutical formulations include those described in *Remington's Pharmaceutical Sciences*, 16th ed. (Mack Publishing Company, Easton, PA, 1980).

**[0107]** The pharmaceutical preparations of the present invention may be administered to a patient, preferably a human, in a manner appropriate to the indication. Thus, for example, the compositions can be administered by intra-

venous injection, local administration, continuous infusion, sustained release from implants, etc. Appropriate dosages and the frequency of administration will depend on such factors as the nature and severity of the indication being treated, the desired response, the condition of the patient and so forth.

**[0108]** In preferred embodiments an LIR polypeptide used in the pharmaceutical compositions of the present invention is purified such that the LIR polypeptide is substantially free of other proteins of natural or endogenous origin, desirably containing less than about 1% by mass of protein contaminants residual of the production processes. Such compositions, however, can contain other proteins added as stabilizers, carriers, excipients or co-therapeutics.

**[0109]** LIR encoding DNAs and DNA fragments disclosed herein find use in the production of LIR polypeptides, as described above. In one embodiment, such fragments comprise at least about 17 consecutive nucleotides, more preferably at least 30 consecutive nucleotides, of LIR DNA. DNA and RNA complements of the fragments have similar utility. Among the uses of LIR nucleic acid fragments are as probes or primers in polymerase chain reactions. For example, a probe corresponding to a fragment of DNA encoding the extracellular domain of LIR may be employed to detect the presence of LIR nucleic acids in *in vitro* assays and in other probing assays such as Northern Blot and Southern blot assays. Cell types expressing an LIR polypeptide can be identified using LIR family nucleic acid probes using probing procedures well known in the art. Those skilled in the art have the knowledge to choose a probe of suitable length and apply conventional PCR techniques to isolate and amplify a DNA sequence.

**[0110]** Nucleic acid fragments may also be used as a probe in cross species hybridization procedures to isolate LIR DNA from other mammalian species. As one example, a probe corresponding to the extracellular domain of an LIR polypeptide may be employed. The probes may be labeled (e.g., with  $^{32}\text{P}$ ) by conventional techniques.

**[0111]** Other useful fragments of LIR nucleic acids are sense or antisense oligonucleotides, which may comprise either RNA or DNA, and which correspond in sequence to an LIR mRNA (sense), to the complement of an LIR mRNA (antisense), or to the non-coding strand of a double-stranded LIR DNA, such as P3G2 DNA (antisense). Thus, an antisense oligonucleotide will form a hybrid duplex with an mRNA sequence. Such oligonucleotides generally are at least 14 nucleotides, and preferably are from about 14 to about 30 nucleotides. The ability to create an antisense or a sense oligonucleotide based upon a cDNA sequence for a given protein is described in, for example, Stein and Cohen, *Cancer Res.* 48:2659, 1988 and van der Krol et al., *BioTechniques* 6:958, 1988.

**[0112]** Binding antisense or sense oligonucleotides to target nucleic acid sequences results in the formation of duplexes that block translation (RNA) or transcription (DNA) by one of several means, including enhanced degradation of the duplexes, premature termination of transcription or translation, or by other means. These oligonucleotides thus may be used to block LIR expression.

**[0113]** In one embodiment antisense or sense LIR oligonucleotides used in binding procedures may encompass oligonucleotides having modified sugar-phosphodiester backbones (or other sugar linkages, such as those described in WO91/06629) and wherein such sugar linkages are resistant to endogenous nucleases. Oligonucleotides having sugar linkages resistant to endogenous nucleases are stable *in vivo* (i.e., capable of resisting enzymatic degradation) but retain sequence specificity to be able to bind to target nucleotide sequences. Other examples of sense or antisense oligonucleotides include those oligonucleotides which are covalently linked to organic moieties, such as those described in WO 90/10448, and other moieties that increase affinity of the oligonucleotide for a target nucleic acid sequence, such as poly-(L-lysine). Further still, intercalating agents, such as ellipticine, and alkylating agents or metal complexes may be attached to sense or antisense oligonucleotides to modify binding specificities of the antisense or sense oligonucleotide for the target nucleotide sequence.

**[0114]** Antisense or sense oligonucleotides may be introduced into a cell containing the target nucleic acid sequence by any gene transfer method, including, for example,  $\text{CaPO}_4$ -mediated DNA transfection, electroporation, or by using gene transfer vectors such as Epstein-Barr virus. Antisense or sense oligonucleotides are preferably introduced into a cell containing the target nucleic acid sequence by inserting the antisense or sense oligonucleotide into a suitable retroviral vector, then contacting the cell with the retroviral vector containing the inserted sequence, either *in vivo* or *ex vivo*. Suitable retroviral vectors include, but are not limited to, those derived from the murine retrovirus M-MuLV, N2 (a retrovirus derived from M-MuLV), or the double copy vectors designated DCT5A, DCT5B and DCT5C (see PCT Application US 90/02656).

**[0115]** Sense or antisense oligonucleotides also may be introduced into a cell containing the target nucleotide sequence by formation of a conjugate with a ligand binding molecule, as described in WO 91/04753. Suitable ligand binding molecules include, but are not limited to, cell surface receptors, growth factors, other cytokines, or other ligands that bind to cell surface receptors. Preferably, conjugating the ligand binding molecule does not substantially interfere with the ability of the ligand binding molecule to bind its corresponding molecule or receptor, or block entry of the sense or antisense oligonucleotide or its conjugated version into the cell.

**[0116]** Alternatively, a sense or an antisense oligonucleotide may be introduced into a cell containing the target nucleic acid sequence by formation of an oligonucleotide-lipid complex, as described in WO 90/10448. The sense or antisense oligonucleotide-lipid complex is preferably dissociated within the cell by an endogenous lipase.

**[0117]** In still a further aspect, the present invention provides antibodies that specifically bind LIR polypeptides, i.e.,

antibodies bind to LIR polypeptides *via* an antigen-binding site of the antibody (as opposed to non-specific binding). Antibodies of the present invention may be generated using LIR polypeptides or immunogenic fragments thereof. Polyclonal and monoclonal antibodies may be prepared by conventional techniques. See, for example, Monoclonal Antibodies, Hybridomas: A New Dimension in Biological Analyses, Kennet et al. (eds.), Plenum Press, New York 1980; and *Antibodies: A Laboratory Manual*, Harlow and Land (eds.), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1988. An exemplary procedure for producing monoclonal antibodies immunoreactive with P3G2-LIR is further illustrated in Example 5 below.

[0118] Included within the scope of the present invention are antigen binding fragments of antibodies which specifically bind to an LIR polypeptide. Such fragments include, but are not limited to, Fab, F(ab')<sub>2</sub>, and F(ab')<sub>2</sub>. Antibody variants and derivatives produced by genetic engineering techniques are contemplated as within the presented invention.

[0119] The monoclonal antibodies of the present invention include chimeric antibodies, e.g., humanized versions of murine monoclonal antibodies. Such antibodies may be prepared by known techniques and offer the advantage of reduced immunogenicity when the antibodies are administered to humans. In one embodiment a humanized monoclonal antibody comprises the variable region of a murine antibody (or just the antigen binding site thereof) and a constant region derived from a human antibody. Alternatively, a humanized antibody fragment may comprise the antigen binding site of a murine monoclonal antibody and a variable region fragment (lacking the antigen-binding site) derived from a human antibody. Procedures for the production of chimeric and further engineered monoclonal antibodies include those described in Riechmann et al., *Nature* 332:232, 1988; Lie et al. *PNAS* 84:3439, 1987; Larrick et al. *Biol Technology* 7:934, 1989; and Winter and Hams *TIPS* 14:139, 1993.

[0120] As mentioned above, antibodies of the present invention are useful in *in vitro* or *in vivo* assays to detect the presence of LIR polypeptides and in purifying an LIR polypeptide by affinity chromatography.

[0121] Additionally, antibodies capable of blocking an LIR from binding to target cells may be used to inhibit a biological activity of an LIR polypeptide. More specifically, therapeutic compositions of an antibody antagonistic to one or more LIR family members having the ITIM motif may be administered to an individual in order to block the interaction of a cell surface LIR with its ligand. The result is an activation of immune function and is particularly beneficial in disease states in which the immune system is hyporesponsive or suppressed. Conversely, therapeutic compositions of an antibody antagonistic to one or more LIR family members lacking the ITIM motif may be used to obtain the opposite effect and be beneficial in disease states in which the immune system is overactive and excessive inflammation or immunopathology is present.

[0122] Pharmaceutical compositions which include at least one antibody that is immunoreactive with an LIR polypeptide and a suitable diluent, excipient, or carrier, are considered with the present invention. Suitable diluents, excipients, and carriers are described in the context of pharmaceutical compositions which include polypeptides of the present invention.

[0123] The following examples are provided to illustrate certain embodiments of the invention, and are not to be construed as limiting the scope of the invention.

## EXAMPLES

### Example 1. Isolating and Expressing Viral Protein

[0124] DNA encoding P3G2 polypeptide of the present invention was identified by isolating and expressing a viral glycoprotein, UL18, known to be expressed on cells infected with HCMV, and then expressing and using a UL18/Fc fusion protein to search for UL18 receptors. DNA encoding UL18 and its amino acid sequence are known and described in Beck, S., B.G. Barrell, *Nature* 331:269-272, 1988. The following describes isolating UL18 and preparing the UL18/Fc fusion protein.

[0125] Using standard techniques, total RNA was isolated from Human Foreskin Fibroblasts infected with HCMV (AD169) at three different transcription stages-immediate early (IE, 8 p.i.h.), early (24 p.i.h.) and late (48 p.i.h.). Because UL18 is known to be transcribed early in the infection, the IE total RNA was polyA<sup>+</sup> selected and used to construct an HCMV-IE cDNA library using a cDNA kit according to the manufacturer's instructions (Pharmacia TIME SAVER cDNA Kit). In order to isolate the full length UL18 gene, two oligonucleotide primers known to include the terminal sequences of the UL18 gene were synthesized and used to isolate and amplify the UL18 gene from the HCMV-IE cDNA library. The primers had the following sequences and included *Not* I restriction sites which incorporate into the PCR product:

Not I  
5' - TAT GCG GCC GCC ATG ATG ACA ATG TGG T - 3' (SEQ ID NO:23)

5' - TAT GCG GCC GCC CCT TGC GAT AGC G - 3' (SEQ ID NO:24)

Not I

The PCR conditions included one 5 minute 95° C cycle followed by 30 cycles of 45 seconds at 95°, 45 seconds at 58° and 45 seconds at 72°, and then one cycle for 5 minutes at 72°C. The PCR product was electrophoresed on a 1% agarose gel and sized using ethidium bromide to visualize the separated DNA products. The presence of DNA of having the expected size of approximately 1.1kb was confirmed.

[0126] The pDC409 expression vector, a vector derived from pDC406 (McMahan et al., *EMBO J.* 10:2821, 1991) but having a single *Bgl* II site was selected for the cloning process. The PCR product was subcloned into a pDC409 expression vector through the *Not* I sites, sequenced and the amino acid sequence deduced from the DNA sequence. The determined nucleotide sequence and amino acid sequence were identical to the previously published sequences (ibid.).

[0127] A fusion protein of the extracellular region of UL18 and a mutein human IgG1 Fc region (UL18:Fc) was prepared by first isolating cDNA encoding the extracellular region of UL18 using primers which flank the extracellular region of UL18. The primers were synthesized with *Sal* I and *Bgl* II restriction sites inserted at the 5' and 3' termini so that the PCR amplified cDNA introduced *Sal* I and *Bgl* II restriction sites at the 5' and 3' ends, respectively. The primers had the following sequences:

5' -ATA GTC GAC AAC GCC ATG ATG ACA ATG TGG TG - 3' (SEQ ID NO:25)

*Sal* I

5' - TAA AGA TCT GGG CTC GTT AGC TGT CGG GT - 3' (SEQ ID NO:26)

*Bgl* II

The conditions for the PCR reaction were as described above except that the template was the full length gene isolated as just described.

[0128] To prepare a vector construct for expressing fusion protein, sUL18:Fc, for use in cell binding studies, a DNA fragment encoding the Fc region of a human IgG1 antibody was isolated from a plasmid using *Bgl* II and *Not* I restriction enzymes. The encoded Fc portion was the mutein Fc described in U.S. 5,457,035 having reduced affinity for immunoglobulin receptors. The *Bgl* II site on the sUL18 gene was used to ligate the sUL18 gene DNA to the *Bgl* II site on the Fc gene to form a sUL18:Fc fusion DNA construction having an N-terminal *Sal* I restriction site and a C-terminal *Not* I restriction site. This fusion sUL18:Fc DNA construct was then ligated into pDC409 expression vector at its *Sal* I and *Not* I sites to form a 409/sUL18/Fc DNA construct.

[0129] The monkey cell line COS-1 (ATCC CRL-1650) was used to confirm expression of the fusion protein. COS-1 cells in 6-well plates (2 X 10<sup>5</sup> cells per well) were transfected with about 2µg of the DNA construct 409/sUL18/Fc per well. The cells were cultured for 2-3 days in 5% FBSMEM/F12 (available from GIBCO), then washed twice with PBS, starved for 1 hour in cysteine/methionine depleted RPMI (available from GIBCO as RPMI 1640) and metabolically labeled with 100 µCi/mL of <sup>35</sup>S-Met/Cys for 4 hours. The supernatant was spun clear to remove loose cells and 150 µL of the supernatant was incubated with 100 µL of RIPA (0.05% Tween 20, 0.1% SDS, 1% Triton X-100, 0.5% deoxycholate in PBS) buffer and 50µL of 50% Protein A-Sepharose solid support beads at 4°C for 1 hour. Protein A-Sepharose is a Sepharose solid support (available from Pharmacia) having immobilized Protein A which binds the Fc portion of the fusion protein. After washing the solid support with RIPA to remove unbound material, fusion protein bound to the Protein A-Sepharose solid support was eluted from the Protein A-Sepharose using 35µL of SDS -PAGE reducing sample buffer and then heated at 100°C for 5 minutes. The eluant was then electrophoresed on a 4-20% SDS polyacrylamide gradient gel with <sup>14</sup>C labeled protein molecular weight markers. After electrophoresis the gel was fixed with 8% acetic acid and enhanced at room temperature for 20 minutes with Amplifier available from Amersham. After drying the gel under vacuum it was exposed to x-ray film. Film analysis confirmed that the expected protein, a 100-120 kDa protein which includes the mutein Fc region of IgG and UL18 extracellular domains fused to the Fc, was expressed.

[0130] Once cells expressing the fusion protein were identified large scale cultures of transfected cells were grown to accumulate supernatant from cells expressing the fusion protein. This procedure involved transfecting COS-1 cells in T175 flasks with 15 µg of the UL18/Fc/409 fusion DNA per flask. After 7 days of culture in medium containing 0.5% low immunoglobulin bovine serum, a solution of 0.2% azide was added to the supernatant and the supernatant was

filtered through a 0.22  $\mu$ m filter. Then approximately 1 L of culture supernatant was passed through a BioCad Protein A HPLC protein purification system using a 4.6 x 100 mm Protein A column (POROS 20A from PerSeptive Biosystems) at 10 mL/min. The Protein A column binds the Fc portion of the sUL18/Fc fusion protein in the supernatant, immobilizing the fusion protein and allowing other components of the supernatant to pass through the column. The column was washed with 30 mL of PBS solution and bound sUL18/Fc was eluted from the HPLC column with citric acid adjusted to pH 3.0. Eluted purified sUL18/Fc was neutralized as it eluted using 1M Hepes solution at pH 7.4. The pooled eluted protein was analyzed using SDS PAGE with silver staining, confirming expression of the 100-120 kDa UL18/Fc fusion protein.

## **Example 2. Screening Cell Lines for Binding to UL18**

[0131] The sUL18/Fc protein isolated as described in Example 1 was used to screen cell lines to which it binds using quantitative binding studies according to standard flow cytometry methodologies. For each cell line screened, the procedure involved incubating approximately 100,000 of the cells blocked with 2% FCS (fetal calf serum), 5% normal goat serum and 5% rabbit serum in PBS for 1 hour. Then the blocked cells were incubated with 5  $\mu$ g/mL of sUL18/Fc fusion protein in 2% FCS, 5% goat serum and 5% rabbit serum in PBS. Following the incubation the sample was washed 2 times with FACS buffer (2% FCS in PBS) and then treated with mouse anti human Fc/biotin (purchased from Jackson Research) and SAPE (streptavidin-phycoerythrin purchased from Molecular Probes). This treatment causes the anti human Fc/biotin to bind to any bound sUL18/Fc and the SAPE to bind to the anti human Fc/biotin resulting in a fluorescent identifying label on sUL18/Fc which is bound to cells. The cells were analyzed for any bound protein using fluorescent detection flow cytometry. The results indicated that UL18 binds well to B cell lines CB23, RAJI and MP-1; monocytic cell lines Thp-1 and U937; and primary B cell and primary monocytes. UL18 does not bind detectably to T cell lines nor does it bind to primary T cells.

## **Example 3. Isolating a P3G2 cDNA and Polypeptide**

[0132] The following describes screening cDNA of one of the cell lines found to bind UL18 and the isolation of a novel polypeptide expressed by the cell line. A CB23 cDNA library in the mammalian expression vector pDC406, prepared as described in U.S. Patent No. 5,350,683 (incorporated herein by reference) was obtained and plasmid DNA was isolated from pools consisting of approximately 2,000 clones per pool. The isolated DNA was transfected into CV1-EBNA cells (ATCC CRL 10478) using DEAE-dextran followed by chloroquine treatment. The CV1-EBNA cells were maintained in complete medium (Du)becco's modified Eagles' media containing 10% (v/v) fetal calf serum, 50 U/mL penicillin, 50 U/mL streptomycin, and 2 mM L-glutamine) and were plated to a density of approximately  $2 \times 10^5$  cells/well in single-well chambered slides. The slides had been pre-treated with 1 mL of a solution of 10  $\mu$ g/mL human fibronectin in PBS for 30 minutes followed by a single washing with PBS. Media was removed from adherent cells growing in a layer and replaced with 1.5 mL complete medium containing 66.6  $\mu$ M chloroquine sulfate. About 0.2 mL of a DNA solution (2  $\mu$ g DNA, 0.5 mg/mL DEAE-dextran in complete medium containing chloroquine) was added to the cells and the mixture was incubated at 37 C for about five hours. Following incubation, the media was removed and the cells were shocked by addition of complete medium containing 10% DMSO (dimethylsulfoxide) for 2.5 minutes. Shocking was followed by replacing the solution with fresh complete medium. The cells were grown in culture for two to three days to permit transient expression of the inserted DNA sequences. These conditions led to a 30% to 80% transfection frequency in surviving CV1-EBNA cells.

[0133] Each slide was incubated with 1 mL of UL18:Fc at a concentration of 1  $\mu$ g/mL in binding buffer (RPMI 1640 containing 25 mg/mL bovine serum albumin, 2 mg/mL sodium azide, 20 mM Hepes at pH 7.2, and 50 mg/mL nonfat dry milk) at room temperature for 1 hour. The incubated slides were washed with the binding buffer and then incubated with Fc specific  $^{125}$ I-mouse anti-human IgG (see Goodwin et al., *Cell* 73:447-456, 1993). This was followed by a second wash with buffer after which the slides were fixed with a 2.5% glutaraldehyde/PBS solution, washed with PBS solution and allowed to air dry. The dried slides were dipped in Kodak GTNB-2 photographic emulsion (6x dilution in water). After air drying, the slides were placed in a dark box and refrigerated. After three days the slides were developed in Kodak D19 developer, rinsed in water and fixed in Agfa G433C fixer. The fixed slides were individually examined under a microscope at 25-40x magnification. Positive cells demonstrating binding of sUL18:Fc were visualized by the presence of autoradiographic silver grains against the film background. Two positive pools were identified. Bacterial clones from each pool were titrated and plated to provide plates containing approximately 200 colonies each. Each plate was scraped to provide pooled plasmid DNA for transfection into CV1-EBNA cells and screening as described above. Following subsequent breakdowns and screenings, two positive individual colonies were obtained. The cDNA inserts of the two positive clones were 2922 and 2777 nucleotides in length as determined by automated DNA sequences. The coding regions of the two inserts, designated P3G2 and 18A3 were 1953 (nucleotides 310-2262) and 1959 (nucleotides 168-2126) nucleotides, respectively. The two cDNA clones encode proteins that are substantially similar and

probably represent different alleles of the same gene.

[0134] The cDNA sequence and encoded amino acid of P3G2 are presented in SEQ ID NO:1 and SEQ ID NO:2, respectively. The cDNA sequence and encoded amino acid of 18A3 are presented in SEQ ID NO:3 and SEQ ID NO:4, respectively. The P3G2 amino acid sequence (SEQ ID NO:2) has a predicted signal peptide of 16 amino acids (amino acids 1-16); an extracellular domain of 442 amino acids (amino acids 17-458); a transmembrane domain of 25 amino acids (amino acids 459-483) and, a cytoplasmic domain of 167 amino acids (amino acids 484-650). The extracellular domain includes four immunoglobulin-like domains. Ig-like domain I includes approximately amino acids 17-118; Ig-like domain II includes approximately amino acids 119-220; Ig-like domain III includes approximately amino acids 221-318; and Ig-like domain IV includes approximately amino acids 319-419. Significantly, the cytoplasmic domain of this polypeptide includes four ITIM motifs, each having the consensus sequence of YxxL/V. The first ITIM motif pair is found at amino acids 533-536 and 562-565 and the second pair is found at amino acids 614-617 and 644-647. The amino acid sequence of 18A3 is nearly identical having the features describes above.

[0135] The features of these encoded polypeptides are consistent with a type I transmembrane glycoprotein.

#### Example 4. Preparing P3G2 Fusion Protein

[0136] The following describes procedures used to generate a P3G2 fusion protein which was then used to identify cell lines to which it binds and finally isolate a normal cell-surface P3G2 ligand which is distinct from UL18. A fusion protein of the extracellular region of P3G2 and the mutein human Fc region (sP3G2:Fc) was prepared by first isolating cDNA encoding the extracellular region of P3G2 using primers which flank the extracellular region of P3G2. The primers were synthesized with *Sal*I and *Bgl*II restriction sites inserted at the 5' and 3' termini so that the PCR amplified cDNA introduced *Sal*I and *Bgl*II restriction sites at the 5' and 3' ends, respectively. The primers had the following sequences:

Sal I  
5' - TAT GTC GAC CAT GAC CCC CAT CCT CAC GGT - 3' (SEQ ID NO:5)  
Bgl II Xa

5' - TAT GGG CTC TGC TCC AGG AGA AGA TCT TCC TTC TAT AAC  
CCC CAG GTG CCT T (SEQ ID NO:6)

The conditions for the PCR reaction were as described above and the template was the full length gene P3G2 gene isolated as described in Example 3 above.

[0137] To prepare a vector construct for expressing fusion protein sP3G2:Fc for use in cell binding studies, the mutein human Fc region of IgG1 was cut from the plasmid described above in Example 1 using *Bgl*II and *Not*I restriction enzymes. The *Bgl*II site on the sP3G2 gene was used to ligate the sP3G2 gene DNA to the *Bgl*II site on the human mutein Fc gene to form a sP3G2/Fc fusion DNA construction having an N-terminal *Sal*I restriction site and a terminal *Not*I restriction site. This fusion sP3G2:Fc DNA construct was then ligated into pDC409 expression vector at its *Sal*I and *Not*I sites to form a 409/sP3G2/Fc DNA construct.

[0138] The monkey cell line COS-1 (ATCC CTL-1650) was used to confirm expression of the fusion protein. COS-1 cells in 6-well plates (2 x 10<sup>5</sup> cells per well) were transfected with about 2μg of the DNA construct 409/sP3G2/Fc per well. The cells were cultured in 5% FBS/DMEM/F12 (available from GIBCO) and at day two or three following transfection, the cells were starved for 1 hour in cysteine/methionine depleted RPMI and the transfected cells were metabolically labeled with 100 μCi/mL of <sup>35</sup>S-Met/Cys for 4 hours. The supernatant was spun clear to removed loose cells and debris and 150 μL of the supernatant was incubated with 100 μL of RIPA buffer and 50 μL of 50% Protein A-Sepharose solid support beads at 4°C for 1 hour. After washing the solid support with RIPA to remove unbound material, fusion protein bound to the Protein A-Sepharose solid support was eluted from the Protein A-Sepharose using 30μL of SDS - PAGE reducing sample buffer and then heated at 100°C for 5 minutes. The eluant was then electrophoresed on a 4-20% SDS polyacrylamide gradient gel with <sup>14</sup>C labeled protein molecular weight markers. After electrophoresis the gel was fixed with 8% acetic acid and enhanced at room temperature for 20 minutes with Amplifier available from Amersham. After drying the gel under vacuum it was exposed to x-ray film. Film analysis confirmed that the expected protein, having a molecular weight of 120-130 kDa, was expressed.

[0139] Once fusion protein expression was verified, large scale cultures of transfected cells were grown to accumulate supernatant from COS-1 cells expressing the fusion protein as described in Example 1 above. The P3G2/Fc fusion protein was purified according to the procedure described in Example 3 above using the BioCad system and the POROS

20A column from PerSeptive Biosystems. The pooled eluted protein was analyzed using SDS PAGE with silver staining, confirming expression.

#### **Example 5. Generating LIR-P3G2 Antibody**

**[0140]** The following example describes generating monoclonal antibody to P3G2 that was used in flow cytometry analysis to identify cells on which P3G2 is expressed. Purified P3G2/Fc fusion protein was prepared by COS-1 cell expression and affinity purification as described in Example 4. The purified protein or cells transfected with an expression vector encoding the full length protein can generate monoclonal antibodies against P3G2 using conventional techniques, for example those techniques described in U.S. Patent 4,411,993. Briefly BALB-C mice were immunized at 0, 2 and 6 weeks with 10 $\mu$ g P3G2/Fc. The primary immunization was prepared with TITERMAX adjuvant, from Vaxcell, Inc., and subsequent immunization were prepared with incomplete Freund's adjuvant (IFA). At 11 weeks, the mice were IV boosted with 3-4  $\mu$ g P3G2 in PBS. Three days after the IV boost, splenocytes were harvested and fused with an Ag8.653 myeloma fusion partner using 50% aqueous PEG 1500 solution. Hybridoma supernatants were screened by ELISA using P3G2 transfected COS-1 cells in PBS at 2 X 10<sup>3</sup> cells per well and dried to polystyrene 96-well microtiter plates as the platecoat antigen. Positive supernatants were subsequently confirmed by FACS analysis and RIP using P3G2 transfected COS-1 cells. Hybridomas were cloned and followed using the same assays. Monoclonal cultures were expanded and supernatants purified by affinity chromatography using BioRad Protein A agarose.

**[0141]** The monoclonal antibodies to P3G2/Fc were used to screen cells and cell lines using standard flow cytometry procedures to identify cells on which P3G2 is expressed. Cell lines and cells screened in the flow cytometry analyses were CB23, CB39, RAJI, AK778, K299, PS-1, U937, THP-1, JURKAT and HSB2. For each cell line or cell sample screened, the procedure involved incubating approximately 100,000 of the cells blocked with 2% PCS (fetal calf serum), 5% normal Goat serum and 5% rabbit serum in PBS with 5 $\mu$ g of FITC conjugated mouse anti-P3G2 antibody for 1 hour. Following the incubation the sample was washed 2 times with FACS buffer (2% FCS in PBS). The cells were analyzed for any bound protein using fluorescent detection flow cytometry to detect FITC. The results indicated that LIR-P3G2 antibody binds well to B cell lines CB23 and RAJI1; monocytic cell lines THP-1 and U937; and primary B cell and primary monocytes. The highest expression of LIR-P3G2 was shown on monocytes that stained brightly for CD16 and less brightly for CD14 and CD64. The antibody does not bind detectably to T cell lines nor does it bind detectably to primary T cells.

**[0142]** In a related experiment, the P3G2 antibody generated as described above was used in immunoprecipitation experiments. The immunoprecipitation analyses involved first surface biotinylating 2.5 x 10<sup>6</sup> monocytes by washing the cells with PBS and suspending the cells in a biotinylation buffer of 10 mM sodium borate and 150 mM NaCl at pH 8.8, followed by adding 5  $\mu$ L of a 10 mg/mL solution of biotin-CNHS-ester (D-biotinoyl-e-aminocaproic acid-N-hydroxysuccinimide ester purchased from Amersham) in DMSO to the cells. After quenching the reaction with 10  $\mu$ L of 1 M ammonium chloride per 1 mL of cells and washing the cells in PBS, the cells were lysed in 1 mL of 0.5% NP40-PBS and the lysate was recovered following centrifugation. Then 100  $\mu$ L of 0.5%NP40-PBS was added to 150  $\mu$ L of the lysate and the resulting mixture was incubated with 2  $\mu$ g/mL of antibody, at 4°C for 16 hours. Fifty microliters of 50% Protein A-Sepharose slurry was added to the antibody mixture and the slurry was shaken at 4°C for 1 hour. The slurry was centrifuged and the resulting pellet was washed with 0.75 mL of 0.5% NP40 in PBS six times. Protein bound to the Protein A-Sepharose was eluted with 30  $\mu$ L of SDS-PAGE reducing sample buffer and heating at 100°C for five minutes.

**[0143]** The eluted proteins were analyzed using 4-20% gradient SDS-PAGE with enhanced chemiluminescence (ECL) protein markers. Then the electrophoreses samples were transferred in a Western Blot onto nitrocellulose membranes. The membranes were treated with blocking reagent (0.1% Tween-20 and 3% nonfat dry milk in PBS) for one hour at room temperature and then they were washed once for 15 minutes followed and twice for 5 minutes with 0.1% Tween-20 in PBS. The washed membranes were incubated with 10 mL of 1:100 HRP-Streptavidin for 30 minutes and then washed 1 times for 15 minutes followed by 4 times for 5 minutes with 0.1% Tween-20 in PBS.

**[0144]** Bound streptavidin HRP was detected with ECL Detection Reagents purchased from Amersham and used according to manufacturer's instructions. The developed membranes were exposed to x-ray film and then visualized. The results showed that LIR-P3G2 was immunoprecipitated from CB23 cells and P3G2 transfected COS-1 cells, indicating that P3G2 is expressed by these cells.

#### **Example 6. Screening Cells and Cell Lines for Binding to P3G2**

**[0145]** The following describes flow cytometry analyses used to identify cells and cell lines which bind to P3G2. The cells and cell lines tested were CB23, HSB2, MP-1, Jurkat, primary T cells, primary B cells, and primary NK cells. For each cell line or cell line tested the procedure involved washing the cells three times with FACS buffer (2% FCS in PBS with 0.2% azide) and incubating each sample (10<sup>5</sup> cells) in 100  $\mu$ L blocking buffer (2% FCS, 5% NGS, 5% rabbit serum



in PBS) for one hour. For each cell line 4 test samples were prepared, one each having 0, 2, 5, or 10 µg of W6/32 (ATCC HB-95) in 100 µL blocking buffer added to the samples, respectively. W6/32 is an antibody against MHC Class I heavy chains (an anti HLA-A, B, and C molecule). Following the addition of the W6/32 solution, the samples were incubated on ice for 1 hour and then washed three times with 200 µL of FACS buffer. Then 5 µg of P3G2/Fc in blocking buffer was added to each sample and they were incubated on ice for one hour. The P3G2/Fc competes with W6/32 for binding sites on the cells.

[0146] Following the incubation, the cells were washed three times with 200 µL of FACS buffer and treated with mouse anti human Fc/biotin and SAPE for 45 minutes. This treatment causes the anti human Fc/biotin to bind to any cell bound sP3G2/Fc and the SAPE to bind to the anti human F/Biotin. Since the SAPE is a fluorescing compound its detection using appropriate excitation and emission conditions positively identifies cell bound P3G2/Fc. Finally the treated cells were washed three times with FACS buffer and subjected to flow cytometry to identify cells bound to protein.

[0147] The results demonstrated that W6/32 competed with P3G2 for binding to all cells and cell lines tested. The P3G2 binding was totally blocked at 5 µg W6/32 indicating that W6/32 and P3G2 are binding to the same or overlapping sites on the MHC Class I heavy chains.

#### **Example 7. Screening HSB2 cDNA Library to Isolate a P3G2 Binding Ligand**

[0148] The following describes screening a cDNA library from one of the cell lines, HSB-2, a T lymphoblastic leukemia cell line, found to bind P3G2, and identifying a P3G2 binding ligand. An HSB2 cDNA library in the mammalian expression vector pDC302, was prepared as generally described in U.S. Patent No. 5,516,658 and specifically in Kozlosky et al. *Oncogene* 10:299-306, 1995. Briefly, mRNA was isolated from sorted HSB-2 cells and a first cDNA strand was synthesized using 5 µg polyA<sup>+</sup> and the reverse transcriptase AMV RTase from Life Science. The second cDNA strand was synthesized using DNA polymerase I from BRL at concentration of 1.5 U/µL. Using standard techniques as described in Haymerle et al., *Nucl. Acids Res.* 14:8615, 1986, the cDNA was ligated into the appropriate site of the pDC302 vector.

[0149] *E. coli* strain DH5α cells were transformed with the cDNA library in pDC302. After amplifying the library a titer check indicated that there was a total of 157,200 clones. The transformed cells were plated into 15 different plates. Plasmid DNA was isolated from pools consisting of approximately 2,000 clones per pool. The isolated DNA was transfected into CV1-EBNA cells (ATCC CRL 10478) using DEAE-dextran followed by chloroquine treatment. The CV1-EBNA cells were maintained in complete medium (Dulbecco's modified Eagles' media containing 10% (v/v) fetal calf serum, 50 U/mL penicillin, 50 U/mL streptomycin, and 2 mM L-glutamine) and were plated to a density of approximately 2 x 10<sup>5</sup> cells/well in single-well chambered slides. The slides had been pre-treated with 1 mL of a solution of 10 µg/mL human fibronectin in PBS for 30 minutes followed by a single washing with PBS. Media was removed from adherent cells growing in a layer and replaced with 1.5 mL complete medium containing 66.6 µM chloroquine sulfate. About 0.2 mL of a DNA solution (2 µg DNA, 0.5 mg/mL DEAE-dextran in complete medium containing chloroquine) was added to the cells and mixture was incubated at 37 °C for about five hours. Following incubation media was removed and the cells were shocked by adding complete medium containing 10% DMSO for 2.5 minutes. After shocking the cells the complete medium was replaced with fresh complete medium and the cells were grown in culture for three days to permit transient expression of the inserted DNA sequences. These conditions led to a 30% to 80% transfection frequency in surviving CV1-EBNA cells.

[0150] Each slide was incubated with 1 mL of P3G2:Fc at a concentration of 0.45 µg/mL in binding buffer (RPMI 1640 containing 25 mg/mL bovine serum albumin, 2 mg/mL sodium azide, 20 mM Hepes at pH 7.2, and 50 mg/mL nonfat dry milk) at room temperature for 1 hour. After incubating the slides, they were washed with binding buffer and then incubated with Fc specific <sup>125</sup>I-mouse anti-human IgG (see Goodwin et al. *Cell* 73:447-456, 1993). This was followed by a second wash with buffer after which the slides were fixed with a 2.5% glutaraldehyde/PBS solution, washed in PBS and allowed to air dry. The slides were dipped in Kodak GTNB-2 photographic emulsion (6x dilution in water). After air drying the slides were placed in a dark box and refrigerated. After three days the slides were developed in Kodak D19 developer, rinsed in water and fixed in Agfa G433C fixer. The fixed slides were individually examined under a microscope at 25-40x magnification. Positive pools demonstrating binding of sP3G2:Fc were visualized by the presence of autoradiographic silver grains against the film background. Two positive pools were titrated and plated to provide plates containing approximately 200 colonies each. Each plate was scraped to provide pooled plasmid DNA for transfection into CV1-EBNA cells and screening as described above. Following subsequent breakdowns and screenings, one positive individual colony was obtained for each pool. The cDNA insert of the positive clones were identified as HLA-B44 and HLA-A2, class I MHC antigens.

#### **Example 8. Northern Blot Analysis**

[0151] Since the experiments described in Example 4 resulted in the detection of LIR-P3G2 surface expression on

a number of cell lines, conventional Northern Blot analysis procedures were used to study the expression of LIR-P3G2 and any LIR-P3G2 related mRNAs in different tissue types. The cell lines selected for Northern Blot analysis were RAJI, PBT, PBM, YT, HEP3B, HELA, KB, KG-1, IMTLH, HPT, HFF, THP-1, and U937. The following describes the Northern Blot analysis and the analysis results.

**[0152]** The cDNA encoding the extracellular region of P3G2 was isolated using primers which flank the extracellular region of P3G2 and having the following sequences:

Sal I  
5' - TAT GTC GAC CAT GAC CCC CAT CCT CAC GGT - 3' (SEQ ID NO:5)

Bgl II  
5' - TAT AGA TCT ACC CCC AGG TGC CTT CCC AGA CCA (SEQ ID NO:27)

The PCR template was the full length P3G2 gene isolated as described in Example 3 above. The conditions for the PCR reaction were as follows: One cycle at 95°C for 5 minutes; 30 cycles which included 95°C for 45 seconds, 64 °C for 45 seconds and 72° C for 45 seconds; and, one cycle at 72°C for 5 minutes. The PCR product was cloned into PCR II vector, purchased from Invitrogen, in accordance with the supplier's instructions. The isolated DNA encoding the extracellular region of P3G2 was used to make a riboprobe with the Ambion MAXISCRIPIT Kit according to the manufacturer's instructions.

**[0153]** Northern blots containing poly A+ selected RNA or total RNA from a variety of human cell lines were prepared by resolving RNA samples on a 1.1% agarose-formaldehyde gel, blotting onto Hybond-N as recommended by the manufacturer (Amersham Corporation) and staining with methylene blue to monitor RNA concentrations. The blots were prepared using 1 µg of the PolyA+ RNA or 10 µg of total RNA and each blot was probed with 10<sup>6</sup> cpm/mL RNA extracellular P3G2 riboprobe, prepared as just described, at 63°C for 16 hours. The probed blots were washed with 2 x SSC at 63°C for 30 minutes 2 times; 1 x SSC at 63°C for 30 minutes 2 times; and, 0.1 x SSC at 63°C for 5 minutes 2 times.

**[0154]** The probed blots were autoradiographically developed. The developed blots showed that the P3G2 RNA hybridized to a 3.5 kb RNA expressed by RAJI, CB23 and U937; an approximately 1.5kb RNA expressed by THP-1; and multiple RNAs ranging from 1.5 kb to 3.5 kb expressed by PBM. These results suggest that different genes having extracellular domains similar in structure to that of P3G2 may be expressed by peripheral blood monocytes.

#### **Example 9. Probing PBM cDNA Library to Isolate LIR Polypeptides**

**[0155]** The following describes steps taken to screen a peripheral blood monocyte cDNA library to isolate polypeptides relating to the P3G2 polypeptide using conventional Southern Blot methodologies. A peripheral blood monocyte cDNA library was prepared using substantially the same procedures described in Example 7.

**[0156]** DNA from an initial 15 pools of cDNA having 10,000 clones per pool was digested with *Bgl*/II restriction enzyme and electrophoresed on a 1 % agarose gel at 100 V for 2 hours. Southern Blots were prepared by electroblotting the electrophoresed DNA in 0.55% TBE buffer onto Hybond membranes. The blotted DNA was denatured in 0.5 M NaOH in 0.6M NaCl solution for 5 minutes and then neutralized in 0.5 M TRIS in 1.5 M NaCl at pH 7.8 for 5 minutes. The membranes were placed in a STRATALINKER UV crosslinker for 20 seconds to crosslink the blotted DNA to the membrane. The membrane and bound DNA were placed in pre-hybridization solution of 10X Denhart's Solution, 0.05M TRIS at pH 7.5, 0.9M NaCl, 0.1% sodium pyrophosphate, 1% SDS and 200 µg/mL salmon sperm DNA at 63° C for 2 hours and then the bound DNA was probed with <sup>32</sup>P labeled probe of DNA encoding the extracellular region of LIR-P3G2, including the signal peptide and *Sal* I and *Bgl* II restriction sites. The concentration of the DNA probe in hybridization solution was 10<sup>6</sup> CPM per mL of hybridization solution. The probed blots were incubated for 16 hours at 63°C and then washed with 2x SSC at 63°C for 1 hour with one solution change; 1x with SSC at 63°C for one hour with one solution change; and, with 0.1x SSC at 68°C for 45 minutes with one solution change. After drying the blots they were autoradiographically developed and visualized for DNA bands which hybridized to the P3G2 extracellular DNA probe.

**[0157]** The results of the autoradiography visualization indicated that all pools contained DNA which hybridized to the probe. One pool showing 7 positive DNA bands was selected and subsequently subdivided to 10 pools having 3,000 clones per pool. Applying subsequent Southern Blotting methodologies to the 10 pools resulted in one pool showing 9 positively hybridizing DNA sequences. Single hybridizing clones were isolated by standard colony hybridi-

zation techniques.

**[0158]** Duplicate bacterial colonies on filters were probed with the P3G2 extracellular probe described above at a concentration of 500,000 cpm/mL at 63°C for 16 hours. The hybridized filters were washed with 2x SSC at 63°C for 30 minutes; with 1x SSC at 63°C for 30 minutes; and finally with 0.1 X SSC at 68°C for 15 minutes.

**[0159]** Forty-eight clones were visualized as hybridizing on duplicate filters by autoradiography and DNA obtained from these clones using standard DNA preparation methodologies was digested with *Bgl* II. Then Southern Blots of the digests were obtained and probed with the P3G2 extracellular probe described above. Seven different sized cloned inserts were identified as positively hybridizing to the P3G2 probe. The nucleotide sequence of each of the inserts was obtained using automated sequencing technology. Of the 8 different cloned inserts, one was identical in sequence to LIR-P3G2. The others were identified as DNA encoding polypeptides of the new LIR family of polypeptides. The nucleotide sequences (cDNA) of the isolated LIR family members are presented in SEQ ID NO:7 (designated pbm25), SEQ ID NO:9 (designated pbm8), SEQ ID NO:11 (designated pbm36-2), SEQ ID NO:13 (designated pbm36-4); SEQ ID NO:15 (designated pbmhh); SEQ ID NO:17 (designated pbm2) and SEQ ID NO:19 (designated pbm17). The amino acid sequences encoded thereby are presented in SEQ ID NO:8 (designated pbm25), SEQ ID NO:10 (designated pbm8), SEQ ID NO: 12 (designated pbm36-2), SEQ ID NO: 14 (designated pbm36-4), SEQ ID NO: 16 (designated pbmhh); SEQ ID NO:18 (designated pbm2); and SEQ ID NO:20 (designated pbm17).

#### **Example 10. Screening a Human Dendritic Cell cDNA Library for LIR cDNA Sequences**

**[0160]** The following describes the isolation and identification of an LIR family member by screening a human bone marrow-derived dendritic cell cDNA library in the  $\lambda$  Zap vector with a radiolabeled Hh0779 cDNA fragment. The Hh0779 cDNA fragment is a 0.7kb insert of the Hh0779 clone previously isolated from a human dendritic cell cDNA library and obtained by restriction digestion with the enzymes PstI and SpeI. The Hh0779 cDNA fragment was labeled with [ $\alpha$ -<sup>32</sup>P] dCTP using the DECAprime II DNA labeling kit purchased from Ambion.

**[0161]** The  $\lambda$  Zap cDNA library was plated at a density of 20,000 pfu per plate to provide a total of 480,000 plaques for the initial screening. The  $\lambda$  Zap cDNA was blotted in duplicate onto Hybond membranes, purchased from Amersham, and then denatured in a solution of 0.5N NaOH and 0.5M NaCl for 5 minutes. The membranes were neutralized in a solution of 0.5M Tris (pH 7.8) and 1.5M NaCl for 5 minutes, and then washed in 2x SSC for 3 minutes. The cDNA was crosslinked to the Hybond membranes using a STRATALINKER UV crosslinker in the auto setting.

**[0162]** The membranes were pre-hybridized at 65°C for 2.25 hours in hybridization buffer containing 10x Denhardt's, 0.05M Tris (pH 7.5), 0.9M NaCl, 0.1% sodium pyrophosphate, 1% SDS and 4 mg/mL heat denatured salmon sperm DNA. After the pre-hybridization, the radiolabeled Hh0779 cDNA was added to the hybridization buffer to a final concentration of  $0.54 \times 10^6$  cpm/mL. After 24 hours of hybridization, the membranes were washed in 0.25xSSC, 0.25% SDS at 65°C for 1.5 hours. The blots were then exposed to autoradiographic film to visual positive clones.

**[0163]** A total of 146 positive clones showing hybridization signals in both membranes of a duplicate set were identified, isolated, and saved for future use. Of the 146 clones, 35 were selected for secondary screening. The selected clones were plated at low density and single clones were isolated after hybridization to the Hh0779 probe using the hybridization conditions described above. The plasmids were then isolated from the  $\lambda$  Zap clones using the VCSM13 helper phage purchased from Stratagene. The plasmid DNA was analyzed by restriction digestion and PCR, and the clones containing the 24 largest inserts were selected and sequenced. Of the 24 sequenced clones, 6 encoded LIR-P3G2, 3 encoded LIR-pbm2, 8 encoded LIR-pbm36-4 and LIR-pbm36-2, 1 encoded LIR-pbm8, 2 encoded LIR-pbmhh, and 1 encoded a novel sequence designated LIR-pbmnew. Three clones were identified as encoding amino acid sequences that are not relevant to the LIR polypeptide family.

#### **Example 11. Association of LIR-P3G2 and LIR-pbm8 with Tyrosine Phosphatase, SHP-1**

**[0164]** The following describes the tests performed to demonstrate that LIR-P3G2 and LIR-pbm8 associate with SHP-1. Human monocytes were cultured in RPMI medium supplemented with 10% FBS, concentrated by centrifugation and finally subdivided into two aliquots. One aliquot was stimulated with a solution of 50 mM/mL sodium pervanadate for 5 minutes. The second aliquot was not stimulated. After stimulation, the cells in each aliquot were immediately lysed in RIPA buffer containing 1% NP-40, 0.5% sodium deoxycholate, 50 mM Tris pH8, 2 mM EDTA, 0.5 mM sodium orthovanadate, 5mM sodium fluoride, 25mM  $\beta$ -glycerol phosphate, and protease inhibitors. Samples of  $24 \times 10^6$  cell equivalents were incubated for 2 hours at 4°C with either 5  $\mu$ g/mL of anti-SHP-1 antibody purchased from Transduction Laboratories, or 5  $\mu$ g/mL of an isotype-matched antibody control (anti-Flag-M5 IgG1). The resulting immunocomplexes were precipitated by incubation with protein G-agarose (Boehringer Mannheim), washed, and resuspended in 40 mL of 2x SDS-PAGE sample buffer. Twenty microliters of each immunoprecipitate were loaded onto electrophoresis gels, electrophoresed under reducing conditions, and transferred to nitrocellulose membranes purchased from Amersham. Western blots were probed with anti-LIR-P3G2 monoclonal antibody sera and anti-LIR-pbm8 monoclonal antibody

antisera and the immunocomplexes were detected by enhanced chemiluminescence (NEN).

[0165] A protein having a molecular weight of approximately 120kDa, corresponding to LIR-P3G2 was readily detected in SHP-1 immunoprecipitates, but not the immunoprecipitates generated with the anti-Flag-M5 antibody control. Similarly, a protein of 90-100kDa, corresponding to LIR-pbm8, was detected in SHP-1 immunoprecipitates, but not in the control immunoprecipitates. Neither the LIR-P3G2 band nor the LIR-pbm8 band was seen in the absence of sodium pervanadate treatment. This confirms that tyrosine phosphorylation of LIR-P3G2 is essential for the association of LIR-P3G2 and SHP-1 and phosphorylation of LIR-pbm8 is essential for the association of LIR-pbm8 and SHP-1.

[0166] To study the inhibition of FcγRI-mediated tyrosine phosphorylation events upon LIR coligation, peripheral blood monocytes were incubated with or without 10 μg/mL of F(ab)<sub>2</sub> version of a number of antibodies (α-LIR-1+α-LIR-2, α-CD11c, αCD14, αCD64, α-CD64+α-LIR-1, α-CD64+α-LIR-2, α-CD64+α-LIR-1+α-LIR-2, αCD64+α-CD11c, α-CD64+α-CD14). This was followed with crosslinking with 30 μ/mL of polyclonal F(ab)<sub>2</sub> goat anti-mouse. Cell lysates were immunoprecipitated overnight with anti-phosphotyrosine conjugated agarose, electrophoresed, and transferred onto nitrocellulose. Western blotting was performed using a combination of PY-20 and 4G10 HRP-conjugated anti-phosphotyrosine mAbs. This data demonstrates the specific inhibition of FcγRI-mediated tyrosine phosphorylation events upon LIR-P3G2 and LIR-pbm8 coligation.

#### **Example 12. Generating Antibodies Immunoreactive with LIR Polypeptides**

[0167] The following describes generating monoclonal antibody immunoreactive with LIR family members. A purified LIR polypeptide is prepared by COS-1 cell expression and affinity purification as described in Example 4. The purified protein or cells transfected with an expression vector encoding the full length protein can generate monoclonal antibodies against the LIR polypeptide using conventional techniques, for example those techniques described in U.S. Patent 4,411,993. Briefly BALB-C mice are immunized at 0, 2 and 6 weeks with 10μg of the LIR polypeptide. The primary immunization is prepared with TITERMAX adjuvant and subsequent immunizations are prepared with incomplete Freund's adjuvant (IFA). At 11 weeks, the mice are IV boosted with 3-4 μg the LIR polypeptide in PBS. Three days after the IV boost, splenocytes are harvested and fused with an Ag8.653 myeloma fusion partner using 50% aqueous PEG 1500 solution. Hybridoma supernatants are -screened by ELISA using the LIR transfected cells in PBS at 7 X 10<sup>3</sup> cells per well and dried to polystyrene 96-well microtiter plates as the platecoat antigen. Positive supernatants are subsequently confirmed by FACS analysis and RIP using LIR transfected cells. Hybridomas are cloned and followed in the same manner of screening. Monoclonal cultures are expanded and supernatants purified by affinity chromatography.

#### **Example 13. Flow Cytometric Analysis For Expression of LIR-P3G2 and LIR-pbm8 on Lymphoid and Myeloid Cells**

[0168] In order to compare the differential expression and distribution of LIR-P3G2 and LIR-pbm8 on lymphocyte populations, freshly isolated peripheral blood mononuclear cells (PBMC) were stained with PE-labeled anti-CD3, anti-CD19, or anti-CD56 mAb in the presence of either biotin labeled anti-LIR-P3G2 or anti LIR-pbm8 mAb. Then the stained cells were treated with APC-labeled streptavidin. Density plots representing 5x10<sup>4</sup> events were collected on a FACScaliber (from Beckton Dickinson). The results demonstrated that LIR-P3G2 is expressed on 80%-95% of CD19<sup>+</sup> B cells, on 5%-15% CD3<sup>+</sup> T cells, and on 10%-30% CD56<sup>+</sup> NK cells. On the cells examiner from the same 12 donors, LIR-pbm8 expression was not detected on CD19<sup>+</sup> B cells, CD3<sup>+</sup> T cells, and CD56<sup>+</sup> NK cells.

[0169] Countercurrent elutriated fractions containing a high percentage of circulating monocytes and dendritic cells (DC) were obtained. The monocytes were characterized according to the phenotypes subsets CD14<sup>+</sup>CD16<sup>-</sup> and CD14<sup>+</sup>CD16<sup>+</sup>. The peripheral blood DC were characterized with the phenotype CD33<sup>+</sup>CD14<sup>+</sup>CD16<sup>+</sup>HLA-DR<sup>+</sup>. The monocytes subsets and DC's were stained with FITC-labeled antiCD14, PE-labeled anti CD3, perCp-labeled antiHLA-DR, and either biotin-labeled anti-CD16, anti-LIR-P3G2, or anti LIR-pbm8. Then the stained cells were treated with APC-labeled streptavidin. Both monocyte subsets co-express similar levels of LIR-P3G2 and LIR-pbm8, with the highest LIR-P3G2 and LIR-pbm8 expression detected on the CD14<sup>+</sup>CD16<sup>+</sup> subset. Blood DC express lower levels of LIR-P3G2 and LIR-pbm8 compared to monocytes. The results of these experiments demonstrate the LIR-P3G2 is expressed on lymphocytes, monocytes and DC, and LIR-pbm8 is expressed on monocytes and DC.

#### **Example 14. Screening LIR-P3G2 and LIR-pbm8 Binding to HLA Class I Alleles**

[0170] The following describes flow cytometry analyses used to screen LIR-P3G2 and LIR-pbm8 for binding to HLA Class I alleles. The B lymphoblastoid class I-deficient 721.221 cell line, untransfected or transfected with a panel of HLA class I alleles was used for staining. LIR-P3G2/Fc and LIR-pbm8/Fc fusion proteins were used in the binding studies and both bound detectably to seven of the eleven HLA-A, HLA-B and HLA-C alleles that were tested. In general,

LIR-P3G2/Fc and LIR-pbm8/Fc bind with higher affinity to HLA-B alleles than to HLA-A or HLA-C alleles. W6/32 (ATCC HB-95), an antibody against MHC Class I heavy chains (an anti HLA-A, B, and C molecule) inhibits LIR-P3G2/Fc and LIR-pbm8/Fc binding to all class I transfectants. Finally, LIR-P3G2 and LIR-pbm8 binding does not correlate with the MHC class I expression levels. Thus, LIR-P3G2 and LIR-pbm8 bind to several HLA-A, -B, and -C alleles, and recognize a similar broad spectrum of MHC class I specificities.

#### **Example 15. Isolation of LIR-9m1, LIR-9m, LIR-9s1, LIR-9s2 and LIR-10**

**[0171]** In the course of high throughput sequencing of a human dendritic cell cDNA library, it was noted that the sequence of an incomplete cDNA (clone ss4894) was strikingly similar to the nucleotide sequences of LIRs 6a, 6b and 7, thus suggesting that ss4894 was a member of the LIR gene family. To obtain the remainder of this cDNA clone, the Rapid Amplification cDNA Extension system (RACE) was used to amplify a human leukocyte cDNA library (Chenchik et al., A new method for full-length cDNA cloning by PCR, In *A Laboratory Guide to RNA: Isolation, Analysis, and Synthesis*, Ed. Kreig, P.A. (Wiley-Liss, Inc.), pages 273-321). The first round of amplification employed one primer corresponding to the RACE adapter at the 5' end of the cDNAs, and a second primer corresponding to sequences near the 3' end of ss4894. This effort yielded several clones that contained sequence that was highly homologous though not identical to that of ss4894 and that extended upstream beyond an initiating methionine codon. These clones, however, lacked some of the sequence at the 3' end of the coding region. In an effort to obtain an entire coding region, another round of RACE sequencing was performed, this time using a first primer from near the 5' end of the first RACE products, and a second primer corresponding to the 3' adapter. This effort yielded five clones containing LIR inserts, four of which are closely related and appear to encode variants of the same gene. These four closely related cDNA sequences were designated LIR-9m1, LIR-9m2, LIR-9s1 and LIR-9s2 (SEQ ID NOS:29, 31, 33 and 35). The fifth of the clones obtained using this last set of primers represented a different gene, which has been designated LIR-10 (SEQ ID NO:37).

**[0172]** All four of the LIR-9 clones encode variants of the same protein, and are presumed to be the products of alternative splicing. The proteins encoded by LIR-9m1 (SEQ ID NO:30) and LIR-9s1 (SEQ ID NO:34) contain a 12 amino acid insert that is absent from LIR-9m2 (SEQ ID NO:32) and LIR-9s2 (SEQ ID NO:36). The soluble forms of the LIR-9 protein, i.e., LIR-9s1 and LIR-9s2, diverge near their carboxy termini from the membrane forms, i.e., LIRs-9m1 and -9m2. This divergence presumably is due to different exons being used by the soluble and membrane forms to encode that region of the protein.

## SEQUENCE LISTING

5 <110> Immunex Corporation  
 Cosman, David J.  
 Anderson, Dirk M.  
 Borges, Luis

10 <120> Family of Immunoregulators Designated Leukocyte Immunoglobulin-  
 Like Receptors (LIR)

<130> 2624A-WO

<140> to be assigned  
 <141> 2000-05-12

15 <150> 08/842,248  
 <151> 1997-04-24

<150> 09/310,463  
 <151> 1999-05-12

20 <160> 39

<170> PatentIn Ver. 2.0

25 <210> 1  
 <211> 2922  
 <212> DNA  
 <213> human

<220>  
 <221> CDS  
 30 <222> (310)..(2262)

<400> 1  
 agggccacgc gtgcatgcgt cgactggaac gagacgacct gctgtgaccc ccttgtgggc 60  
 actccattgg ttttatggcg cctctacttt ctggagtttg tgtaaaacaa aaatattatg 120  
 35 gtctttgtgc acatttacat caagctcagc ctggcgcgca cagccagatg cgagatgcgt 180  
 ctctgctgat ctgagctcgc ctgcagcatg gacctgggtc ttccctgaag catctccagg 240  
 gctggaggga cgactgccat gcaccgaggg ctcatccatc cacagagcag ggcagtggga 300  
 40 ggagacgcc atg acc ccc atc ctc acg gtc ctg atc tgt ctc ggg ctg 348  
 Met Thr Pro Ile Leu Thr Val Leu Ile Cys Leu Gly Leu  
 1 5 10

45 agt ctg ggc ccc cgg acc cac gtg cag gca ggg cac ctc ccc aag ccc 396  
 Ser Leu Gly Pro Arg Thr His Val Gln Ala Gly His Leu Pro Lys Pro  
 15 20 25

acc ctc tgg gct gaa cca ggc tct gtg atc acc cag ggg agt cct gtg 444  
 Thr Leu Trp Ala Glu Pro Gly Ser Val Ile Thr Gln Gly Ser Pro Val  
 50 30 35 40 45

acc ctc agg tgt cag ggg ggc cag gag acc cag gag tac cgt cta tat 492  
 Thr Leu Arg Cys Gln Gly Gly Gln Glu Thr Gln Glu Tyr Arg Leu Tyr  
 50 55 60

EP 1 199 362 A2

	aga gaa aag aaa aca gca ccc tgg att aca cgg atc cca cag gag ctt	540
	Arg Glu Lys Lys Thr Ala Pro Trp Ile Thr Arg Ile Pro Gln Glu Leu	
	65 70 75	
5	gtg aag aag ggc cag ttc ccc atc cca tcc atc acc tgg gaa cat gca	588
	Val Lys Lys Gly Gln Phe Pro Ile Pro Ser Ile Thr Trp Glu His Ala	
	80 85 90	
10	ggg cgg tat cgc tgt tac tat ggt agc gac act gca ggc cgc tca gag	636
	Gly Arg Tyr Arg Cys Tyr Tyr Gly Ser Asp Thr Ala Gly Arg Ser Glu	
	95 100 105	
15	agc agt gac ccc ctg gag ctg gtg gtg aca gga gcc tac atc aaa ccc	684
	Ser Ser Asp Pro Leu Glu Leu Val Val Thr Gly Ala Tyr Ile Lys Pro	
	110 115 120 125	
20	acc ctc tca gcc cag ccc agc ccc gtg gtg aac tca gga ggg aat gta	732
	Thr Leu Ser Ala Gln Pro Ser Pro Val Val Asn Ser Gly Gly Asn Val	
	130 135 140	
25	acc ctc cag tgt gac tca cag gtg gca ttt gat ggc ttc att ctg tgt	780
	Thr Leu Gln Cys Asp Ser Gln Val Ala Phe Asp Gly Phe Ile Leu Cys	
	145 150 155	
30	aag gaa gga gaa gat gaa cac cca caa tgc ctg aac tcc cag ccc cat	828
	Lys Glu Gly Glu Asp Glu His Pro Gln Cys Leu Asn Ser Gln Pro His	
	160 165 170	
35	gcc cgt ggg tcg tcc cgc gcc atc ttc tcc gtg ggc ccc gtg agc ccg	876
	Ala Arg Gly Ser Ser Arg Ala Ile Phe Ser Val Gly Pro Val Ser Pro	
	175 180 185	
40	agt cgc agg tgg tgg tac agg tgc tat gct tat gac tcg aac tct ccc	924
	Ser Arg Arg Trp Trp Tyr Arg Cys Tyr Ala Tyr Asp Ser Asn Ser Pro	
	190 195 200 205	
45	tat gag tgg tct cta ccc agt gat ctc ctg gag ctc ctg gtc cta ggt	972
	Tyr Glu Trp Ser Pro Ser Asp Leu Leu Leu Val Leu Gly	
	210 215 220	
50	gtt tct aag aag cca tca ctc tca gtg cag cca ggt cct atc gtg gcc	1020
	Val Ser Lys Lys Pro Ser Leu Ser Val Gln Pro Gly Pro Ile Val Ala	
	225 230 235	
55	cct gag gag acc ctg act ctg cag tgt ggc tct gat gct ggc tac aac	1068
	Pro Glu Glu Thr Leu Thr Leu Gln Cys Gly Ser Asp Ala Gly Tyr Asn	
	240 245 250	
60	aga ttt gtt ctg tat aag gac ggg gaa cgt gac ttc ctt cag ctc gct	1116
	Arg Phe Val Leu Tyr Lys Asp Gly Glu Arg Asp Phe Leu Gln Leu Ala	
	255 260 265	
65	ggc gca cag ccc cag gct ggg ctc tcc cag gcc aac ttc acc ctg ggc	1164
	Gly Ala Gln Pro Gln Ala Gly Leu Ser Gln Ala Asn Phe Thr Leu Gly	
	270 275 280 285	
70	cct gtg agc cgc tcc tac ggg ggc cag tac aga tgc tac ggt gca cac	1212
	Pro Val Ser Arg Ser Tyr Gly Gly Gln Tyr Arg Cys Tyr Gly Ala His	
	290 295 300	



EP 1 199 362 A2

	aac ctc tcc tcc gag tgg tgc gcc ccc agc gac ccc ctg gac atc ctg	1260
	Asn Leu Ser Ser Glu Trp Ser Ala Pro Ser Asp Pro Leu Asp Ile Leu	
	305 310 315	
5	atc gca gga cag ttc tat gac aga gtc tcc ctc tgc gtg cag ccg ggc	1308
	Ile Ala Gly Gln Phe Tyr Asp Arg Val Ser Leu Ser Val Gln Pro Gly	
	320 325 330	
10	ccc acg gtg gcc tca gga gag aac gtg acc ctg ctg tgt cag tca cag	1356
	Pro Thr Val Ala Ser Gly Glu Asn Val Thr Leu Leu Cys Gln Ser Gln	
	335 340 345	
15	gga tgg atg caa act ttc ctt ctg acc aag gag ggg gca gct gat gac	1404
	Gly Trp Met Gln Thr Phe Leu Leu Thr Lys Glu Gly Ala Ala Asp Asp	
	350 355 360 365	
20	cca tgg cgt cta aga tca acg tac caa tct caa aaa tac cag gct gaa	1452
	Pro Trp Arg Leu Arg Ser Thr Tyr Gln Ser Gln Lys Tyr Gln Ala Glu	
	370 375 380	
25	ttc ccc atg ggt cct gtg acc tca gcc cat gcg ggg acc tac agg tgc	1500
	Phe Pro Met Gly Pro Val Thr Ser Ala His Ala Gly Thr Tyr Arg Cys	
	385 390 395	
30	tac ggc tca cag agc tcc aaa ccc tac ctg ctg act cac ccc agt gac	1548
	Tyr Gly Ser Gln Ser Ser Lys Pro Tyr Leu Leu Thr His Pro Ser Asp	
	400 405 410	
35	ccc ctg gag ctc gtg gtc tca gga ccg tct ggg ggc ccc agc tcc ccg	1596
	Pro Leu Glu Leu Val Val Ser Gly Pro Ser Gly Gly Pro Ser Ser Pro	
	415 420 425	
40	aca aca ggc ccc acc tcc aca tct ggc cct gag gac cag ccc ctc acc	1644
	Thr Thr Gly Pro Thr Ser Thr Ser Gly Pro Glu Asp Gln Pro Leu Thr	
	430 435 440 445	
45	ccc acc ggg tgc gat ccc cag agt ggt ctg gga agg cac ctg ggg gtt	1692
	Pro Thr Gly Ser Asp Pro Gln Ser Gly Leu Gly Arg His Leu Gly Val	
	450 455 460	
50	gtg atc ggc atc ttg gtg gcc gtc atc cta ctg ctc ctc ctc ctc ctc	1740
	Val Ile Gly Ile Leu Val Ala Val Ile Leu Leu Leu Leu Leu Leu Leu	
	465 470 475	
55	ctc ctc ttc ctc atc ctc cga cat cga cgt cag ggc aaa cac tgg aca	1788
	Leu Leu Phe Leu Ile Leu Arg His Arg Arg Gln Gly Lys His Trp Thr	
	480 485 490	
60	tcg acc cag aga aag gct gat ttc caa cat cct gca ggg gct gtg ggg	1836
	Ser Thr Gln Arg Lys Ala Asp Phe Gln His Pro Ala Gly Ala Val Gly	
	495 500 505	
65	cca gag ccc aca gac aga ggc ctg cag tgg agg tcc agc cca gct gcc	1884
	Pro Glu Pro Thr Asp Arg Gly Leu Gln Trp Arg Ser Ser Pro Ala Ala	
	510 515 520 525	
70	gat gcc cag gaa gaa aac ctc tat gct gcc gtg aag cac aca cag cct	1932
	Asp Ala Gln Glu Glu Asn Leu Tyr Ala Ala Val Lys His Thr Gln Pro	
	530 535 540	
75	gag gat ggg gtg gag atg gac act cgg agc cca cac gat gaa gac ccc	1980

EP 1 199 362 A2

	Glu Asp Gly Val Glu Met Asp Thr Arg Ser Pro His Asp Glu Asp Pro	
	545 550 555	
5	cag gca gtg acg tat gcc gag gtg aaa cac tcc aga cct agg aga gaa Gln Ala Val Thr Tyr Ala Glu Val Lys His Ser Arg Pro Arg Arg Glu	2028
	560 565 570	
10	atg gcc tct cct cct tcc cca ctg tct ggg gaa ttc ctg gac aca aag Met Ala Ser Pro Pro Ser Pro Leu Ser Gly Glu Phe Leu Asp Thr Lys	2076
	575 580 585	
15	gac aga cag gcg gaa gag gac agg cag atg gac act gag gct gct gca Asp Arg Gln Ala Glu Glu Asp Arg Gln Met Asp Thr Glu Ala Ala Ala	2124
	590 595 600 605	
20	tct gaa gcc ccc cag gat gtg acc tac gcc cag ctg cac agc ttg acc Ser Glu Ala Pro Gln Asp Val Thr Tyr Ala Gln Leu His Ser Leu Thr	2172
	610 615 620	
25	ctt aga cgg aag gca act gag cct cct cca tcc cag gaa ggg ccc tct Leu Arg Arg Lys Ala Thr Glu Pro Pro Pro Ser Gln Glu Gly Pro Ser	2220
	625 630 635	
30	cca gct gtg ccc agc atc tac gcc act ctg gcc atc cac tag Pro Ala Val Pro Ser Ile Tyr Ala Thr Leu Ala Ile His	2262
	640 645 650	
35	cccagggggg gacgcagacc ccacactcca tggagtctgg aatgcatggg agctgcccc	2322
40	ccagtggaca ccattggacc ccacccagcc tggatctacc ccaggagact ctgggaactt	2382
45	ttaggggtca ctcaattctg cagtataaat aactaatgtc tctacaattt tgaaataaag	2442
50	caacagactt ctcaataatc aatgaagtag ctgagaaaac taagtcagaa agtgcattaa	2502
55	actgaatcac aatgtaaata ttacacatca agcgatgaaa ctggaaaact acaagccacg	2562
60	aatgaatgaa ttaggaaaga aaaaaagtag gaaatgaatg atcttggtt tcctataaga	2622
65	aatttagggc agggcacggt ggctcacgcc tgtaattcca gcactttggg aggccgaggc	2682
70	gggcagatca cgagttcagg agatcgagac catcttggcc aacatggtga aacctgtct	2742
75	ctcctaaaaa tacaaaaatt agctggatgt ggtggcagt cctgtaatcc cagctatttg	2802
80	ggaggctgag gcaggagaat cgcttgaacc agggagtcag aggtttcagt gagccaagat	2862
85	cgcaccactg ctctccagcc tggcgacaag caggtcgtct cgttccagtc gacggcccat	2922
90	<210> 2	
95	<211> 650	
100	<212> PRT	
105	<213> human	
110	<400> 2	
115	Met Thr Pro Ile Leu Thr Val Leu Ile Cys Leu Gly Leu Ser Leu Gly	
120	1 5 10 15	
125	Pro Arg Thr His Val Gln Ala Gly His Leu Pro Lys Pro Thr Leu Trp	
130	20 25 30	
135		
140		
145		
150		
155		
160		
165		
170		
175		
180		
185		
190		
195		
200		
205		
210		
215		
220		
225		
230		
235		
240		
245		
250		
255		
260		
265		
270		
275		
280		
285		
290		
295		
300		
305		
310		
315		
320		
325		
330		
335		
340		
345		
350		
355		
360		
365		
370		
375		
380		
385		
390		
395		
400		
405		
410		
415		
420		
425		
430		
435		
440		
445		
450		
455		
460		
465		
470		
475		
480		
485		
490		
495		
500		
505		
510		
515		
520		
525		
530		
535		
540		
545		
550		
555		
560		
565		
570		
575		
580		
585		
590		
595		
600		
605		
610		
615		
620		
625		
630		
635		
640		
645		
650		
655		
660		
665		
670		
675		
680		
685		
690		
695		
700		
705		
710		
715		
720		
725		
730		
735		
740		
745		
750		
755		
760		
765		
770		
775		
780		
785		
790		
795		
800		
805		
810		
815		
820		
825		
830		
835		
840		
845		
850		
855		
860		
865		
870		
875		
880		
885		
890		
895		
900		
905		
910		
915		
920		
925		
930		
935		
940		
945		
950		
955		
960		
965		
970		
975		
980		
985		
990		
995		

EP 1 199 362 A2

Ala Glu Pro Gly Ser Val Ile Thr Gln Gly Ser Pro Val Thr Leu Arg  
35 40 45

5 Cys Gln Gly Gly Gln Glu Thr Gln Glu Tyr Arg Leu Tyr Arg Glu Lys  
50 55 60

Lys Thr Ala Pro Trp Ile Thr Arg Ile Pro Gln Glu Leu Val Lys Lys  
65 70 75 80

10 Gly Gln Phe Pro Ile Pro Ser Ile Thr Trp Glu His Ala Gly Arg Tyr  
85 90 95

Arg Cys Tyr Tyr Gly Ser Asp Thr Ala Gly Arg Ser Glu Ser Ser Asp  
100 105 110

15 Pro Leu Glu Leu Val Val Thr Gly Ala Tyr Ile Lys Pro Thr Leu Ser  
115 120 125

Ala Gln Pro Ser Pro Val Val Asn Ser Gly Gly Asn Val Thr Leu Gln  
130 135 140

20 Cys Asp Ser Gln Val Ala Phe Asp Gly Phe Ile Leu Cys Lys Glu Gly  
145 150 155 160

Glu Asp Glu His Pro Gln Cys Leu Asn Ser Gln Pro His Ala Arg Gly  
165 170 175

25 Ser Ser Arg Ala Ile Phe Ser Val Gly Pro Val Ser Pro Ser Arg Arg  
180 185 190

Trp Trp Tyr Arg Cys Tyr Ala Tyr Asp Ser Asn Ser Pro Tyr Glu Trp  
195 200 205

30 Ser Leu Pro Ser Asp Leu Leu Glu Leu Leu Val Leu Gly Val Ser Lys  
210 215 220

Lys Pro Ser Leu Ser Val Gln Pro Gly Pro Ile Val Ala Pro Glu Glu  
225 230 235 240

35 Thr Leu Thr Leu Gln Cys Gly Ser Asp Ala Gly Tyr Asn Arg Phe Val  
245 250 255

Leu Tyr Lys Asp Gly Glu Arg Asp Phe Leu Gln Leu Ala Gly Ala Gln  
260 265 270

40 Pro Gln Ala Gly Leu Ser Gln Ala Asn Phe Thr Leu Gly Pro Val Ser  
275 280 285

Arg Ser Tyr Gly Gly Gln Tyr Arg Cys Tyr Gly Ala His Asn Leu Ser  
290 295 300

45 Ser Glu Trp Ser Ala Pro Ser Asp Pro Leu Asp Ile Leu Ile Ala Gly  
305 310 315 320

Gln Phe Tyr Asp Arg Val Ser Leu Ser Val Gln Pro Gly Pro Thr Val  
325 330 335

50 Ala Ser Gly Glu Asn Val Thr Leu Leu Cys Gln Ser Gln Gly Trp Met  
340 345 350

55

EP 1 199 362 A2

	Gln Thr Phe Leu Leu Thr Lys Glu Gly Ala Ala Asp Asp Pro Trp Arg	
	355	360 365
5	Leu Arg Ser Thr Tyr Gln Ser Gln Lys Tyr Gln Ala Glu Phe Pro Met	
	370	375 380
	Gly Pro Val Thr Ser Ala His Ala Gly Thr Tyr Arg Cys Tyr Gly Ser	
	385	390 395 400
10	Gln Ser Ser Lys Pro Tyr Leu Leu Thr His Pro Ser Asp Pro Leu Glu	
		405 410 415
	Leu Val Val Ser Gly Pro Ser Gly Gly Pro Ser Ser Pro Thr Thr Gly	
		420 425 430
15	Pro Thr Ser Thr Ser Gly Pro Glu Asp Gln Pro Leu Thr Pro Thr Gly	
		435 440 445
	Ser Asp Pro Gln Ser Gly Leu Gly Arg His Leu Gly Val Val Ile Gly	
		450 455 460
20	Ile Leu Val Ala Val Ile Leu Leu Leu Leu Leu Leu Leu Leu Phe	
		465 470 475 480
	Leu Ile Leu Arg His Arg Arg Gln Gly Lys His Trp Thr Ser Thr Gln	
		485 490 495
25	Arg Lys Ala Asp Phe Gln His Pro Ala Gly Ala Val Gly Pro Glu Pro	
		500 505 510
	Thr Asp Arg Gly Leu Gln Trp Arg Ser Ser Pro Ala Ala Asp Ala Gln	
		515 520 525
30	Glu Glu Asn Leu Tyr Ala Ala Val Lys His Thr Gln Pro Glu Asp Gly	
		530 535 540
	Val Glu Met Asp Thr Arg Ser Pro His Asp Glu Asp Pro Gln Ala Val	
		545 550 555 560
35	Thr Tyr Ala Glu Val Lys His Ser Arg Pro Arg Arg Glu Met Ala Ser	
		565 570 575
	Pro Pro Ser Pro Leu Ser Gly Glu Phe Leu Asp Thr Lys Asp Arg Gln	
		580 585 590
40	Ala Glu Glu Asp Arg Gln Met Asp Thr Glu Ala Ala Ala Ser Glu Ala	
		595 600 605
	Pro Gln Asp Val Thr Tyr Ala Gln Leu His Ser Leu Thr Leu Arg Arg	
		610 615 620
45	Lys Ala Thr Glu Pro Pro Pro Ser Gln Glu Gly Pro Ser Pro Ala Val	
		625 630 635 640
	Pro Ser Ile Tyr Ala Thr Leu Ala Ile His	
		645 650
50	<210> 3	
	<211> 2777	
	<212> DNA	
	<213> human	

55

5 <220>  
 <221> CDS  
 <222> (168)..(2126)  
  
 <400> 3  
 agctcagcct gggcgggcaca gccagatgcg agatgcgtct ctgctgatct gagtctgcct 60  
 10 gcagcatgga cctgggtctt ccctgaagca tctccagggc tggagggacg actgccatgc 120  
 accgaggggt catccatcca cagagcaggg cagtgggagg agacgcc atg acc ccc 176  
 Met Thr Pro  
 1  
 15 atc ctc acg gtc ctg atc tgt ctc ggg ctg agt ctg ggc ccc agg acc 224  
 Ile Leu Thr Val Leu Ile Cys Leu Gly Leu Ser Leu Gly Pro Arg Thr  
 5 10 15  
 20 cac gtg cag gca ggg cac ctc ccc aag ccc acc ctc tgg gct gaa cca 272  
 His Val Gln Ala Gly His Leu Pro Lys Pro Thr Leu Trp Ala Glu Pro  
 20 25 30 35  
 ggc tct gtg atc acc cag ggg agt cct gtg acc ctc agg tgt cag ggg 320  
 Gly Ser Val Ile Thr Gln Gly Ser Pro Val Thr Leu Arg Cys Gln Gly  
 40 45 50  
 25 ggc cag gag acc cag gag tac cgt cta tat aga gaa aag aaa aca gca 368  
 Gly Gln Glu Thr Gln Glu Tyr Arg Leu Tyr Arg Glu Lys Lys Thr Ala  
 55 60 65  
 30 ctc tgg att aca cgg atc cca cag gag ctt gtg aag aag ggc cag ttc 416  
 Leu Trp Ile Thr Arg Ile Pro Gln Glu Leu Val Lys Lys Gly Gln Phe  
 70 75 80  
 ccc atc cca tcc atc acc tgg gaa cat gca ggg cgg tat cgc tgt tac 464  
 Pro Ile Pro Ser Ile Thr Trp Glu His Ala Gly Arg Tyr Arg Cys Tyr  
 85 90 95  
 35 tat ggt agc gac act gca ggc cgc tca gag agc agt gac ccc ctg gag 512  
 Tyr Gly Ser Asp Thr Ala Gly Arg Ser Glu Ser Ser Asp Pro Leu Glu  
 100 105 110 115  
 40 ctg gtg gtg aca gga gcc tac atc aaa ccc acc ctc tca gcc cag ccc 560  
 Leu Val Val Thr Gly Ala Tyr Ile Lys Pro Thr Leu Ser Ala Gln Pro  
 120 125 130  
 agc ccc gtg gtg aac tca gga ggg aat gta atc ctc cag tgt gac tca 608  
 Ser Pro Val Val Asn Ser Gly Gly Asn Val Ile Leu Gln Cys Asp Ser  
 135 140 145  
 45 cag gtg gca ttt gat ggc ttc agt ctg tgt aag gaa gga gaa gat gaa 656  
 Gln Val Ala Phe Asp Gly Phe Ser Leu Cys Lys Glu Gly Glu Asp Glu  
 150 155 160  
 50 cac cca caa tgc ctg aac tcc cag ccc cat gcc cgt ggg tgc tcc cgc 704  
 His Pro Gln Cys Leu Asn Ser Gln Pro His Ala Arg Gly Ser Ser Arg  
 165 170 175  
 gcc atc ttc tcc gtg ggc ccc gtg agc ccg agt cgc agg tgg tgg tac 752

EP 1 199 362 A2

	Ala	Ile	Phe	Ser	Val	Gly	Pro	Val	Ser	Pro	Ser	Arg	Arg	Trp	Trp	Tyr	
	180					185					190					195	
5	agg	tgc	tat	gct	tat	gac	tcg	aac	tct	ccc	tat	gag	tgg	tct	cta	ccc	800
	Arg	Cys	Tyr	Ala	Tyr	Asp	Ser	Asn	Ser	Pro	Tyr	Glu	Trp	Ser	Leu	Pro	
					200					205					210		
	agt	gat	ctc	ctg	gag	ctc	ctg	gtc	cta	ggc	gtt	tct	aag	aag	cca	tca	848
	Ser	Asp	Leu	Leu	Glu	Leu	Leu	Val	Leu	Gly	Val	Ser	Lys	Lys	Pro	Ser	
10					215				220					225			
	ctc	tca	gtg	cag	cca	ggc	cct	atc	gtg	gcc	cct	gag	gag	acc	ctg	act	896
	Leu	Ser	Val	Gln	Pro	Gly	Pro	Ile	Val	Ala	Pro	Glu	Glu	Thr	Leu	Thr	
					230			235						240			
15	ctg	cag	tgt	ggc	tct	gat	gct	ggc	tac	aac	aga	ttt	gtt	ctg	tat	aag	944
	Leu	Gln	Cys	Gly	Ser	Asp	Ala	Gly	Tyr	Asn	Arg	Phe	Val	Leu	Tyr	Lys	
		245					250					255					
	gac	ggg	gaa	cgt	gac	ttc	ctt	cag	ctc	gct	ggc	gca	cag	ccc	cag	gct	992
	Asp	Gly	Glu	Arg	Asp	Phe	Leu	Gln	Leu	Ala	Gly	Ala	Gln	Pro	Gln	Ala	
20	260					265					270					275	
	ggg	ctc	tcc	cag	gcc	aac	ttc	acc	ctg	ggc	cct	gtg	agc	cgc	tcc	tac	1040
	Gly	Leu	Ser	Gln	Ala	Asn	Phe	Thr	Leu	Gly	Pro	Val	Ser	Arg	Ser	Tyr	
					280					285					290		
25	ggg	ggc	cag	tac	aga	tgc	tac	ggc	gca	cac	aac	ctc	tcc	tcc	gag	tgg	1088
	Gly	Gly	Gln	Tyr	Arg	Cys	Tyr	Gly	Ala	His	Asn	Leu	Ser	Ser	Glu	Trp	
				295					300					305			
	tcg	gcc	ccc	agt	gac	ccc	ctg	gac	atc	ctg	atc	gca	gga	cag	ttc	tat	1136
	Ser	Ala	Pro	Ser	Asp	Pro	Leu	Asp	Ile	Leu	Ile	Ala	Gly	Gln	Phe	Tyr	
30			310					315					320				
	gac	aga	gtc	tcc	ctc	tcg	gtg	cag	ccg	ggc	ccc	acg	gtg	gcc	tca	gga	1184
	Asp	Arg	Val	Ser	Leu	Ser	Val	Gln	Pro	Gly	Pro	Thr	Val	Ala	Ser	Gly	
		325					330					335					
35	gag	aac	gtg	acc	ctg	ctg	tgt	cag	tca	cag	gga	tgg	atg	caa	act	ttc	1232
	Glu	Asn	Val	Thr	Leu	Leu	Cys	Gln	Ser	Gln	Gly	Trp	Met	Gln	Thr	Phe	
	340					345					350					355	
	ctt	ctg	acc	aag	gag	ggg	gca	gct	gat	gac	cca	tgg	cgt	cta	aga	tca	1280
	Leu	Leu	Thr	Lys	Glu	Gly	Ala	Ala	Asp	Asp	Pro	Trp	Arg	Leu	Arg	Ser	
40					360					365					370		
	acg	tac	caa	tct	caa	aaa	tac	cag	gct	gaa	ttc	ccc	atg	ggc	cct	gtg	1328
	Thr	Tyr	Gln	Ser	Gln	Lys	Tyr	Gln	Ala	Glu	Phe	Pro	Met	Gly	Pro	Val	
				375					380					385			
45	acc	tca	gcc	cat	gcg	ggg	acc	tac	agg	tgc	tac	ggc	tca	cag	agc	tcc	1376
	Thr	Ser	Ala	His	Ala	Gly	Thr	Tyr	Arg	Cys	Tyr	Gly	Ser	Gln	Ser	Ser	
			390					395					400				
	aaa	ccc	tac	ctg	ctg	act	cac	ccc	agt	gac	ccc	ctg	gag	ctc	gtg	gtc	1424
	Lys	Pro	Tyr	Leu	Leu	Thr	His	Pro	Ser	Asp	Pro	Leu	Glu	Leu	Val	Val	
50		405					410					415					
	tca	gga	ccg	tct	ggg	ggc	ccc	agc	tcc	ccg	aca	aca	ggc	ccc	acc	tcc	1472
	Ser	Gly	Pro	Ser	Gly	Gly	Pro	Ser	Ser	Pro	Thr	Thr	Gly	Pro	Thr	Ser	

55

EP 1 199 362 A2

	420		425		430		435	
5	aca tct gca ggc cct gag gac cag ccc ctc acc ccc acc ggg tcg gat							1520
	Thr Ser Ala Gly Pro Glu Asp Gln Pro Leu Thr Pro Thr Gly Ser Asp							
		440			445		450	
	ccc cag agt ggt ctg gga agg cac ctg ggg gtt gtg atc ggc atc ttg							1568
	Pro Gln Ser Gly Leu Gly Arg His Leu Gly Val Val Ile Gly Ile Leu							
		455			460		465	
10	gtg gcc gtc atc cta ctg ctc ctc ctc ctc ctc ctc ttc ctc atc							1616
	Val Ala Val Ile Leu Leu Leu Leu Leu Leu Leu Leu Phe Leu Ile							
		470			475		480	
	ctc cga cat cga cgt cag ggc aaa cac tgg aca tcg acc cag aga aag							1664
15	Leu Arg His Arg Arg Gln Gly Lys His Trp Thr Ser Thr Gln Arg Lys							
		485			490		495	
	gct gat ttc caa cat cct gca ggg gct gtg ggg cca gag ccc aca gac							1712
	Ala Asp Phe Gln His Pro Ala Gly Ala Val Gly Pro Glu Pro Thr Asp							
	500			505		510		515
20	aga ggc ctg cag tgg agg tcc agc cca gct gcc gat gcc cag gaa gaa							1760
	Arg Gly Leu Gln Trp Arg Ser Ser Pro Ala Ala Asp Ala Gln Glu Glu							
		520			525		530	
	aac ctc tat gct gcc gtg aag cac aca cag cct gag gat ggg gtg gag							1808
25	Asn Leu Tyr Ala Ala Val Lys His Thr Gln Pro Glu Asp Gly Val Glu							
		535			540		545	
	atg gac act cgg cag agc cca cac gat gaa gac ccc cag gca gtg acg							1856
	Met Asp Thr Arg Gln Ser Pro His Asp Glu Asp Pro Gln Ala Val Thr							
		550			555		560	
30	tat gcc gag gtg aaa cac tcc aga cct agg aga gaa atg gcc tct cct							1904
	Tyr Ala Glu Val Lys His Ser Arg Pro Arg Arg Glu Met Ala Ser Pro							
		565			570		575	
	cct tcc cca ctg tct ggg gaa ttc ctg gac aca aag gac aga cag gcg							1952
35	Pro Ser Pro Leu Ser Gly Glu Phe Leu Asp Thr Lys Asp Arg Gln Ala							
		580			585		590	595
	gaa gag gac agg cag atg gac act gag gct gct gca tct gaa gcc ccc							2000
	Glu Glu Asp Arg Gln Met Asp Thr Glu Ala Ala Ala Ser Glu Ala Pro							
		600			605		610	
40	cag gat gtg acc tac gcc cag ctg cac agc ttg acc ctc aga cgg gag							2048
	Gln Asp Val Thr Tyr Ala Gln Leu His Ser Leu Thr Leu Arg Arg Glu							
		615			620		625	
	gca act gag cct cct cca tcc cag gaa ggg ccc tct cca gct gtg ccc							2096
45	Ala Thr Glu Pro Pro Pro Ser Gln Glu Gly Pro Ser Pro Ala Val Pro							
		630			635		640	
	agc atc tac gcc act ctg gcc atc cac tag cccagggggg gacgcagacc							2146
	Ser Ile Tyr Ala Thr Leu Ala Ile His							
50		645			650			
	ccacactcca tggagtctgg aatgcatggg agctgcccc ccagtggaca ccattggacc							2206
	ccaccagcc tggatctacc ccaggagact ctgggaactt ttaggggtca ctcaattctg							2266
55								



EP 1 199 362 A2

cagtataaat aactaatgtc tctacaattt tgaaataaag caatagactt ctcaataatc 2326  
 aatgaagtag ctgagaaaac taagtcagaa agtgcattaa actgaatcac aatgtaaata 2386  
 5 ttacacatca agcgatgaaa ctggaaaact acaagccacg aatgaatgaa ttaggaaaga 2446  
 aaaaaagtag gaaatgaatg atcttggctt tcctataaga aatttagggc agggcacggt 2506  
 ggctcacgcc tgtaattcca gcactttggg aggccgaggc gggcagatca cgagttcagg 2566  
 10 agatcgagac catcttggcc aacatggtga aaccctgtct ctccataaaa tacaaaaatt 2626  
 agctggatgt ggtggcagtg cctgtaatcc cagctatttg ggaggctgag gcaggagaat 2686  
 cgcttgaacc agggagtcag aggtttcagt gagccaagat cgcaccactg ctctccagcc 2746  
 15 tggcgacaga gggagactcc atctcaaatt a 2777  
  
 <210> 4  
 <211> 652  
 20 <212> PRT  
 <213> human  
  
 <400> 4  
 Met Thr Pro Ile Leu Thr Val Leu Ile Cys Leu Gly Leu Ser Leu Gly  
 1 5 10 15  
 25 Pro Arg Thr His Val Gln Ala Gly His Leu Pro Lys Pro Thr Leu Trp  
 20 25 30  
 Ala Glu Pro Gly Ser Val Ile Thr Gln Gly Ser Pro Val Thr Leu Arg  
 35 35 40 45  
 Cys Gln Gly Gly Gln Glu Thr Gln Glu Tyr Arg Leu Tyr Arg Glu Lys  
 50 55 60  
 Lys Thr Ala Leu Trp Ile Thr Arg Ile Pro Gln Glu Leu Val Lys Lys  
 65 70 75 80  
 35 Gly Gln Phe Pro Ile Pro Ser Ile Thr Trp Glu His Ala Gly Arg Tyr  
 85 90 95  
 Arg Cys Tyr Tyr Gly Ser Asp Thr Ala Gly Arg Ser Glu Ser Ser Asp  
 100 105 110  
 40 Pro Leu Glu Leu Val Val Thr Gly Ala Tyr Ile Lys Pro Thr Leu Ser  
 115 120 125  
 Ala Gln Pro Ser Pro Val Val Asn Ser Gly Gly Asn Val Ile Leu Gln  
 130 135 140  
 45 Cys Asp Ser Gln Val Ala Phe Asp Gly Phe Ser Leu Cys Lys Glu Gly  
 145 150 155 160  
 Glu Asp Glu His Pro Gln Cys Leu Asn Ser Gln Pro His Ala Arg Gly  
 165 170 175  
 50 Ser Ser Arg Ala Ile Phe Ser Val Gly Pro Val Ser Pro Ser Arg Arg  
 180 185 190  
 55

EP 1 199 362 A2

Trp Trp Tyr Arg Cys Tyr Ala Tyr Asp Ser Asn Ser Pro Tyr Glu Trp  
 195 200 205  
 5 Ser Leu Pro Ser Asp Leu Leu Glu Leu Leu Val Leu Gly Val Ser Lys  
 210 215 220  
 Lys Pro Ser Leu Ser Val Gln Pro Gly Pro Ile Val Ala Pro Glu Glu  
 225 230 235 240  
 10 Thr Leu Thr Leu Gln Cys Gly Ser Asp Ala Gly Tyr Asn Arg Phe Val  
 245 250 255  
 Leu Tyr Lys Asp Gly Glu Arg Asp Phe Leu Gln Leu Ala Gly Ala Gln  
 260 265 270  
 15 Pro Gln Ala Gly Leu Ser Gln Ala Asn Phe Thr Leu Gly Pro Val Ser  
 275 280 285  
 Arg Ser Tyr Gly Gly Gln Tyr Arg Cys Tyr Gly Ala His Asn Leu Ser  
 290 295 300  
 20 Ser Glu Trp Ser Ala Pro Ser Asp Pro Leu Asp Ile Leu Ile Ala Gly  
 305 310 315 320  
 Gln Phe Tyr Asp Arg Val Ser Leu Ser Val Gln Pro Gly Pro Thr Val  
 325 330 335  
 25 Ala Ser Gly Glu Asn Val Thr Leu Leu Cys Gln Ser Gln Gly Trp Met  
 340 345 350  
 Gln Thr Phe Leu Leu Thr Lys Glu Gly Ala Ala Asp Asp Pro Trp Arg  
 355 360 365  
 30 Leu Arg Ser Thr Tyr Gln Ser Gln Lys Tyr Gln Ala Glu Phe Pro Met  
 370 375 380  
 Gly Pro Val Thr Ser Ala His Ala Gly Thr Tyr Arg Cys Tyr Gly Ser  
 385 390 395 400  
 35 Gln Ser Ser Lys Pro Tyr Leu Leu Thr His Pro Ser Asp Pro Leu Glu  
 405 410 415  
 Leu Val Val Ser Gly Pro Ser Gly Gly Pro Ser Ser Pro Thr Thr Gly  
 420 425 430  
 40 Pro Thr Ser Thr Ser Ala Gly Pro Glu Asp Gln Pro Leu Thr Pro Thr  
 435 440 445  
 Gly Ser Asp Pro Gln Ser Gly Leu Gly Arg His Leu Gly Val Val Ile  
 450 455 460  
 45 Gly Ile Leu Val Ala Val Ile Leu Leu Leu Leu Leu Leu Leu Leu  
 465 470 475 480  
 Phe Leu Ile Leu Arg His Arg Arg Gln Gly Lys His Trp Thr Ser Thr  
 485 490 495  
 50 Gln Arg Lys Ala Asp Phe Gln His Pro Ala Gly Ala Val Gly Pro Glu  
 500 505 510

55

EP 1 199 362 A2

	Pro Thr Asp Arg Gly Leu Gln Trp Arg Ser Ser Pro Ala Ala Asp Ala	
	515 520 525	
5	Gln Glu Glu Asn Leu Tyr Ala Ala Val Lys His Thr Gln Pro Glu Asp	
	530 535 540	
	Gly Val Glu Met Asp Thr Arg Gln Ser Pro His Asp Glu Asp Pro Gln	
	545 550 555 560	
10	Ala Val Thr Tyr Ala Glu Val Lys His Ser Arg Pro Arg Arg Glu Met	
	565 570 575	
	Ala Ser Pro Pro Ser Pro Leu Ser Gly Glu Phe Leu Asp Thr Lys Asp	
	580 585 590	
15	Arg Gln Ala Glu Glu Asp Arg Gln Met Asp Thr Glu Ala Ala Ala Ser	
	595 600 605	
	Glu Ala Pro Gln Asp Val Thr Tyr Ala Gln Leu His Ser Leu Thr Leu	
	610 615 620	
20	Arg Arg Glu Ala Thr Glu Pro Pro Pro Ser Gln Glu Gly Pro Ser Pro	
	625 630 635 640	
	Ala Val Pro Ser Ile Tyr Ala Thr Leu Ala Ile His	
	645 650	
25	<210> 5	
	<211> 30	
	<212> DNA	
	<213> human	
30	<400> 5	
	tatgtcgacc atgacccccca tcttcacggt	30
35	<210> 6	
	<211> 52	
	<212> DNA	
	<213> human	
	<400> 6	
	tatgggctct gctccaggag aagatcttcc ttctataacc cccaggtgcc tt	52
40	<210> 7	
	<211> 1605	
	<212> DNA	
	<213> human	
45	<220>	
	<221> CDS	
	<222> (93)..(1412)	
	<400> 7	
	gagcctccaa gtgtccacac cctgtgtgtc ctctgtcctg ccagcaccga gggctcatcc	60
50	atccacagag cagtgcagtg ggaggagacg cc atg acc ccc atc ctc acg gtc	113
	Met Thr Pro Ile Leu Thr Val	
	1 5	
55		

EP 1 199 362 A2

	ctg atc tgt ctc ggg ctg agc ctg gac ccc agg acc cac gtg cag gca	161
	Leu Ile Cys Leu Gly Leu Ser Leu Asp Pro Arg Thr His Val Gln Ala	
	10 15 20	
5	ggg ccc ctc ccc aag ccc acc ctc tgg gct gag cca ggc tct gtg atc	209
	Gly Pro Leu Pro Lys Pro Thr Leu Trp Ala Glu Pro Gly Ser Val Ile	
	25 30 35	
	acc caa ggg agt cct gtg acc ctc agg tgt cag ggg agc ctg gag acg	257
10	Thr Gln Gly Ser Pro Val Thr Leu Arg Cys Gln Gly Ser Leu Glu Thr	
	40 45 50 55	
	cag gag tac cat cta tat aga gaa aag aaa aca gca ctc tgg att aca	305
	Gln Glu Tyr His Leu Tyr Arg Glu Lys Thr Ala Leu Trp Ile Thr	
	60 65 70	
15	cgg atc cca cag gag ctt gtg aag aag ggc cag ttc ccc atc cta tcc	353
	Arg Ile Pro Gln Glu Leu Val Lys Lys Gly Gln Phe Pro Ile Leu Ser	
	75 80 85	
	atc acc tgg gaa cat gca ggg cgg tat tgc tgt atc tat ggc agc cac	401
20	Ile Thr Trp Glu His Ala Gly Arg Tyr Cys Cys Ile Tyr Gly Ser His	
	90 95 100	
	act gca ggc ctc tca gag agc agt gac ccc ctg gag ctg gtg gtg aca	449
	Thr Ala Gly Leu Ser Glu Ser Ser Asp Pro Leu Glu Leu Val Val Thr	
	105 110 115	
25	gga gcc tac agc aaa ccc acc ctc tca gct ctg ccc agc cct gtg gtg	497
	Gly Ala Tyr Ser Lys Pro Thr Leu Ser Ala Leu Pro Ser Pro Val Val	
	120 125 130 135	
	acc tca gga agg aat gtg acc atc cag tgt gac tca cag gtg gca ttt	545
30	Thr Ser Gly Arg Asn Val Thr Ile Gln Cys Asp Ser Gln Val Ala Phe	
	140 145 150	
	gat ggc ttc att ctg tgt aag gaa gga gaa gat gaa cac cca caa tgc	593
	Asp Gly Phe Ile Leu Cys Lys Glu Gly Glu Asp Glu His Pro Gln Cys	
	155 160 165	
35	ctg aac tcc cat tcc cat gcc cgt ggg tca tcc cgg gcc atc ttc tcc	641
	Leu Asn Ser His Ser His Ala Arg Gly Ser Ser Arg Ala Ile Phe Ser	
	170 175 180	
	gtg ggc ccc gtg agc cca agt cgc agg tgg tgc tac agg tgc tat ggt	689
40	Val Gly Pro Val Ser Pro Ser Arg Arg Trp Ser Tyr Arg Cys Tyr Gly	
	185 190 195	
	tat gac tcg cgc gct ccc tat gtg tgg tct cta ccc agt gat ctc ctg	737
	Tyr Asp Ser Arg Ala Pro Tyr Val Trp Ser Leu Pro Ser Asp Leu Leu	
	200 205 210 215	
45	ggg ctc ctg gtc cca ggt gtt tct aag aag cca tca ctc tca gtg cag	785
	Gly Leu Leu Val Pro Gly Val Ser Lys Lys Pro Ser Leu Ser Val Gln	
	220 225 230	
	ccg ggt cct gtc gtg gcc cct ggg gag aag ctg acc ttc cag tgt ggc	833
50	Pro Gly Pro Val Val Ala Pro Gly Glu Lys Leu Thr Phe Gln Cys Gly	
	235 240 245	
	tct gat gcc ggc tac gac aga ttt gtt ctg tac aag gag tgg gga cgt	881

55

EP 1 199 362 A2

	Ser	Asp	Ala	Gly	Tyr	Asp	Arg	Phe	Val	Leu	Tyr	Lys	Glu	Trp	Gly	Arg	
			250					255					260				
5	gac	ttc	ctc	cag	cgc	cct	ggc	cgg	cag	ccc	cag	gct	ggg	ctc	tcc	cag	929
	Asp	Phe	Leu	Gln	Arg	Pro	Gly	Arg	Gln	Pro	Gln	Ala	Gly	Leu	Ser	Gln	
		265					270					275					
	gcc	aac	ttc	acc	ctg	ggc	cct	gtg	agc	cgc	tcc	tac	ggg	ggc	cag	tac	977
10	Ala	Asn	Phe	Thr	Leu	Gly	Pro	Val	Ser	Arg	Ser	Tyr	Gly	Gly	Gln	Tyr	
	280					285					290					295	
	aca	tgc	tcc	ggt	gca	tac	aac	ctc	tcc	tcc	gag	tgg	tcg	gcc	ccc	agc	1025
	Thr	Cys	Ser	Gly	Ala	Tyr	Asn	Leu	Ser	Ser	Glu	Trp	Ser	Ala	Pro	Ser	
					300					305						310	
15	gac	ccc	ctg	gac	atc	ctg	atc	aca	gga	cag	atc	cgt	gcc	aga	ccc	ttc	1073
	Asp	Pro	Leu	Asp	Ile	Leu	Ile	Thr	Gly	Gln	Ile	Arg	Ala	Arg	Pro	Phe	
				315					320					325			
	ctc	tcc	gtg	cgg	ccg	ggc	ccc	aca	gtg	gcc	tca	gga	gag	aac	gtg	acc	1121
20	Leu	Ser	Val	Arg	Pro	Gly	Pro	Thr	Val	Ala	Ser	Gly	Glu	Asn	Val	Thr	
			330					335					340				
	ctg	ctg	tgt	cag	tca	cag	gga	ggg	atg	cac	act	ttc	ctt	ttg	acc	aag	1169
	Leu	Leu	Cys	Gln	Ser	Gln	Gly	Gly	Met	His	Thr	Phe	Leu	Leu	Thr	Lys	
			345				350					355					
25	gag	ggg	gca	gct	gat	tcc	ccg	ctg	cgt	cta	aaa	tca	aag	cgc	caa	tct	1217
	Glu	Gly	Ala	Ala	Asp	Ser	Pro	Leu	Arg	Leu	Lys	Ser	Lys	Arg	Gln	Ser	
	360					365					370					375	
	cat	aag	tac	cag	gct	gaa	ttc	ccc	atg	agt	cct	gtg	acc	tcg	gcc	cac	1265
30	His	Lys	Tyr	Gln	Ala	Glu	Phe	Pro	Met	Ser	Pro	Val	Thr	Ser	Ala	His	
				380					385						390		
	gcg	ggg	acc	tac	agg	tgc	tac	ggc	tca	ctc	agc	tcc	aac	ccc	tac	ctg	1313
	Ala	Gly	Thr	Tyr	Arg	Cys	Tyr	Gly	Ser	Leu	Ser	Ser	Asn	Pro	Tyr	Leu	
			395					400					405				
35	ctg	act	cac	ccc	agt	gac	ccc	ctg	gag	ctc	gtg	gtc	tca	gga	gca	gct	1361
	Leu	Thr	His	Pro	Ser	Asp	Pro	Leu	Glu	Leu	Val	Val	Ser	Gly	Ala	Ala	
			410				415						420				
	gag	acc	ctc	agc	cca	cca	caa	aac	aag	tcc	gac	tcc	aag	gct	ggt	gag	1409
40	Glu	Thr	Leu	Ser	Pro	Pro	Gln	Asn	Lys	Ser	Asp	Ser	Lys	Ala	Gly	Glu	
		425					430				435						
	tga	ggagatgctt	gccgtgatga	cgtggggcac	agagggtcag	gtcctgtcaa											1462
	gaggagctgg	gtgtcctggg	tggacatttg	aagaattata	ttcattccaa	cttgaagaat											1522
45	tattcaacac	ctttaacaat	gtatatgtga	agtactttat	tctttcatat	tttaaaaaata											1582
	aaagataaatt	atccatgaga	aaa														1605

<210> 8

55

EP 1 199 362 A2

<211> 439  
<212> PRT  
<213> human

5  
<400> 8  
Met Thr Pro Ile Leu Thr Val Leu Ile Cys Leu Gly Leu Ser Leu Asp  
1 5 10 15  
Pro Arg Thr His Val Gln Ala Gly Pro Leu Pro Lys Pro Thr Leu Trp  
20 25 30  
10 Ala Glu Pro Gly Ser Val Ile Thr Gln Gly Ser Pro Val Thr Leu Arg  
35 40 45  
Cys Gln Gly Ser Leu Glu Thr Gln Glu Tyr His Leu Tyr Arg Glu Lys  
50 55 60  
15 Lys Thr Ala Leu Trp Ile Thr Arg Ile Pro Gln Glu Leu Val Lys Lys  
65 70 75 80  
Gly Gln Phe Pro Ile Leu Ser Ile Thr Trp Glu His Ala Gly Arg Tyr  
85 90 95  
20 Cys Cys Ile Tyr Gly Ser His Thr Ala Gly Leu Ser Glu Ser Ser Asp  
100 105 110  
Pro Leu Glu Leu Val Val Thr Gly Ala Tyr Ser Lys Pro Thr Leu Ser  
115 120 125  
25 Ala Leu Pro Ser Pro Val Val Thr Ser Gly Arg Asn Val Thr Ile Gln  
130 135 140  
Cys Asp Ser Gln Val Ala Phe Asp Gly Phe Ile Leu Cys Lys Glu Gly  
145 150 155 160  
30 Glu Asp Glu His Pro Gln Cys Leu Asn Ser His Ser His Ala Arg Gly  
165 170 175  
Ser Ser Arg Ala Ile Phe Ser Val Gly Pro Val Ser Pro Ser Arg Arg  
180 185 190  
35 Trp Ser Tyr Arg Cys Tyr Gly Tyr Asp Ser Arg Ala Pro Tyr Val Trp  
195 200 205  
Ser Leu Pro Ser Asp Leu Leu Gly Leu Leu Val Pro Gly Val Ser Lys  
210 215 220  
40 Lys Pro Ser Leu Ser Val Gln Pro Gly Pro Val Val Ala Pro Gly Glu  
225 230 235 240  
Lys Leu Thr Phe Gln Cys Gly Ser Asp Ala Gly Tyr Asp Arg Phe Val  
245 250 255  
45 Leu Tyr Lys Glu Trp Gly Arg Asp Phe Leu Gln Arg Pro Gly Arg Gln  
260 265 270  
Pro Gln Ala Gly Leu Ser Gln Ala Asn Phe Thr Leu Gly Pro Val Ser  
275 280 285  
50 Arg Ser Tyr Gly Gly Gln Tyr Thr Cys Ser Gly Ala Tyr Asn Leu Ser  
290 295 300

55

EP 1 199 362 A2

Ser Glu Trp Ser Ala Pro Ser Asp Pro Leu Asp Ile Leu Ile Thr Gly  
305 310 315 320

5 Gln Ile Arg Ala Arg Pro Phe Leu Ser Val Arg Pro Gly Pro Thr Val  
325 330 335

Ala Ser Gly Glu Asn Val Thr Leu Leu Cys Gln Ser Gln Gly Gly Met  
340 345 350

10 His Thr Phe Leu Leu Thr Lys Glu Gly Ala Ala Asp Ser Pro Leu Arg  
355 360 365

Leu Lys Ser Lys Arg Gln Ser His Lys Tyr Gln Ala Glu Phe Pro Met  
370 375 380

15 Ser Pro Val Thr Ser Ala His Ala Gly Thr Tyr Arg Cys Tyr Gly Ser  
385 390 395 400

Leu Ser Ser Asn Pro Tyr Leu Leu Thr His Pro Ser Asp Pro Leu Glu  
405 410 415

20 Leu Val Val Ser Gly Ala Ala Glu Thr Leu Ser Pro Pro Gln Asn Lys  
420 425 430

Ser Asp Ser Lys Ala Gly Glu  
435

25 <210> 9  
<211> 2221  
<212> DNA  
<213> human

30 <220>  
<221> CDS  
<222> (184)..(1980)

<400> 9  
gctcactgcc acacgcagct cagcctgggc ggcacagcca gatgcgagat gcgtctctgc 60

35 tgatctgagt ctgcctgcag catggacctg ggtcttcctt gaagcatctc cagggctgga 120  
gggacgactg ccatgcaccg agggctcatc catccgcaga gcagggcagt gggaggagac 180

40 gcc atg acc ccc atc gtc aca gtc ctg atc tgt ctc ggg ctg agt ctg 228  
Met Thr Pro Ile Val Thr Val Leu Ile Cys Leu Gly Leu Ser Leu  
1 5 10 15

ggc ccc agg acc cac gtg cag aca ggg acc atc ccc aag ccc acc ctg 276  
Gly Pro Arg Thr His Val Gln Thr Gly Thr Ile Pro Lys Pro Thr Leu  
20 25 30

45 tgg gct gag cca gac tct gtg atc acc cag ggg agt ccc gtc acc ctc 324  
Trp Ala Glu Pro Asp Ser Val Ile Thr Gln Gly Ser Pro Val Thr Leu  
35 40 45

50 agt tgt cag ggg agc ctt gaa gcc cag gag tac cgt cta tat agg gag 372  
Ser Cys Gln Gly Ser Leu Glu Ala Gln Glu Tyr Arg Leu Tyr Arg Glu  
50 55 60

aaa aaa tca gca tct tgg att aca cgg ata cga cca gag ctt gtg aag 420

55



EP 1 199 362 A2

	Lys	Lys	Ser	Ala	Ser	Trp	Ile	Thr	Arg	Ile	Arg	Pro	Glu	Leu	Val	Lys	
	65						70					75					
5	aac	ggc	cag	ttc	cac	atc	cca	tcc	atc	acc	tgg	gaa	cac	aca	ggg	cga	468
	Asn	Gly	Gln	Phe	His	Ile	Pro	Ser	Ile	Thr	Trp	Glu	His	Thr	Gly	Arg	
	80					85					90					95	
	tat	ggc	tgt	cag	tat	tac	agc	cgc	gct	cgg	tgg	tct	gag	ctc	agt	gac	516
	Tyr	Gly	Cys	Gln	Tyr	Tyr	Ser	Arg	Ala	Arg	Trp	Ser	Glu	Leu	Ser	Asp	
10					100					105					110		
	ccc	ctg	gtg	ctg	gtg	atg	aca	gga	gcc	tac	cca	aaa	ccc	acc	ctc	tca	564
	Pro	Leu	Val	Leu	Val	Met	Thr	Gly	Ala	Tyr	Pro	Lys	Pro	Thr	Leu	Ser	
				115					120					125			
15	gcc	cag	ccc	agc	cct	gtg	gtg	acc	tca	gga	gga	agg	gtg	acc	ctc	cag	612
	Ala	Gln	Pro	Ser	Pro	Val	Val	Thr	Ser	Gly	Gly	Arg	Val	Thr	Leu	Gln	
			130					135					140				
	tgt	gag	tca	cag	gtg	gca	ttt	ggc	ggc	ttc	att	ctg	tgt	aag	gaa	gga	660
	Cys	Glu	Ser	Gln	Val	Ala	Phe	Gly	Gly	Phe	Ile	Leu	Cys	Lys	Glu	Gly	
20		145					150					155					
	gaa	gat	gaa	cac	cca	caa	tgc	ctg	aac	tcc	cag	ccc	cat	gcc	cgt	ggg	708
	Glu	Asp	Glu	His	Pro	Gln	Cys	Leu	Asn	Ser	Gln	Pro	His	Ala	Arg	Gly	
	160					165					170					175	
25	tcg	tcc	cgc	gcc	atc	ttc	tcc	gtg	ggc	ccc	gtg	agc	ccg	aat	cgc	agg	756
	Ser	Ser	Arg	Ala	Ile	Phe	Ser	Val	Gly	Pro	Val	Ser	Pro	Asn	Arg	Arg	
					180					185					190		
	tgg	tcg	cac	agg	tgc	tat	ggt	tat	gac	ttg	aac	tct	ccc	tat	gtg	tgg	804
	Trp	Ser	His	Arg	Cys	Tyr	Gly	Tyr	Asp	Leu	Asn	Ser	Pro	Tyr	Val	Trp	
30				195					200					205			
	tct	tca	ccc	agt	gat	ctc	ctg	gag	ctc	ctg	gtc	cca	ggt	gtt	tct	aag	852
	Ser	Ser	Pro	Ser	Asp	Leu	Leu	Leu	Leu	Val	Pro	Gly	Val	Ser	Lys		
			210				215					220					
35	aag	cca	tca	ctc	tca	gtg	cag	ccg	ggt	cct	gtc	gtg	gcc	cct	ggg	gaa	900
	Lys	Pro	Ser	Leu	Ser	Val	Gln	Pro	Gly	Pro	Val	Val	Ala	Pro	Gly	Glu	
		225					230					235					
	agc	ctg	acc	ctc	cag	tgt	gtc	tct	gat	gtc	ggc	tat	gac	aga	ttt	gtt	948
	Ser	Leu	Thr	Leu	Gln	Cys	Val	Ser	Asp	Val	Gly	Tyr	Asp	Arg	Phe	Val	
40	240				245					250					255		
	ctg	tac	aag	gag	ggg	gaa	cgt	gac	ctt	cgc	cag	ctc	cct	ggc	cgg	cag	996
	Leu	Tyr	Lys	Glu	Gly	Glu	Arg	Asp	Leu	Arg	Gln	Leu	Pro	Gly	Arg	Gln	
				260					265						270		
45	ccc	cag	gct	ggg	ctc	tcc	cag	gcc	aac	ttc	acc	ctg	ggc	cct	gtg	agc	1044
	Pro	Gln	Ala	Gly	Leu	Ser	Gln	Ala	Asn	Phe	Thr	Leu	Gly	Pro	Val	Ser	
				275					280					285			
	cgc	tcc	tac	ggg	ggc	cag	tac	aga	tgc	tac	ggt	gca	tac	aac	ctc	tcc	1092
	Arg	Ser	Tyr	Gly	Gly	Gln	Tyr	Arg	Cys	Tyr	Gly	Ala	Tyr	Asn	Leu	Ser	
50			290					295					300				
	tcc	gag	tgg	tcg	gcc	ccc	agc	gac	ccc	ctg	gac	atc	ctg	atc	aca	gga	1140
	Ser	Glu	Trp	Ser	Ala	Pro	Ser	Asp	Pro	Leu	Asp	Ile	Leu	Ile	Thr	Gly	

55

EP 1 199 362 A2

	305		310		315	
5	cag atc cat ggc aca ccc ttc atc tca gtg cag cca ggc ccc aca gtg Gln Ile His Gly Thr Pro Phe Ile Ser Val Gln Pro Gly Pro Thr Val 320 325 330 335	1188				
	gcc tca gga gag aac gtg acc ctg ctg tgt cag tca tgg cgg cag ttc Ala Ser Gly Glu Asn Val Thr Leu Leu Cys Gln Ser Trp Arg Gln Phe 340 345 350	1236				
10	cac act ttc ctt ctg acc aag gcg gga gca gct gat gcc cca ctc cgt His Thr Phe Leu Leu Thr Lys Ala Gly Ala Ala Asp Ala Pro Leu Arg 355 360 365	1284				
15	cta aga tca ata cac gaa tat cct aag tac cag gct gaa ttc ccc atg Leu Arg Ser Ile His Glu Tyr Pro Lys Tyr Gln Ala Glu Phe Pro Met 370 375 380	1332				
	agt cct gtg acc tca gcc cac gcg ggg acc tac agg tgc tac ggc tca Ser Pro Val Thr Ser Ala His Ala Gly Thr Tyr Arg Cys Tyr Gly Ser 385 390 395	1380				
20	ctc aac tcc gac ccc tac ctg ctg tct cac ccc agt gag ccc ctg gag Leu Asn Ser Asp Pro Tyr Leu Leu Ser His Pro Ser Glu Pro Leu Glu 400 405 410 415	1428				
25	ctc gtg gtc tca gga ccc tcc atg ggt tcc agc ccc cca ccc acc ggt Leu Val Val Ser Gly Pro Ser Met Gly Ser Ser Pro Pro Pro Thr Gly 420 425 430	1476				
	ccc atc tcc aca cct gca ggc cct gag gac cag ccc ctc acc ccc act Pro Ile Ser Thr Pro Ala Gly Pro Glu Asp Gln Pro Leu Thr Pro Thr 435 440 445	1524				
30	ggg tcg gat ccc caa agt ggt ctg gga agg cac ctg ggg gtt gtg atc Gly Ser Asp Pro Gln Ser Gly Leu Gly Arg His Leu Gly Val Val Ile 450 455 460	1572				
35	ggc atc ttg gtg gcc gtc gtc cta ctg ctc ctc ctc ctc ctc ctc Gly Ile Leu Val Ala Val Val Leu Leu Leu Leu Leu Leu Leu Leu 465 470 475	1620				
	ttc ctc atc ctc cga cat cga cgt cag ggc aaa cac tgg aca tcg acc Phe Leu Ile Leu Arg His Arg Arg Gln Gly Lys His Trp Thr Ser Thr 480 485 490 495	1668				
40	cag aga aag gct gat ttc caa cat cct gca ggg gct gtg ggg cca gag Gln Arg Lys Ala Asp Phe Gln His Pro Ala Gly Ala Val Gly Pro Glu 500 505 510	1716				
45	ccc aca gac aga ggc ctg cag tgg agg tcc agc cca gct gcc gac gcc Pro Thr Asp Arg Gly Leu Gln Trp Arg Ser Ser Pro Ala Ala Asp Ala 515 520 525	1764				
	cag gaa gaa aac ctc tat gct gcc gtg aag gac aca cag cct gaa gat Gln Glu Glu Asn Leu Tyr Ala Ala Val Lys Asp Thr Gln Pro Glu Asp 530 535 540	1812				
50	ggg gtg gag atg gac act cgg gct gct gca tct gaa gcc ccc cag gat Gly Val Glu Met Asp Thr Arg Ala Ala Ala Ser Glu Ala Pro Gln Asp 545 550 555	1860				
55						

**EP 1 199 362 A2**

5

EP 1 199 362 A2

Ser Arg Ala Ile Phe Ser Val Gly Pro Val Ser Pro Asn Arg Arg Trp  
 180 185 190  
 5 Ser His Arg Cys Tyr Gly Tyr Asp Leu Asn Ser Pro Tyr Val Trp Ser  
 195 200 205  
 Ser Pro Ser Asp Leu Leu Glu Leu Leu Val Pro Gly Val Ser Lys Lys  
 210 215 220  
 10 Pro Ser Leu Ser Val Gln Pro Gly Pro Val Val Ala Pro Gly Glu Ser  
 225 230 235 240  
 Leu Thr Leu Gln Cys Val Ser Asp Val Gly Tyr Asp Arg Phe Val Leu  
 245 250 255  
 15 Tyr Lys Glu Gly Glu Arg Asp Leu Arg Gln Leu Pro Gly Arg Gln Pro  
 260 265 270  
 Gln Ala Gly Leu Ser Gln Ala Asn Phe Thr Leu Gly Pro Val Ser Arg  
 275 280 285  
 20 Ser Tyr Gly Gly Gln Tyr Arg Cys Tyr Gly Ala Tyr Asn Leu Ser Ser  
 290 295 300  
 Glu Trp Ser Ala Pro Ser Asp Pro Leu Asp Ile Leu Ile Thr Gly Gln  
 305 310 315 320  
 25 Ile His Gly Thr Pro Phe Ile Ser Val Gln Pro Gly Pro Thr Val Ala  
 325 330 335  
 Ser Gly Glu Asn Val Thr Leu Leu Cys Gln Ser Trp Arg Gln Phe His  
 340 345 350  
 30 Thr Phe Leu Leu Thr Lys Ala Gly Ala Ala Asp Ala Pro Leu Arg Leu  
 355 360 365  
 Arg Ser Ile His Glu Tyr Pro Lys Tyr Gln Ala Glu Phe Pro Met Ser  
 370 375 380  
 35 Pro Val Thr Ser Ala His Ala Gly Thr Tyr Arg Cys Tyr Gly Ser Leu  
 385 390 395 400  
 Asn Ser Asp Pro Tyr Leu Leu Ser His Pro Ser Glu Pro Leu Glu Leu  
 405 410 415  
 40 Val Val Ser Gly Pro Ser Met Gly Ser Ser Pro Pro Pro Thr Gly Pro  
 420 425 430  
 Ile Ser Thr Pro Ala Gly Pro Glu Asp Gln Pro Leu Thr Pro Thr Gly  
 435 440 445  
 45 Ser Asp Pro Gln Ser Gly Leu Gly Arg His Leu Gly Val Val Ile Gly  
 450 455 460  
 Ile Leu Val Ala Val Val Leu Leu Leu Leu Leu Leu Leu Leu Phe  
 465 470 475 480  
 50 Leu Ile Leu Arg His Arg Arg Gln Gly Lys His Trp Thr Ser Thr Gln  
 485 490 495

55

EP 1 199 362 A2

	Arg Lys Ala Asp Phe Gln His Pro Ala Gly Ala Val Gly Pro Glu Pro	
	500 505 510	
5	Thr Asp Arg Gly Leu Gln Trp Arg Ser Ser Pro Ala Ala Asp Ala Gln	
	515 520 525	
	Glu Glu Asn Leu Tyr Ala Ala Val Lys Asp Thr Gln Pro Glu Asp Gly	
	530 535 540	
10	Val Glu Met Asp Thr Arg Ala Ala Ala Ser Glu Ala Pro Gln Asp Val	
	545 550 555 560	
	Thr Tyr Ala Gln Leu His Ser Leu Thr Leu Arg Arg Lys Ala Thr Glu	
	565 570 575	
15	Pro Pro Pro Ser Gln Glu Arg Glu Pro Pro Ala Glu Pro Ser Ile Tyr	
	580 585 590	
	Ala Thr Leu Ala Ile His	
	595	
20	<210> 11	
	<211> 2446	
	<212> DNA	
	<213> human	
	<220>	
25	<221> CDS	
	<222> (171)..(1040)	
	<400> 11	
30	cgcagctcaa cctgagctac acagccagat gcgagatgct tctctgctga tctgagtctg	60
	cctgcagcat ggaccttggt cttccctgaa gcatctccag ggctggaggg acgactgcca	120
	tgcacctagg gcttatccat cgcagagca gggcagtggg aggagacgct atg acc	176
	Met Thr	
	1	
35	ccc atc ctc acg gtc ctg atc tgt ctc ggg ctg agt ctg ggc ccc cgg	224
	Pro Ile Leu Thr Val Leu Ile Cys Leu Gly Leu Ser Leu Gly Pro Arg	
	5 10 15	
40	acc cac gtg cag gca ggg acc ctc ccc aag ccc aca ctc tgg gct gag	272
	Thr His Val Gln Ala Gly Thr Leu Pro Lys Pro Thr Leu Trp Ala Glu	
	20 25 30	
	cca ggc tct gtg atc acc cag ggg agt ccc gtg acc ctc tgg tgt cag	320
	Pro Gly Ser Val Ile Thr Gln Gly Ser Pro Val Thr Leu Trp Cys Gln	
45	35 40 45 50	
	ggg atc ctg gag acc cag gag tac cgt ctg tat aga gaa aag aaa aca	368
	Gly Ile Leu Glu Thr Gln Glu Tyr Arg Leu Tyr Arg Glu Lys Lys Thr	
	55 60 65	
50	gca ccc tgg att aca cgg atc cca cag gag att gtg aag aag ggc cag	416
	Ala Pro Trp Ile Thr Arg Ile Pro Gln Glu Ile Val Lys Lys Gly Gln	
	70 75 80	
	ttc ccc atc ccg tcc atc acc tgg gaa cac acc ggg cgg tat cgc tgt	464
55		

EP 1 199 362 A2

	Phe	Pro	Ile	Pro	Ser	Ile	Thr	Trp	Glu	His	Thr	Gly	Arg	Tyr	Arg	Cys	
			85					90					95				
5	ttc	tac	ggg	agc	cac	act	gca	ggc	tgg	tca	gag	ccc	agt	gac	ccc	ctg	512
	Phe	Tyr	Gly	Ser	His	Thr	Ala	Gly	Trp	Ser	Glu	Pro	Ser	Asp	Pro	Leu	
	100						105					110					
	gag	ctg	gtg	gtg	aca	gga	gcc	tac	atc	aaa	ccc	acc	ctc	tcg	gct	cta	560
10	Glu	Leu	Val	Val	Thr	Gly	Ala	Tyr	Ile	Lys	Pro	Thr	Leu	Ser	Ala	Leu	
	115					120					125					130	
	ccc	agc	cct	gtg	gtg	acc	tca	gga	ggg	aac	gtg	acc	ctc	cat	tgt	gtc	608
	Pro	Ser	Pro	Val	Val	Thr	Ser	Gly	Gly	Asn	Val	Thr	Leu	His	Cys	Val	
					135					140					145		
15	tca	cag	gtg	gca	ttt	ggc	agc	ttc	att	ctg	tgt	aag	gaa	gga	gaa	gat	656
	Ser	Gln	Val	Ala	Phe	Gly	Ser	Phe	Ile	Leu	Cys	Lys	Glu	Gly	Glu	Asp	
				150				155						160			
	gaa	cac	cca	caa	tgc	ctg	aac	tca	cag	ccc	cgt	acc	cat	ggg	tgg	tcc	704
20	Glu	His	Pro	Gln	Cys	Leu	Asn	Ser	Gln	Pro	Arg	Thr	His	Gly	Trp	Ser	
			165					170					175				
	cgg	gcc	atc	ttc	tct	gtg	ggc	ccc	gtg	agc	ccg	agt	cgc	agg	tgg	tcg	752
	Arg	Ala	Ile	Phe	Ser	Val	Gly	Pro	Val	Ser	Pro	Ser	Arg	Arg	Trp	Ser	
	180						185					190					
25	tac	agg	tgc	tat	gct	tat	gac	tcg	aac	tct	ccc	cat	gtg	tgg	tct	cta	800
	Tyr	Arg	Cys	Tyr	Ala	Tyr	Asp	Ser	Asn	Ser	Pro	His	Val	Trp	Ser	Leu	
	195					200					205					210	
	ccc	agt	gat	ctc	ctg	gag	ctc	ctg	gtc	cca	gga	gca	gct	gag	acc	ctc	848
30	Pro	Ser	Asp	Leu	Leu	Glu	Leu	Val	Pro	Gly	Ala	Ala	Glu	Thr	Leu		
				215					220					225			
	agc	cca	cca	caa	aac	aag	tcc	gat	tcc	aag	gct	gga	gca	gct	aac	acc	896
	Ser	Pro	Pro	Gln	Asn	Lys	Ser	Asp	Ser	Lys	Ala	Gly	Ala	Ala	Asn	Thr	
				230					235					240			
35	ctc	agc	cca	tca	caa	aac	aag	act	gcc	tca	cac	ccc	cag	gat	tac	aca	944
	Leu	Ser	Pro	Ser	Gln	Asn	Lys	Thr	Ala	Ser	His	Pro	Gln	Asp	Tyr	Thr	
			245				250					255					
	gtg	gag	aat	ctc	atc	cgc	atg	ggc	ata	gct	ggc	ttg	gtc	ctg	gtg	gtc	992
40	Val	Glu	Asn	Leu	Ile	Arg	Met	Gly	Ile	Ala	Gly	Leu	Val	Leu	Val	Val	
	260						265				270						
	ctc	ggg	att	ctg	cta	ttt	gag	gct	cag	cac	agc	cag	aga	agc	ctc	tga	1040
	Leu	Gly	Ile	Leu	Leu	Phe	Glu	Ala	Gln	His	Ser	Gln	Arg	Ser	Leu		
	275					280					285						
45	gatgcagccg	ggaggtgaac	agcagagaga	agaatgtacc	cttcagagtg	gtggagcctt											1100
	gggaacagat	ctgatgatgc	caggagggttc	cgaggagacaa	tttagggctg	atgctatctg											1160
	gactgtctgc	caatcatttt	tagagggagg	aatcagtgtt	ggattgcaga	gacattttct											1220
50	ggagtgatcc	atgaaggacc	attaacatgt	gatacctttc	ctctctatta	atgttgactt											1280
	cccttggttg	gacccctctt	tttccccacc	cccagacaga	catgaggcta	catccacat											1340
55																	

EP 1 199 362 A2

5 ggcagcgttg ggtccacacc tctgcacatc tgtgtgctct ggtccatggt gtgtaacaca 1400  
 gtcttcttta ttactcattg ccatactccc tgggtgtgctt tactgagcct ccatctcttc 1460  
 aattcagagt tccaaacgtg cttcagtaac taaatcaatg ggagagtatc ggatttcaac 1520  
 caggaaaaga taaatccacc ctgatgccct gacaccctct ctgaacccta cgagcccttc 1580  
 10 cctccttctc acatgctacc tgtgcagctt ctccttagat cattgtgtaa ccatcactgc 1640  
 catcctgttc cacacatggt catcaccta caccattca gcagccactc cccattccct 1700  
 cttccctcca gcacctgcta accacaaatg tgctttctgt ctctacggat ttgcctattc 1760  
 15 tgtctgaaaa catttcaatc tcctttgacc tgtgagctcc tcacttcgag acttctgccc 1820  
 tttccaggca gaaccaaagt acaccacgtc aaaagcaatg ataggcattt gcagtgtggt 1880  
 ggtgatccac gaaaggaaaa tcacggaagc aggatagaaa tccagctgca gacaagacct 1940  
 20 caggctgatg aatcttgaca agcagttgag ctgttttttt ctactcacct aggacagtca 2000  
 ggcagaagta tgcaaaatga ctggggctga ttcttttctg aattgtcgca aacagcaaga 2060  
 ggacttgagt cctagcatta aagagttcaa catgtctagg tccaagacca ctgttgtgtt 2120  
 25 tgaaggatgt aaaaccctgc tgcataggat ggaatatattg gagggaggat cctgaaaaac 2180  
 atgagggatc aaatagtect caactttcta ggacaaaggg agcagctatt tgccatctac 2240  
 cctccagaat aaagaaatct tatcattcac catctaccct ctagaataaa gaaatcttat 2300  
 30 cattcgccat ctaccctgta gaataaagaa atcttatcat tcaccgtcta cctcttagag 2360  
 taaacaaatc ttatcattca ccatctaccc tctagaataa agaaatctta tcattcgcca 2420  
 tctaccctct agaataaaga aatctt 2446

35 <210> 12

<211> 289

<212> PRT

<213> human

40 <400> 12

Met Thr Pro Ile Leu Thr Val Leu Ile Cys Leu Gly Leu Ser Leu Gly  
           1                  5                  10                  15

Pro Arg Thr His Val Gln Ala Gly Thr Leu Pro Lys Pro Thr Leu Trp  
                   20                  25                  30

Ala Glu Pro Gly Ser Val Ile Thr Gln Gly Ser Pro Val Thr Leu Trp  
           35                  40                  45

Cys Gln Gly Ile Leu Glu Thr Gln Glu Tyr Arg Leu Tyr Arg Glu Lys  
           50                  55                  60

Lys Thr Ala Pro Trp Ile Thr Arg Ile Pro Gln Glu Ile Val Lys Lys  
           65                  70                  75                  80

55

EP 1 199 362 A2

	Gly	Gln	Phe	Pro	Ile	Pro	Ser	Ile	Thr	Trp	Glu	His	Thr	Gly	Arg	Tyr	
					85					90					95		
5	Arg	Cys	Phe	Tyr	Gly	Ser	His	Thr	Ala	Gly	Trp	Ser	Glu	Pro	Ser	Asp	
				100					105					110			
	Pro	Leu	Glu	Leu	Val	Val	Thr	Gly	Ala	Tyr	Ile	Lys	Pro	Thr	Leu	Ser	
			115					120					125				
10	Ala	Leu	Pro	Ser	Pro	Val	Val	Thr	Ser	Gly	Gly	Asn	Val	Thr	Leu	His	
		130						135				140					
	Cys	Val	Ser	Gln	Val	Ala	Phe	Gly	Ser	Phe	Ile	Leu	Cys	Lys	Glu	Gly	
	145					150					155				160		
15	Glu	Asp	Glu	His	Pro	Gln	Cys	Leu	Asn	Ser	Gln	Pro	Arg	Thr	His	Gly	
					165					170					175		
	Trp	Ser	Arg	Ala	Ile	Phe	Ser	Val	Gly	Pro	Val	Ser	Pro	Ser	Arg	Arg	
				180					185					190			
20	Trp	Ser	Tyr	Arg	Cys	Tyr	Ala	Tyr	Asp	Ser	Asn	Ser	Pro	His	Val	Trp	
			195					200					205				
	Ser	Leu	Pro	Ser	Asp	Leu	Leu	Glu	Leu	Val	Pro	Gly	Ala	Ala	Glu		
		210					215				220						
25	Thr	Leu	Ser	Pro	Pro	Gln	Asn	Lys	Ser	Asp	Ser	Lys	Ala	Gly	Ala	Ala	
	225					230				235					240		
	Asn	Thr	Leu	Ser	Pro	Ser	Gln	Asn	Lys	Thr	Ala	Ser	His	Pro	Gln	Asp	
					245					250				255			
30	Tyr	Thr	Val	Glu	Asn	Leu	Ile	Arg	Met	Gly	Ile	Ala	Gly	Leu	Val	Leu	
				260					265					270			
	Val	Val	Leu	Gly	Ile	Leu	Leu	Phe	Glu	Ala	Gln	His	Ser	Gln	Arg	Ser	
			275					280					285				
35	Leu																
	<210>	13															
	<211>	1910															
40	<212>	DNA															
	<213>	human															
	<220>																
	<221>	CDS															
	<222>	(183) .. (1652)															
45	<400>	13															
	ctcactgcca	cacgcagctc	aacctgagct	acacagccag	atgcgagatg	cttctctgct										60	
	gatctgagtc	tgccctgcagc	atggaccttg	gtcttccctg	aagcatctcc	agggctggag										120	
50	ggacgactgc	catgcaccga	gggctcatcc	atccgcagag	cagggcagtg	ggaggagacg										180	
	ct atg acc ccc atc gtc aca gtc ctg atc tgt ctc agg ctg agt ctg															227	
	Met Thr Pro Ile Val Thr Val Leu Ile Cys Leu Arg Leu Ser Leu																
	1				5					10					15		
55																	



EP 1 199 362 A2

5	ggc ccc cgg acc cac gtg cag gca ggg acc ctc ccc aag ccc aca ctc Gly Pro Arg Thr His Val Gln Ala Gly Thr Leu Pro Lys Pro Thr Leu 20 25 30	275
	tgg gct gag cca ggc tct gtg atc acc cag ggg agt ccc gtg acc ctc Trp Ala Glu Pro Gly Ser Val Ile Thr Gln Gly Ser Pro Val Thr Leu 35 40 45	323
10	tgg tgt cag ggg atc ctg gag acc cag gag tac cgt ctg tat aga gaa Trp Cys Gln Gly Ile Leu Glu Thr Gln Glu Tyr Arg Leu Tyr Arg Glu 50 55 60	371
15	aag aaa aca gca ccc tgg att aca cgg atc cca cag gag att gtg aag Lys Lys Thr Ala Pro Trp Ile Thr Arg Ile Pro Gln Glu Ile Val Lys 65 70 75	419
	aag ggc cag ttc ccc atc cca tcc atc acc tgg gaa cac aca ggg cgg Lys Gly Gln Phe Pro Ile Pro Ser Ile Thr Trp Glu His Thr Gly Arg 80 85 90 95	467
20	tat cgc tgt ttc tac ggt agc cac act gca ggc tgg tca gag ccc agt Tyr Arg Cys Phe Tyr Gly Ser His Thr Ala Gly Trp Ser Glu Pro Ser 100 105 110	515
25	gac ccc ctg gag ctg gtg gtg aca gga gcc tac atc aaa ccc acc ctc Asp Pro Leu Glu Leu Val Val Thr Gly Ala Tyr Ile Lys Pro Thr Leu 115 120 125	563
	tca gct cta ccc agc cct gtg gtg acc tca gga ggg aac gtg acc ctc Ser Ala Leu Pro Ser Pro Val Val Thr Ser Gly Gly Asn Val Thr Leu 130 135 140	611
30	cat tgt gtc tca cag gtg gca ttt ggc agc ttc att ctg tgt aag gaa His Cys Val Ser Gln Val Ala Phe Gly Ser Phe Ile Leu Cys Lys Glu 145 150 155	659
35	gga gaa gat gaa cac cca caa tgc ctg aac tca cag ccc cgt acc cat Gly Glu Asp Glu His Pro Gln Cys Leu Asn Ser Gln Pro Arg Thr His 160 165 170 175	707
	ggg tgg tcc cgg gcc atc ttc tct gtg ggc ccc gtg agc ccg agt cgc Gly Trp Ser Arg Ala Ile Phe Ser Val Gly Pro Val Ser Pro Ser Arg 180 185 190	755
40	agg tgg tcg tac agg tgc tat gct tat gac tcg aac tct ccc cat gtg Arg Trp Ser Tyr Arg Cys Tyr Ala Tyr Asp Ser Asn Ser Pro His Val 195 200 205	803
45	tgg tct cta ccc agt gat ctc ctg gag ctc ctg gtc cta ggt gtt tct Trp Ser Leu Pro Ser Asp Leu Leu Glu Leu Leu Val Leu Gly Val Ser 210 215 220	851
	aag aag cca tca ctc tca gtg cag cca ggt cct ata gtg gcc cct ggg Lys Lys Pro Ser Leu Ser Val Gln Pro Gly Pro Ile Val Ala Pro Gly 225 230 235	899
50	gag agc ctg acc ctc cag tgt gtt tct gat gtc agc tac gac aga ttt Glu Ser Leu Thr Leu Gln Cys Val Ser Asp Val Ser Tyr Asp Arg Phe 240 245 250 255	947

55

EP 1 199 362 A2

	gtt ctg tat aag gag gga gaa cgt gac ttc ctc cag ctc cct ggc cca	995
	Val Leu Tyr Lys Glu Gly Glu Arg Asp Phe Leu Gln Leu Pro Gly Pro	
	260 265 270	
5	cag ccc cag gct ggg ctc tcc cag gcc aac ttc acc ctg ggc cct gtg	1043
	Gln Pro Gln Ala Gly Leu Ser Gln Ala Asn Phe Thr Leu Gly Pro Val	
	275 280 285	
10	agc cgc tcc tac ggg ggc cag tac aga tgc tcc ggt gca tac aac ctc	1091
	Ser Arg Ser Tyr Gly Gly Gln Tyr Arg Cys Ser Gly Ala Tyr Asn Leu	
	290 295 300	
	tcc tcc gag tgg tgg gcc ccc agc gac ccc ctg gac atc ctg atc gca	1139
	Ser Ser Glu Trp Ser Ala Pro Ser Asp Pro Leu Asp Ile Leu Ile Ala	
	305 310 315	
15	gga cag ttc cgt ggc aga ccc ttc atc tgg gtg cat ccg ggc ccc acg	1187
	Gly Gln Phe Arg Gly Arg Pro Phe Ile Ser Val His Pro Gly Pro Thr	
	320 325 330 335	
20	gtg gcc tca gga gag aac gtg acc ctg ctg tgt cag tca tgg ggg ccg	1235
	Val Ala Ser Gly Glu Asn Val Thr Leu Leu Cys Gln Ser Trp Gly Pro	
	340 345 350	
	ttc cac act ttc ctt ctg acc aag gcg gga gca gct gat gcc ccc ctc	1283
	Phe His Thr Phe Leu Leu Thr Lys Ala Gly Ala Ala Asp Ala Pro Leu	
	355 360 365	
25	cgt ctc aga tca ata cac gaa tat cct aag tac cag gct gaa ttc cct	1331
	Arg Leu Arg Ser Ile His Glu Tyr Pro Lys Tyr Gln Ala Glu Phe Pro	
	370 375 380	
30	atg agt cct gtg acc tca gcc cac tgg ggg acc tac agg tgc tac ggc	1379
	Met Ser Pro Val Thr Ser Ala His Ser Gly Thr Tyr Arg Cys Tyr Gly	
	385 390 395	
	tca ctc agc tcc aac ccc tac ctg ctg tct cac ccc agt gac tcc ctg	1427
	Ser Leu Ser Ser Asn Pro Tyr Leu Leu Ser His Pro Ser Asp Ser Leu	
	400 405 410 415	
35	gag ctc atg gtc tca gga gca gct gag acc ctc agc cca cca caa aac	1475
	Glu Leu Met Val Ser Gly Ala Ala Glu Thr Leu Ser Pro Pro Gln Asn	
	420 425 430	
40	aag tcc gat tcc aag gct gga gca gct aac acc ctc agc cca tca caa	1523
	Lys Ser Asp Ser Lys Ala Gly Ala Ala Asn Thr Leu Ser Pro Ser Gln	
	435 440 445	
	aac aag act gcc tca cac ccc cag gat tac aca gtg gag aat ctc atc	1571
	Asn Lys Thr Ala Ser His Pro Gln Asp Tyr Thr Val Glu Asn Leu Ile	
	450 455 460	
45	cgc atg ggc ata gct ggc ttg gtc ctg gtg gtc ctc ggg att ctg cta	1619
	Arg Met Gly Ile Ala Gly Leu Val Leu Val Val Leu Gly Ile Leu Leu	
	465 470 475	
50	ttt gag gct cag cac agc cag aga agc ctc tga gatgcagccg ggaggtgaac	1672
	Phe Glu Ala Gln His Ser Gln Arg Ser Leu	
	480 485	
	agcagagaga agaattgtacc cttcagagtg gtggagcctt gggaacagat ctgatgatgc	1732
55		

EP 1 199 362 A2

caggaggttc cgggagacaa tttagggctg atgttatctg gactgtctgc caatcatttt 1792

5 tagagggagg aatcagtgtt ggattgcaga gacattttct ggagtgatcc atgaaggacc 1852

attaacatgt gatacctttc ctctctatta atgttgactt cccttggttg gatcctct 1910

<210> 14  
 <211> 489  
 <212> PRT  
 <213> human

<400> 14

15 Met Thr Pro Ile Val Thr Val Leu Ile Cys Leu Arg Leu Ser Leu Gly  
 1 5 10 15

Pro Arg Thr His Val Gln Ala Gly Thr Leu Pro Lys Pro Thr Leu Trp  
 20 25 30

Ala Glu Pro Gly Ser Val Ile Thr Gln Gly Ser Pro Val Thr Leu Trp  
 20 35 40 45

Cys Gln Gly Ile Leu Glu Thr Gln Glu Tyr Arg Leu Tyr Arg Glu Lys  
 50 55 60

Lys Thr Ala Pro Trp Ile Thr Arg Ile Pro Gln Glu Ile Val Lys Lys  
 25 65 70 75 80

Gly Gln Phe Pro Ile Pro Ser Ile Thr Trp Glu His Thr Gly Arg Tyr  
 85 90 95

Arg Cys Phe Tyr Gly Ser His Thr Ala Gly Trp Ser Glu Pro Ser Asp  
 30 100 105 110

Pro Leu Glu Leu Val Val Thr Gly Ala Tyr Ile Lys Pro Thr Leu Ser  
 115 120 125

Ala Leu Pro Ser Pro Val Val Thr Ser Gly Gly Asn Val Thr Leu His  
 35 130 135 140

Cys Val Ser Gln Val Ala Phe Gly Ser Phe Ile Leu Cys Lys Glu Gly  
 145 150 155 160

Glu Asp Glu His Pro Gln Cys Leu Asn Ser Gln Pro Arg Thr His Gly  
 40 165 170 175

Trp Ser Arg Ala Ile Phe Ser Val Gly Pro Val Ser Pro Ser Arg Arg  
 180 185 190

Trp Ser Tyr Arg Cys Tyr Ala Tyr Asp Ser Asn Ser Pro His Val Trp  
 45 195 200 205

Ser Leu Pro Ser Asp Leu Leu Glu Leu Val Leu Gly Val Ser Lys  
 210 215 220

Lys Pro Ser Leu Ser Val Gln Pro Gly Pro Ile Val Ala Pro Gly Glu  
 50 225 230 235 240

Ser Leu Thr Leu Gln Cys Val Ser Asp Val Ser Tyr Asp Arg Phe Val  
 245 250 255

55

EP 1 199 362 A2

Leu Tyr Lys Glu Gly Glu Arg Asp Phe Leu Gln Leu Pro Gly Pro Gln  
 260 265 270  
 5 Pro Gln Ala Gly Leu Ser Gln Ala Asn Phe Thr Leu Gly Pro Val Ser  
 275 280 285  
 Arg Ser Tyr Gly Gly Gln Tyr Arg Cys Ser Gly Ala Tyr Asn Leu Ser  
 290 295 300  
 10 Ser Glu Trp Ser Ala Pro Ser Asp Pro Leu Asp Ile Leu Ile Ala Gly  
 305 310 315 320  
 Gln Phe Arg Gly Arg Pro Phe Ile Ser Val His Pro Gly Pro Thr Val  
 325 330 335  
 15 Ala Ser Gly Glu Asn Val Thr Leu Leu Cys Gln Ser Trp Gly Pro Phe  
 340 345 350  
 His Thr Phe Leu Leu Thr Lys Ala Gly Ala Ala Asp Ala Pro Leu Arg  
 355 360 365  
 20 Leu Arg Ser Ile His Glu Tyr Pro Lys Tyr Gln Ala Glu Phe Pro Met  
 370 375 380  
 Ser Pro Val Thr Ser Ala His Ser Gly Thr Tyr Arg Cys Tyr Gly Ser  
 385 390 395 400  
 25 Leu Ser Ser Asn Pro Tyr Leu Leu Ser His Pro Ser Asp Ser Leu Glu  
 405 410 415  
 Leu Met Val Ser Gly Ala Ala Glu Thr Leu Ser Pro Pro Gln Asn Lys  
 420 425 430  
 30 Ser Asp Ser Lys Ala Gly Ala Ala Asn Thr Leu Ser Pro Ser Gln Asn  
 435 440 445  
 Lys Thr Ala Ser His Pro Gln Asp Tyr Thr Val Glu Asn Leu Ile Arg  
 450 455 460  
 35 Met Gly Ile Ala Gly Leu Val Leu Val Val Leu Gly Ile Leu Leu Phe  
 465 470 475 480  
 Glu Ala Gln His Ser Gln Arg Ser Leu  
 485

<210> 15  
 <211> 1725  
 <212> DNA  
 <213> human

<220>  
 <221> CDS  
 <222> (40)..(1491)

<400> 15  
 ctcacccatc cgcagagcag ggcagtgagg ggagacgcc atg acc ccc atc ctc  
 Met Thr Pro Ile Leu  
 1 5

54

EP 1 199 362 A2

	acg gtc ctg atc tgt ctc ggg ctg agt ctg ggc ccc agg acc cac gtg	102
	Thr Val Leu Ile Cys Leu Gly Leu Ser Leu Gly Pro Arg Thr His Val	
	10 15 20	
5	cag gca ggg cac ctc ccc aag ccc acc ctc tgg gct gag cca ggc tct	150
	Gln Ala Gly His Leu Pro Lys Pro Thr Leu Trp Ala Glu Pro Gly Ser	
	25 30 35	
10	gtg atc atc cag gga agt cct gtg acc ctc agg tgt cag ggg agc ctt	198
	Val Ile Ile Gln Gly Ser Pro Val Thr Leu Arg Cys Gln Gly Ser Leu	
	40 45 50	
15	cag gct gag gag tac cat cta tat agg gaa aac aaa tca gca tcc tgg	246
	Gln Ala Glu Glu Tyr His Leu Tyr Arg Glu Asn Lys Ser Ala Ser Trp	
	55 60 65	
20	gtt aga cgg ata caa gag cct ggg aag aat ggc cag ttc ccc atc cca	294
	Val Arg Arg Ile Gln Glu Pro Gly Lys Asn Gly Gln Phe Pro Ile Pro	
	70 75 80 85	
25	tcc atc acc tgg gaa cac gca ggg cgg tat cac tgt cag tac tac agc	342
	Ser Ile Thr Trp Glu His Ala Gly Arg Tyr His Cys Gln Tyr Tyr Ser	
	90 95 100	
30	cac aat cac tca tca gag tac agt gac ccc ctg gag ctg gtg gtg aca	390
	His Asn His Ser Ser Glu Tyr Ser Asp Pro Leu Glu Leu Val Val Thr	
	105 110 115	
35	gga gcc tac agc aaa ccc acc ctc tca gct ctg ccc agc cct gtg gtg	438
	Gly Ala Tyr Ser Lys Pro Thr Leu Ser Ala Leu Pro Ser Pro Val Val	
	120 125 130	
40	acc tta gga ggg aac gtg acc ctc cag tgt gtc tca cag gtg gca ttt	486
	Thr Leu Gly Gly Asn Val Thr Leu Gln Cys Val Ser Gln Val Ala Phe	
	135 140 145	
45	gac ggc ttc att ctg tgt aag gaa gga gaa gat gaa cac cca caa cgc	534
	Asp Gly Phe Ile Leu Cys Lys Glu Gly Glu Asp Glu His Pro Gln Arg	
	150 155 160 165	
50	ctg aac tcc cat tcc cat gcc cgt ggg tgg tcc tgg gcc atc ttc tcc	582
	Leu Asn Ser His Ser His Ala Arg Gly Trp Ser Trp Ala Ile Phe Ser	
	170 175 180	
55	gtg ggc ccc gtg agc ccg agt cgc agg tgg tcg tac agg tgc tat gct	630
	Val Gly Pro Val Ser Pro Ser Arg Arg Trp Ser Tyr Arg Cys Tyr Ala	
	185 190 195	
60	tat gac tcg aac tct ccc tat gtg tgg tct cta ccc agt gat ctc ctg	678
	Tyr Asp Ser Asn Ser Pro Tyr Val Trp Ser Leu Pro Ser Asp Leu Leu	
	200 205 210	
65	gag ctc ctg gtc cca ggt gtt tct aag aag cca tca ctc tca gtg cag	726
	Glu Leu Leu Val Pro Gly Val Ser Lys Lys Pro Ser Leu Ser Val Gln	
	215 220 225	
70	cca ggt cct atg gtg gcc ccc ggg gag agc ctg acc ctc cag tgt gtc	774
	Pro Gly Pro Met Val Ala Pro Gly Glu Ser Leu Thr Leu Gln Cys Val	
	230 235 240 245	
75	tct gat gtc ggc tac gac aga ttt gtt ctg tat aag gag gga gaa cgt	822

EP 1 199 362 A2

	Ser Asp Val Gly Tyr Asp Arg Phe Val Leu Tyr Lys Glu Gly Glu Arg	
	250 255 260	
5	gac ttc ctc cag cgc cct ggt tgg cag ccc cag gct ggg ctc tcc cag Asp Phe Leu Gln Arg Pro Gly Trp Gln Pro Gln Ala Gly Leu Ser Gln	870
	265 270 275	
10	gcc aac ttc acc ctg ggc cct gtg agc ccc tcc cac ggg ggc cag tac Ala Asn Phe Thr Leu Gly Pro Val Ser Pro Ser His Gly Gly Gln Tyr	918
	280 285 290	
15	aga tgc tac agt gca cac aac ctc tcc tcc gag tgg tcg gcc ccc agt Arg Cys Tyr Ser Ala His Asn Leu Ser Ser Glu Trp Ser Ala Pro Ser	966
	295 300 305	
20	gac ccc ctg gac atc ctg atc aca gga cag ttc tat gac aga ccc tct Asp Pro Leu Asp Ile Leu Ile Thr Gly Gln Phe Tyr Asp Arg Pro Ser	1014
	310 315 320 325	
25	ctc tcg gtg cag ccg gtc ccc aca gta gcc cca gga aag aac gtg acc Leu Ser Val Gln Pro Val Pro Thr Val Ala Pro Gly Lys Asn Val Thr	1062
	330 335 340	
30	ctg ctg tgt cag tca cgg ggg cag ttc cac act ttc ctt ctg acc aag Leu Leu Cys Gln Ser Arg Gly Gln Phe His Thr Phe Leu Leu Thr Lys	1110
	345 350 355	
35	gag ggg gca ggc cat ccc cca ctg cat ctg aga tca gag cac caa gct Glu Gly Ala Gly His Pro Pro Leu His Leu Arg Ser Glu His Gln Ala	1158
	360 365 370	
40	cag cag aac cag gct gaa ttc cgc atg ggt cct gtg acc tca gcc cac Gln Gln Asn Gln Ala Glu Phe Arg Met Gly Pro Val Thr Ser Ala His	1206
	375 380 385	
45	gtg ggg acc tac aga tgc tac agc tca ctc agc tcc aac ccc tac ctg Val Gly Thr Tyr Arg Cys Tyr Ser Ser Leu Ser Ser Asn Pro Tyr Leu	1254
	390 395 400 405	
50	ctg tct ctc ccc agt gac ccc ctg gag ctc gtg gtc tca gaa gca gct Leu Ser Leu Pro Ser Asp Pro Leu Glu Leu Val Val Ser Glu Ala Ala	1302
	410 415 420	
55	gag acc ctc agc cca tca caa aac aag aca gac tcc acg act aca tcc Glu Thr Leu Ser Pro Ser Gln Asn Lys Thr Asp Ser Thr Thr Thr Ser	1350
	425 430 435	
60	cta ggc caa cac ccc cag gat tac aca gtg gag aat ctc atc cgc atg Leu Gly Gln His Pro Gln Asp Tyr Thr Val Glu Asn Leu Ile Arg Met	1398
	440 445 450	
65	ggt gtg gct ggc ttg gtc ctg gtg gtc ctc ggg att ctg cta ttt gag Gly Val Ala Gly Leu Val Leu Val Val Leu Gly Ile Leu Leu Phe Glu	1446
	455 460 465	
70	gct cag cac agc cag aga agc cta caa gat gca gcc ggg agg tga Ala Gln His Ser Gln Arg Ser Leu Gln Asp Ala Ala Gly Arg	1491
	470 475 480	
75	acagcagaga ggacaatgca tccttcagcg tgggtggagcc tcagggacag atctgatgat	1551

EP 1 199 362 A2

cccaggaggc tctggaggac aatctaggac ctacattatc tggactgtat gctgggcatt 1611  
 tctagagaca gcaatcaata tttgagtgtgta aggaaactgt ctgggggtgat tcctagaaga 1671  
 5 tcattaaact gtggtacatt tttttgtcta aaaagcaggt cgtctcgttc caag 1725  
  
 <210> 16  
 <211> 483  
 <212> PRT  
 <213> human  
 10  
  
 <400> 16  
 Met Thr Pro Ile Leu Thr Val Leu Ile Cys Leu Gly Leu Ser Leu Gly  
 1 5 10 15  
 15 Pro Arg Thr His Val Gln Ala Gly His Leu Pro Lys Pro Thr Leu Trp  
 20 25 30  
 Ala Glu Pro Gly Ser Val Ile Ile Gln Gly Ser Pro Val Thr Leu Arg  
 35 40 45  
 20 Cys Gln Gly Ser Leu Gln Ala Glu Glu Tyr His Leu Tyr Arg Glu Asn  
 50 55 60  
 Lys Ser Ala Ser Trp Val Arg Arg Ile Gln Glu Pro Gly Lys Asn Gly  
 65 70 75 80  
 25 Gln Phe Pro Ile Pro Ser Ile Thr Trp Glu His Ala Gly Arg Tyr His  
 85 90 95  
 Cys Gln Tyr Tyr Ser His Asn His Ser Ser Glu Tyr Ser Asp Pro Leu  
 100 105 110  
 30 Glu Leu Val Val Thr Gly Ala Tyr Ser Lys Pro Thr Leu Ser Ala Leu  
 115 120 125  
 Pro Ser Pro Val Val Thr Leu Gly Gly Asn Val Thr Leu Gln Cys Val  
 130 135 140  
 35 Ser Gln Val Ala Phe Asp Gly Phe Ile Leu Cys Lys Glu Gly Glu Asp  
 145 150 155 160  
 Glu His Pro Gln Arg Leu Asn Ser His Ser His Ala Arg Gly Trp Ser  
 165 170 175  
 40 Trp Ala Ile Phe Ser Val Gly Pro Val Ser Pro Ser Arg Arg Trp Ser  
 180 185 190  
 Tyr Arg Cys Tyr Ala Tyr Asp Ser Asn Ser Pro Tyr Val Trp Ser Leu  
 195 200 205  
 45 Pro Ser Asp Leu Leu Glu Leu Leu Val Pro Gly Val Ser Lys Lys Pro  
 210 215 220  
 Ser Leu Ser Val Gln Pro Gly Pro Met Val Ala Pro Gly Glu Ser Leu  
 225 230 235 240  
 50 Thr Leu Gln Cys Val Ser Asp Val Gly Tyr Asp Arg Phe Val Leu Tyr  
 245 250 255  
 55

EP 1 199 362 A2

	Lys	Glu	Gly	Glu	Arg	Asp	Phe	Leu	Gln	Arg	Pro	Gly	Trp	Gln	Pro	Gln	
				260					265					270			
5	Ala	Gly	Leu	Ser	Gln	Ala	Asn	Phe	Thr	Leu	Gly	Pro	Val	Ser	Pro	Ser	
			275					280					285				
	His	Gly	Gly	Gln	Tyr	Arg	Cys	Tyr	Ser	Ala	His	Asn	Leu	Ser	Ser	Glu	
		290					295					300					
10	Trp	Ser	Ala	Pro	Ser	Asp	Pro	Leu	Asp	Ile	Leu	Ile	Thr	Gly	Gln	Phe	
	305					310					315					320	
	Tyr	Asp	Arg	Pro	Ser	Leu	Ser	Val	Gln	Pro	Val	Pro	Thr	Val	Ala	Pro	
					325					330					335		
15	Gly	Lys	Asn	Val	Thr	Leu	Leu	Cys	Gln	Ser	Arg	Gly	Gln	Phe	His	Thr	
				340					345					350			
	Phe	Leu	Leu	Thr	Lys	Glu	Gly	Ala	Gly	His	Pro	Pro	Leu	His	Leu	Arg	
			355					360					365				
20	Ser	Glu	His	Gln	Ala	Gln	Gln	Asn	Gln	Ala	Glu	Phe	Arg	Met	Gly	Pro	
		370					375					380					
	Val	Thr	Ser	Ala	His	Val	Gly	Thr	Tyr	Arg	Cys	Tyr	Ser	Ser	Leu	Ser	
	385					390					395					400	
25	Ser	Asn	Pro	Tyr	Leu	Leu	Ser	Leu	Pro	Ser	Asp	Pro	Leu	Glu	Leu	Val	
					405					410					415		
	Val	Ser	Glu	Ala	Ala	Glu	Thr	Leu	Ser	Pro	Ser	Gln	Asn	Lys	Thr	Asp	
				420					425					430			
30	Ser	Thr	Thr	Thr	Ser	Leu	Gly	Gln	His	Pro	Gln	Asp	Tyr	Thr	Val	Glu	
			435					440					445				
	Asn	Leu	Ile	Arg	Met	Gly	Val	Ala	Gly	Leu	Val	Leu	Val	Val	Leu	Gly	
		450					455					460					
35	Ile	Leu	Leu	Phe	Glu	Ala	Gln	His	Ser	Gln	Arg	Ser	Leu	Gln	Asp	Ala	
	465					470					475					480	
	Ala	Gly	Arg														
40	<210>	17															
	<211>	1625															
	<212>	DNA															
	<213>	human															
45	<220>																
	<221>	CDS															
	<222>	(30)..(1376)															
	<400>	17															
50	cacagctggg	gcccctggga	ggagacgcc	atg	atc	ccc	acc	ttc	acg	gct	ctg					53	
				Met	Ile	Pro	Thr	Phe	Thr	Ala	Leu						
					1				5								
	ctc	tgc	ctc	ggg	ctg	agt	ctg	ggc	ccc	agg	acc	cac	atg	cag	gca	ggg	101
55																	



EP 1 199 362 A2

	Leu	Cys	Leu	Gly	Leu	Ser	Leu	Gly	Pro	Arg	Thr	His	Met	Gln	Ala	Gly	
	10						15					20					
5	ccc	ctc	ccc	aaa	ccc	acc	ctc	tgg	gct	gag	cca	ggc	tct	gtg	atc	agc	149
	Pro	Leu	Pro	Lys	Pro	Thr	Leu	Trp	Ala	Glu	Pro	Gly	Ser	Val	Ile	Ser	
	25					30					35					40	
	tgg	ggg	aac	tct	gtg	acc	atc	tgg	tgt	cag	ggg	acc	ctg	gag	gct	cgg	197
10	Trp	Gly	Asn	Ser	Val	Thr	Ile	Trp	Cys	Gln	Gly	Thr	Leu	Glu	Ala	Arg	
					45					50					55		
	gag	tac	cgt	ctg	gat	aaa	gag	gaa	agc	cca	gca	ccc	tgg	gac	aga	cag	245
	Glu	Tyr	Arg	Leu	Asp	Lys	Glu	Glu	Ser	Pro	Ala	Pro	Trp	Asp	Arg	Gln	
				60					65					70			
15	aac	cca	ctg	gag	ccc	aag	aac	aag	gcc	aga	ttc	tcc	atc	cca	tcc	atg	293
	Asn	Pro	Leu	Pro	Lys	Asn	Lys	Ala	Arg	Phe	Ser	Ile	Pro	Ser	Met		
			75				80					85					
	aca	gag	gac	tat	gca	ggg	aga	tac	cgc	tgt	tac	tat	cgc	agc	cct	gta	341
20	Thr	Glu	Asp	Tyr	Ala	Gly	Arg	Tyr	Arg	Cys	Tyr	Tyr	Arg	Ser	Pro	Val	
	90					95						100					
	ggc	tgg	tca	cag	ccc	agt	gac	ccc	ctg	gag	ctg	gtg	atg	aca	gga	gcc	389
	Gly	Trp	Ser	Gln	Pro	Ser	Asp	Pro	Leu	Glu	Leu	Val	Met	Thr	Gly	Ala	
	105					110					115				120		
25	tac	agt	aaa	ccc	acc	ctt	tca	gcc	ctg	ccg	agt	cct	ctt	gtg	acc	tca	437
	Tyr	Ser	Lys	Pro	Thr	Leu	Ser	Ala	Leu	Pro	Ser	Pro	Leu	Val	Thr	Ser	
					125					130					135		
	gga	aag	agc	gtg	acc	ctg	ctg	tgt	cag	tca	cgg	agc	cca	atg	gac	act	485
30	Gly	Lys	Ser	Val	Thr	Leu	Leu	Cys	Gln	Ser	Arg	Ser	Pro	Met	Asp	Thr	
				140					145					150			
	ttt	ctt	ctg	atc	aag	gag	cgg	gca	gcc	cat	ccc	cta	ctg	cat	ctg	aga	533
	Phe	Leu	Leu	Ile	Lys	Glu	Arg	Ala	Ala	His	Pro	Leu	Leu	His	Leu	Arg	
				155				160					165				
35	tca	gag	cac	gga	gct	cag	cag	cac	cag	gct	gaa	ttc	ccc	atg	agt	cct	581
	Ser	Glu	His	Gly	Ala	Gln	Gln	His	Gln	Ala	Glu	Phe	Pro	Met	Ser	Pro	
				170			175					180					
	gtg	acc	tca	gtg	cac	ggg	ggg	acc	tac	agg	tgc	ttc	agc	tca	cac	ggc	629
40	Val	Thr	Ser	Val	His	Gly	Gly	Thr	Tyr	Arg	Cys	Phe	Ser	Ser	His	Gly	
	185					190					195					200	
	ttc	tcc	cac	tac	ctg	ctg	tca	cac	ccc	agt	gac	ccc	ctg	gag	ctc	ata	677
	Phe	Ser	His	Tyr	Leu	Leu	Ser	His	Pro	Ser	Asp	Pro	Leu	Glu	Leu	Ile	
					205					210					215		
45	gtc	tca	gga	tcc	ttg	gag	ggt	ccc	agg	ccc	tca	ccc	aca	agg	tcc	gtc	725
	Val	Ser	Gly	Ser	Leu	Glu	Gly	Pro	Arg	Pro	Ser	Pro	Thr	Arg	Ser	Val	
				220					225					230			
	tca	aca	gct	gca	ggc	cct	gag	gac	cag	ccc	ctc	atg	cct	aca	ggg	tca	773
50	Ser	Thr	Ala	Ala	Gly	Pro	Glu	Asp	Gln	Pro	Leu	Met	Pro	Thr	Gly	Ser	
				235				240					245				
	gtc	ccc	cac	agt	ggt	ctg	aga	agg	cac	tgg	gag	gta	ctg	atc	ggg	gtc	821
	Val	Pro	His	Ser	Gly	Leu	Arg	Arg	His	Trp	Glu	Val	Leu	Ile	Gly	Val	

55

EP 1 199 362 A2

	250	255	260	
5	ttg gtg gtc tcc atc ctg ctt ctc tcc ctc ctc ctc ttc ctc ctc ctc			869
	Leu Val Val Ser Ile Leu Leu Leu Ser Leu Leu Leu Phe Leu Leu Leu			
	265	270	275	280
	caa cac tgg cgt cag gga aaa cac agg aca ttg gcc cag aga cag gct			917
	Gln His Trp Arg Gln Gly Lys His Arg Thr Leu Ala Gln Arg Gln Ala			
		285	290	295
10	gat ttc caa cgt cct cca ggg gct gcc gag cca gag ccc aag gac ggg			965
	Asp Phe Gln Arg Pro Pro Gly Ala Ala Glu Pro Glu Pro Lys Asp Gly			
		300	305	310
15	ggc cta cag agg agg tcc agc cca gct gct gac gtc cag gga gaa aac			1013
	Gly Leu Gln Arg Arg Ser Ser Pro Ala Ala Asp Val Gln Gly Glu Asn			
		315	320	325
	ttc tgt gct gcc gtg aag aac aca cag cct gag gac ggg gtg gaa atg			1061
	Phe Cys Ala Ala Val Lys Asn Thr Gln Pro Glu Asp Gly Val Glu Met			
		330	335	340
20	gac act cgg cag agc cca cac gat gaa gac ccc cag gca gtg acg tat			1109
	Asp Thr Arg Gln Ser Pro His Asp Glu Asp Pro Gln Ala Val Thr Tyr			
		345	350	355
25	gcc aag gtg aaa cac tcc aga cct agg aga gaa atg gcc tct cct ccc			1157
	Ala Lys Val Lys His Ser Arg Pro Arg Glu Met Ala Ser Pro Pro			
		365	370	375
	tcc cca ctg tct ggg gaa ttc ctg gac aca aag gac aga cag gca gaa			1205
	Ser Pro Leu Ser Gly Glu Phe Leu Asp Thr Lys Asp Arg Gln Ala Glu			
		380	385	390
30	gag gac aga cag atg gac act gag gct gct gca tct gaa gcc ccc cag			1253
	Glu Asp Arg Gln Met Asp Thr Glu Ala Ala Ala Ser Glu Ala Pro Gln			
		395	400	405
35	gat gtg acc tac gcc cgg ctg cac agc ttt acc ctc aga cag aag gca			1301
	Asp Val Thr Tyr Ala Arg Leu His Ser Phe Thr Leu Arg Gln Lys Ala			
		410	415	420
	act gag cct cct cca tcc cag gaa ggg gcc tct cca gct gag ccc agt			1349
	Thr Glu Pro Pro Pro Ser Gln Glu Gly Ala Ser Pro Ala Glu Pro Ser			
		425	430	435
40	gtc tat gcc act ctg gcc atc cac taa tccagggggg acccagaccc			1396
	Val Tyr Ala Thr Leu Ala Ile His			
		445		
45	cacaagccat ggagactcag gaccccagaa ggcatggaag ctgcctccag tagacatcac			1456
	tgaaccccag ccagcccaga cccctgacac agaccactag aagattccgg gaacgttggg			1516
	agtcacctga ttctgcaaag ataaataata tccctgcatt atcaaaataa agtagcagac			1576
50	ctctcaattc acaatgagtt aactgataaa acaaaacaga agtcaaaaa			1625
	<210> 18			
	<211> 448			

55

EP 1 199 362 A2

<212> PRT  
<213> human

5  
<400> 18  
Met Ile Pro Thr Phe Thr Ala Leu Leu Cys Leu Gly Leu Ser Leu Gly  
1 5 10 15  
Pro Arg Thr His Met Gln Ala Gly Pro Leu Pro Lys Pro Thr Leu Trp  
10 20 25 30  
Ala Glu Pro Gly Ser Val Ile Ser Trp Gly Asn Ser Val Thr Ile Trp  
35 40 45  
Cys Gln Gly Thr Leu Glu Ala Arg Glu Tyr Arg Leu Asp Lys Glu Glu  
15 50 55 60  
Ser Pro Ala Pro Trp Asp Arg Gln Asn Pro Leu Glu Pro Lys Asn Lys  
65 70 75 80  
Ala Arg Phe Ser Ile Pro Ser Met Thr Glu Asp Tyr Ala Gly Arg Tyr  
20 85 90 95  
Arg Cys Tyr Tyr Arg Ser Pro Val Gly Trp Ser Gln Pro Ser Asp Pro  
100 105 110  
Leu Glu Leu Val Met Thr Gly Ala Tyr Ser Lys Pro Thr Leu Ser Ala  
25 115 120 125  
Leu Pro Ser Pro Leu Val Thr Ser Gly Lys Ser Val Thr Leu Leu Cys  
130 135 140  
Gln Ser Arg Ser Pro Met Asp Thr Phe Leu Leu Ile Lys Glu Arg Ala  
30 145 150 155 160  
Ala His Pro Leu Leu His Leu Arg Ser Glu His Gly Ala Gln Gln His  
165 170 175  
Gln Ala Glu Phe Pro Met Ser Pro Val Thr Ser Val His Gly Gly Thr  
35 180 185 190  
Tyr Arg Cys Phe Ser Ser His Gly Phe Ser His Tyr Leu Leu Ser His  
195 200 205  
Pro Ser Asp Pro Leu Glu Leu Ile Val Ser Gly Ser Leu Glu Gly Pro  
40 210 215 220  
Arg Pro Ser Pro Thr Arg Ser Val Ser Thr Ala Ala Gly Pro Glu Asp  
225 230 235 240  
Gln Pro Leu Met Pro Thr Gly Ser Val Pro His Ser Gly Leu Arg Arg  
45 245 250 255  
His Trp Glu Val Leu Ile Gly Val Leu Val Val Ser Ile Leu Leu Leu  
260 265 270  
Ser Leu Leu Leu Phe Leu Leu Leu Gln His Trp Arg Gln Gly Lys His  
50 275 280 285  
Arg Thr Leu Ala Gln Arg Gln Ala Asp Phe Gln Arg Pro Pro Gly Ala  
290 295 300

55

EP 1 199 362 A2

Ala Glu Pro Glu Pro Lys Asp Gly Gly Leu Gln Arg Arg Ser Ser Pro  
305 310 315 320

5 Ala Ala Asp Val Gln Gly Glu Asn Phe Cys Ala Ala Val Lys Asn Thr  
325 330 335

Gln Pro Glu Asp Gly Val Glu Met Asp Thr Arg Gln Ser Pro His Asp  
340 345 350

10 Glu Asp Pro Gln Ala Val Thr Tyr Ala Lys Val Lys His Ser Arg Pro  
355 360 365

Arg Arg Glu Met Ala Ser Pro Pro Ser Pro Leu Ser Gly Glu Phe Leu  
370 375 380

15 Asp Thr Lys Asp Arg Gln Ala Glu Glu Asp Arg Gln Met Asp Thr Glu  
385 390 395 400

Ala Ala Ala Ser Glu Ala Pro Gln Asp Val Thr Tyr Ala Arg Leu His  
405 410 415

20 Ser Phe Thr Leu Arg Gln Lys Ala Thr Glu Pro Pro Pro Ser Gln Glu  
420 425 430

Gly Ala Ser Pro Ala Glu Pro Ser Val Tyr Ala Thr Leu Ala Ile His  
435 440 445

25 <210> 19  
<211> 2194  
<212> DNA  
<213> human

30 <220>  
<221> CDS  
<222> (67)..(1962)

<400> 19  
tctctgtcct gccagcactg agggctcatc cctctgcaga gcgcggggtc accggaagga 60

35 gacgcc atg acg ccc gcc ctc aca gcc ctg ctc tgc ctt ggg ctg agt 108  
Met Thr Pro Ala Leu Thr Ala Leu Leu Cys Leu Gly Leu Ser  
1 5 10

40 ctg ggc ccc agg acc cgc gtg cag gca ggg ccc ttc ccc aaa ccc acc 156  
Leu Gly Pro Arg Thr Arg Val Gln Ala Gly Pro Phe Pro Lys Pro Thr  
15 20 25 30

ctc tgg gct gag cca ggc tct gtg atc agc tgg ggg agc ccc gtg acc 204  
Leu Trp Ala Glu Pro Gly Ser Val Ile Ser Trp Gly Ser Pro Val Thr  
35 40 45

45 atc tgg tgt cag ggg agc ctg gag gcc cag gag tac caa ctg gat aaa 252  
Ile Trp Cys Gln Gly Ser Leu Glu Ala Gln Glu Tyr Gln Leu Asp Lys  
50 55 60

50 gag gga agc cca gag ccc ttg gac aga aat aac cca ctg gaa ccc aag 300  
Glu Gly Ser Pro Glu Pro Leu Asp Arg Asn Asn Pro Leu Glu Pro Lys  
65 70 75

aac aag gcc aga ttc tcc atc cca tcc atg aca cag cac cat gca ggg 348

55

EP 1 199 362 A2

	Asn	Lys	Ala	Arg	Phe	Ser	Ile	Pro	Ser	Met	Thr	Gln	His	His	Ala	Gly	
		80					85					90					
5	aga	tac	cgc	tgc	cac	tat	tac	agc	tct	gca	ggc	tgg	tca	gag	ccc	agc	396
	Arg	Tyr	Arg	Cys	His	Tyr	Tyr	Ser	Ser	Ala	Gly	Trp	Ser	Glu	Pro	Ser	
	95					100					105					110	
	gac	ccc	ctg	gag	ctg	gtg	atg	aca	gga	gcc	tat	agc	aaa	ccc	acc	ctc	444
10	Asp	Pro	Leu	Glu	Leu	Val	Met	Thr	Gly	Ala	Tyr	Ser	Lys	Pro	Thr	Leu	
					115					120					125		
	tca	gcc	ctg	ccc	agc	cct	gtg	gtg	gcc	tca	ggg	ggg	aat	atg	acc	ctc	492
	Ser	Ala	Leu	Pro	Ser	Pro	Val	Val	Ala	Ser	Gly	Gly	Asn	Met	Thr	Leu	
				130					135					140			
15	cga	tgt	ggc	tca	cag	aag	aga	tat	cac	cat	ttt	gtt	ctg	atg	aag	gaa	540
	Arg	Cys	Gly	Ser	Gln	Lys	Arg	Tyr	His	His	Phe	Val	Leu	Met	Lys	Glu	
			145					150					155				
	gga	gaa	cac	cag	ctc	ccc	cgg	acc	ctg	gac	tca	cag	cag	ctc	cac	agt	588
20	Gly	Glu	His	Gln	Leu	Pro	Arg	Thr	Leu	Asp	Ser	Gln	Gln	Leu	His	Ser	
		160					165					170					
	ggg	ggg	ttc	cag	gcc	ctg	ttc	cct	gtg	ggc	ccc	gtg	aac	ccc	agc	cac	636
	Gly	Gly	Phe	Gln	Ala	Leu	Phe	Pro	Val	Gly	Pro	Val	Asn	Pro	Ser	His	
		175				180					185					190	
25	agg	tgg	agg	ttc	aca	tgc	tat	tac	tat	tat	atg	aac	acc	ccc	cgg	gtg	684
	Arg	Trp	Arg	Phe	Thr	Cys	Tyr	Tyr	Tyr	Tyr	Met	Asn	Thr	Pro	Arg	Val	
					195					200					205		
	tgg	tcc	cac	ccc	agt	gac	ccc	ctg	gag	att	ctg	ccc	tca	ggc	gtg	tct	732
30	Trp	Ser	His	Pro	Ser	Asp	Pro	Leu	Glu	Ile	Leu	Pro	Ser	Gly	Val	Ser	
				210					215					220			
	agg	aag	ccc	tcc	ctc	ctg	acc	ctg	cag	ggc	cct	gtc	ctg	gcc	cct	ggg	780
	Arg	Lys	Pro	Ser	Leu	Leu	Thr	Leu	Gln	Gly	Pro	Val	Leu	Ala	Pro	Gly	
			225					230					235				
35	cag	agt	ctg	acc	ctc	cag	tgt	ggc	tct	gat	gtc	ggc	tac	gac	aga	ttt	828
	Gln	Ser	Leu	Thr	Leu	Gln	Cys	Gly	Ser	Asp	Val	Gly	Tyr	Asp	Arg	Phe	
			240				245					250					
	gtt	ctg	tat	aag	gag	ggg	gaa	cgt	gac	ttc	ctc	cag	cgc	cct	ggc	cag	876
40	Val	Leu	Tyr	Lys	Glu	Gly	Glu	Arg	Asp	Phe	Leu	Gln	Arg	Pro	Gly	Gln	
		255				260					265					270	
	cag	ccc	cag	gct	ggg	ctc	tcc	cag	gcc	aac	ttc	acc	ctg	ggc	cct	gtg	924
	Gln	Pro	Gln	Ala	Gly	Leu	Ser	Gln	Ala	Asn	Phe	Thr	Leu	Gly	Pro	Val	
					275					280					285		
45	agc	ccc	tcc	aat	ggg	ggc	cag	tac	agg	tgc	tac	ggt	gca	cac	aac	ctc	972
	Ser	Pro	Ser	Asn	Gly	Gly	Gln	Tyr	Arg	Cys	Tyr	Gly	Ala	His	Asn	Leu	
				290					295					300			
	tcc	tcc	gag	tgg	tgc	gcc	ccc	agc	gac	ccc	ctg	aac	atc	ctg	atg	gca	1020
50	Ser	Ser	Glu	Trp	Ser	Ala	Pro	Ser	Asp	Pro	Leu	Asn	Ile	Leu	Met	Ala	
			305					310					315				
	gga	cag	atc	tat	gac	acc	gtc	tcc	ctg	tca	gca	cag	ccg	ggc	ccc	aca	1068
	Gly	Gln	Ile	Tyr	Asp	Thr	Val	Ser	Leu	Ser	Ala	Gln	Pro	Gly	Pro	Thr	

55

EP 1 199 362 A2

	320	325	330	
5	gtg gcc tca gga gag aac gtg acc ctg ctg tgt cag tca tgg tgg cag Val Ala Ser Gly Glu Asn Val Thr Leu Leu Cys Gln Ser Trp Trp Gln 335 340 345 350	1116		
	ttt gac act ttc ctt ctg acc aaa gaa ggg gca gcc cat ccc cca ctg Phe Asp Thr Phe Leu Leu Thr Lys Glu Gly Ala Ala His Pro Pro Leu 355 360 365	1164		
10	cgt ctg aga tca atg tac gga gct cat aag tac cag gct gaa ttc ccc Arg Leu Arg Ser Met Tyr Gly Ala His Lys Tyr Gln Ala Glu Phe Pro 370 375 380	1212		
15	atg agt cct gtg acc tca gcc cac gcg ggg acc tac agg tgc tac ggc Met Ser Pro Val Thr Ser Ala His Ala Gly Thr Tyr Arg Cys Tyr Gly 385 390 395	1260		
	tca cgc agc tcc aac ccc tac ctg ctg tct cac ccc agt gag ccc ctg Ser Arg Ser Ser Asn Pro Tyr Leu Leu Ser His Pro Ser Glu Pro Leu 400 405 410	1308		
20	gag ctc gtg gtc tca gga cac tct gga ggc tcc agc ctc cca ccc aca Glu Leu Val Val Ser Gly His Ser Gly Gly Ser Ser Leu Pro Pro Thr 415 420 425 430	1356		
25	ggg ccg ccc tcc aca cct ggt ctg gga aga tac ctg gag gtt ttg att Gly Pro Pro Ser Thr Pro Gly Leu Gly Arg Tyr Leu Glu Val Leu Ile 435 440 445	1404		
	ggg gtc tcg gtg gcc ttc gtc ctg ctg ctc ttc ctc ctc ctc ttc ctc Gly Val Ser Val Ala Phe Val Leu Leu Leu Phe Leu Leu Phe Leu 450 455 460	1452		
30	ctc ctc cga cgt cag cgt cac agc aaa cac agg aca tct gac cag aga Leu Leu Arg Arg Gln Arg His Ser Lys His Arg Thr Ser Asp Gln Arg 465 470 475	1500		
35	aag act gat ttc cag cgt cct gca ggg gct gcg gag aca gag ccc aag Lys Thr Asp Phe Gln Arg Pro Ala Gly Ala Ala Glu Thr Glu Pro Lys 480 485 490	1548		
	gac agg ggc ctg ctg agg agg tcc agc cca gct gct gac gtc cag gaa Asp Arg Gly Leu Leu Arg Arg Ser Ser Pro Ala Ala Asp Val Gln Glu 495 500 505 510	1596		
40	gaa aac ctc tat gct gcc gtg aag gac aca cag tct gag gac ggg gtg Glu Asn Leu Tyr Ala Ala Val Lys Asp Thr Gln Ser Glu Asp Gly Val 515 520 525	1644		
45	gag ctg gac agt cag agc cca cac gat gaa gac ccc cac gca gtg acg Glu Leu Asp Ser Gln Ser Pro His Asp Glu Asp Pro His Ala Val Thr 530 535 540	1692		
	tat gcc ccg gtg aaa cac tcc agt cct agg aga gaa atg gcc tct cct Tyr Ala Pro Val Lys His Ser Ser Pro Arg Arg Glu Met Ala Ser Pro 545 550 555	1740		
50	cct tcc cca ctg tct ggg gaa ttc ctg gac aca aag gac aga cag gca Pro Ser Pro Leu Ser Gly Glu Phe Leu Asp Thr Lys Asp Arg Gln Ala 560 565 570	1788		
55				

EP 1 199 362 A2

5 gaa gag gac aga cag atg gac act gag gct gct gca tct gaa gcc tcc 1836  
 Glu Glu Asp Arg Gln Met Asp Thr Glu Ala Ala Ala Ser Glu Ala Ser  
 575 580 585 590

cag gat gtg acc tac gcc cag ctg cac agc ttg acc ctt aga cgg aag 1884  
 Gln Asp Val Thr Tyr Ala Gln Leu His Ser Leu Thr Leu Arg Arg Lys  
 595 600 605

10 gca act gag cct cct cca tcc cag gaa ggg gaa cct cca gct gag ccc 1932  
 Ala Thr Glu Pro Pro Pro Ser Gln Glu Gly Glu Pro Pro Ala Glu Pro  
 610 615 620

15 agc atc tac gcc act ctg gcc atc cac tag cccgggggggt acgcagaccc 1982  
 Ser Ile Tyr Ala Thr Leu Ala Ile His  
 625 630

cacactcagc agaaggagac tcaggactgc tgaaggacgg gagctgcccc cagtggacac 2042

20 cagtgaaccc cagtcagcct ggaccacctaa cacagaccat gaggagacgc tgggaacttg 2102

tgggactcac ctgactcaaa gatgactaat atcgtcccat tttggaaata aagcaacaga 2162

cttctcaagc aggtcgtctc gttccaagat ct 2194

25 <210> 20  
 <211> 631  
 <212> PRT  
 <213> human

30 <400> 20  
 Met Thr Pro Ala Leu Thr Ala Leu Leu Cys Leu Gly Leu Ser Leu Gly  
 1 5 10 15

Pro Arg Thr Arg Val Gln Ala Gly Pro Phe Pro Lys Pro Thr Leu Trp  
 20 25 30

35 Ala Glu Pro Gly Ser Val Ile Ser Trp Gly Ser Pro Val Thr Ile Trp  
 35 40 45

Cys Gln Gly Ser Leu Glu Ala Gln Glu Tyr Gln Leu Asp Lys Glu Gly  
 50 55 60

40 Ser Pro Glu Pro Leu Asp Arg Asn Asn Pro Leu Glu Pro Lys Asn Lys  
 65 70 75 80

Ala Arg Phe Ser Ile Pro Ser Met Thr Gln His His Ala Gly Arg Tyr  
 85 90 95

45 Arg Cys His Tyr Tyr Ser Ser Ala Gly Trp Ser Glu Pro Ser Asp Pro  
 100 105 110

Leu Glu Leu Val Met Thr Gly Ala Tyr Ser Lys Pro Thr Leu Ser Ala  
 115 120 125

50 Leu Pro Ser Pro Val Val Ala Ser Gly Gly Asn Met Thr Leu Arg Cys  
 130 135 140

Gly Ser Gln Lys Arg Tyr His His Phe Val Leu Met Lys Glu Gly Glu  
 145 150 155 160

55

EP 1 199 362 A2

5 His Gln Leu Pro Arg Thr Leu Asp Ser Gln Gln Leu His Ser Gly Gly  
 165 170 175  
 Phe Gln Ala Leu Phe Pro Val Gly Pro Val Asn Pro Ser His Arg Trp  
 180 185 190  
 Arg Phe Thr Cys Tyr Tyr Tyr Tyr Met Asn Thr Pro Arg Val Trp Ser  
 195 200 205  
 10 His Pro Ser Asp Pro Leu Glu Ile Leu Pro Ser Gly Val Ser Arg Lys  
 210 215 220  
 Pro Ser Leu Leu Thr Leu Gln Gly Pro Val Leu Ala Pro Gly Gln Ser  
 225 230 235 240  
 15 Leu Thr Leu Gln Cys Gly Ser Asp Val Gly Tyr Asp Arg Phe Val Leu  
 245 250 255  
 Tyr Lys Glu Gly Glu Arg Asp Phe Leu Gln Arg Pro Gly Gln Gln Pro  
 260 265 270  
 20 Gln Ala Gly Leu Ser Gln Ala Asn Phe Thr Leu Gly Pro Val Ser Pro  
 275 280 285  
 Ser Asn Gly Gly Gln Tyr Arg Cys Tyr Gly Ala His Asn Leu Ser Ser  
 290 295 300  
 25 Glu Trp Ser Ala Pro Ser Asp Pro Leu Asn Ile Leu Met Ala Gly Gln  
 305 310 315 320  
 Ile Tyr Asp Thr Val Ser Leu Ser Ala Gln Pro Gly Pro Thr Val Ala  
 325 330 335  
 30 Ser Gly Glu Asn Val Thr Leu Leu Cys Gln Ser Trp Trp Gln Phe Asp  
 340 345 350  
 Thr Phe Leu Leu Thr Lys Glu Gly Ala Ala His Pro Pro Leu Arg Leu  
 355 360 365  
 35 Arg Ser Met Tyr Gly Ala His Lys Tyr Gln Ala Glu Phe Pro Met Ser  
 370 375 380  
 Pro Val Thr Ser Ala His Ala Gly Thr Tyr Arg Cys Tyr Gly Ser Arg  
 385 390 395 400  
 40 Ser Ser Asn Pro Tyr Leu Leu Ser His Pro Ser Glu Pro Leu Glu Leu  
 405 410 415  
 Val Val Ser Gly His Ser Gly Gly Ser Ser Leu Pro Pro Thr Gly Pro  
 420 425 430  
 45 Pro Ser Thr Pro Gly Leu Gly Arg Tyr Leu Glu Val Leu Ile Gly Val  
 435 440 445  
 Ser Val Ala Phe Val Leu Leu Leu Phe Leu Leu Phe Leu Leu Leu  
 450 455 460  
 50 Arg Arg Gln Arg His Ser Lys His Arg Thr Ser Asp Gln Arg Lys Thr  
 465 470 475 480

55



EP 1 199 362 A2

Asp Phe Gln Arg Pro Ala Gly Ala Ala Glu Thr Glu Pro Lys Asp Arg  
 485 490 495  
 5 Gly Leu Leu Arg Arg Ser Ser Pro Ala Ala Asp Val Gln Glu Glu Asn  
 500 505 510  
 Leu Tyr Ala Ala Val Lys Asp Thr Gln Ser Glu Asp Gly Val Glu Leu  
 515 520 525  
 10 Asp Ser Gln Ser Pro His Asp Glu Asp Pro His Ala Val Thr Tyr Ala  
 530 535 540  
 Pro Val Lys His Ser Ser Pro Arg Arg Glu Met Ala Ser Pro Pro Ser  
 545 550 555 560  
 15 Pro Leu Ser Gly Glu Phe Leu Asp Thr Lys Asp Arg Gln Ala Glu Glu  
 565 570 575  
 Asp Arg Gln Met Asp Thr Glu Ala Ala Ala Ser Glu Ala Ser Gln Asp  
 580 585 590  
 20 Val Thr Tyr Ala Gln Leu His Ser Leu Thr Leu Arg Arg Lys Ala Thr  
 595 600 605  
 Glu Pro Pro Pro Ser Gln Glu Gly Glu Pro Pro Ala Glu Pro Ser Ile  
 610 615 620  
 25 Tyr Ala Thr Leu Ala Ile His  
 625 630  
 <210> 21  
 <211> 2061  
 30 <212> DNA  
 <213> human  
 <220>  
 <221> CDS  
 <222> (67)..(1839)  
 35 <400> 21  
 tttgtgtcct gccaggcacc gtggtctcat ccgcctgcac agctgagtc agtgggagct 60  
 gaggcc atg acc ctc acc ctc tca gtc ctg att tgc ctc ggg ctg agt 108  
 Met Thr Leu Thr Leu Ser Val Leu Ile Cys Leu Gly Leu Ser  
 1 5 10  
 40 gtg ggc ccc agg acc tgc gtg cag gca ggc acc ctc ccc aaa ccc acc 156  
 Val Gly Pro Arg Thr Cys Val Gln Ala Gly Thr Leu Pro Lys Pro Thr  
 15 20 25 30  
 45 ctc tgg gct gag cca gcc tct gtg ata gct cgg ggg aag ccc gtg acc 204  
 Leu Trp Ala Glu Pro Ala Ser Val Ile Ala Arg Gly Lys Pro Val Thr  
 35 40 45  
 ctc tgg tgt cag ggg ccc ctg gag act gag gag tac cgt ctg gat aag 252  
 50 Leu Trp Cys Gln Gly Pro Leu Glu Thr Glu Tyr Arg Leu Asp Lys  
 50 55 60  
 gag gga ctc cca tgg gcc cgg aag aga cag aac cca ctg gag cct gga 300  
 Glu Gly Leu Pro Trp Ala Arg Lys Arg Gln Asn Pro Leu Glu Pro Gly  
 65 70 75  
 55

EP 1 199 362 A2

5	gcc aag gcc aag ttc cac att cca tcc acg gtg tat gac agt gca ggg Ala Lys Ala Lys Phe His Ile Pro Ser Thr Val Tyr Asp Ser Ala Gly 80 85 90	348
	cga tac cgc tgc tac tat gag acc cct gca ggc tgg tca gag ccc agt Arg Tyr Arg Cys Tyr Tyr Glu Thr Pro Ala Gly Trp Ser Glu Pro Ser 95 100 105 110	396
10	gac ccc ctg gag ctg gtg gcg aca gga ttc tat gca gaa ccc act ctt Asp Pro Leu Glu Leu Val Ala Thr Gly Phe Tyr Ala Glu Pro Thr Leu 115 120 125	444
15	tta gcc ctg ccg agt cct gtg gtg gcc tca gga gga aat gtg acc ctc Leu Ala Leu Pro Ser Pro Val Val Ala Ser Gly Gly Asn Val Thr Leu 130 135 140	492
	cag tgt gat aca ctg gac gga ctt ctc acg ttt gtt ctt gtt gag gaa Gln Cys Asp Thr Leu Asp Gly Leu Leu Thr Phe Val Leu Val Glu Glu 145 150 155	540
20	gaa cag aag ctc ccc agg acc ctg tac tca cag aag ctc ccc aaa ggg Glu Gln Lys Leu Pro Arg Thr Leu Tyr Ser Gln Lys Leu Pro Lys Gly 160 165 170	588
25	cca tcc cag gcc ctg ttc cct gtg ggt ccc gtg acc ccc agc tgc agg Pro Ser Gln Ala Leu Phe Pro Val Gly Pro Val Thr Pro Ser Cys Arg 175 180 185 190	636
	tgg agg ttc aga tgc tat tac tat tac agg aaa aac cct cag gtg tgg Trp Arg Phe Arg Cys Tyr Tyr Tyr Tyr Arg Lys Asn Pro Gln Val Trp 195 200 205	684
30	tcg aac ccc agt gac ctc ctg gag att ctg gtc cca ggc gtg tct agg Ser Asn Pro Ser Asp Leu Leu Glu Ile Leu Val Pro Gly Val Ser Arg 210 215 220	732
35	aag ccc tcc ctc ctg atc ccg cag ggc tct gtc gtg gcc cgc gga ggc Lys Pro Ser Leu Leu Ile Pro Gln Gly Ser Val Val Ala Arg Gly Gly 225 230 235	780
	agc ctg acc ctg cag tgt cgc tct gat gtc ggc tat gac ata ttc gtt Ser Leu Thr Leu Gln Cys Arg Ser Asp Val Gly Tyr Asp Ile Phe Val 240 245 250	828
40	ctg tac aag gag ggg gaa cat gac ctc gtc cag ggc tct ggc cag cag Leu Tyr Lys Glu Gly Glu His Asp Leu Val Gln Gly Ser Gly Gln Gln 255 260 265 270	876
45	ccc cag gct ggg ctc tcc cag gcc aac ttc acc ctg ggc cct gtg agc Pro Gln Ala Gly Leu Ser Gln Ala Asn Phe Thr Leu Gly Pro Val Ser 275 280 285	924
	cgc tcc cac ggg ggc cag tac aga tgc tac ggt gca cac aac ctc tcc Arg Ser His Gly Gly Gln Tyr Arg Cys Tyr Gly Ala His Asn Leu Ser 290 295 300	972
50	cct agg tgg tcg gcc ccc agc gac ccc ctg gac atc ctg atc gca gga Pro Arg Trp Ser Ala Pro Ser Asp Pro Leu Asp Ile Leu Ile Ala Gly 305 310 315	1020

55

EP 1 199 362 A2

5	ctg atc cct gac ata ccc gcc ctc tcg gtg cag ccg ggc ccc aag gtg Leu Ile Pro Asp Ile Pro Ala Leu Ser Val Gln Pro Gly Pro Lys Val 320 325 330	1068
10	gcc tca gga gag aac gtg acc ctg ctg tgt cag tca tgg cat cag ata Ala Ser Gly Glu Asn Val Thr Leu Leu Cys Gln Ser Trp His Gln Ile 335 340 345 350	1116
15	gac act ttc ttt ttg acc aag gag ggg gca gcc cat ccc ccg ctg tgt Asp Thr Phe Phe Leu Thr Lys Glu Gly Ala Ala His Pro Pro Leu Cys 355 360 365	1164
20	cta aag tca aag tac cag tct tat aga cac cag gct gaa ttc tcc atg Leu Lys Ser Lys Tyr Gln Ser Tyr Arg His Gln Ala Glu Phe Ser Met 370 375 380	1212
25	agt cct gtg acc tca gcc cag ggt gga acc tac cga tgc tac agc gca Ser Pro Val Thr Ser Ala Gln Gly Gly Thr Tyr Arg Cys Tyr Ser Ala 385 390 395	1260
30	atc agg tcc tac ccc tac ctg ctg tcc agc cct agt tac ccc cag gag Ile Arg Ser Tyr Pro Tyr Leu Leu Ser Ser Pro Ser Tyr Pro Gln Glu 400 405 410	1308
35	ctc gtg gtc tca gga ccc tct ggg gat ccc agc ctc tca cct aca ggc Leu Val Val Ser Gly Pro Ser Gly Asp Pro Ser Leu Ser Pro Thr Gly 415 420 425 430	1356
40	tcc acc ccc aca cct ggc cct gag gac cag ccc ctc acc ccc acg ggg Ser Thr Pro Thr Pro Gly Pro Glu Asp Gln Pro Leu Thr Pro Thr Gly 435 440 445	1404
45	ttg gat ccc cag agt ggt ctg gga agg cac ctg ggg gtt gtg act ggg Leu Asp Pro Gln Ser Gly Leu Gly Arg His Leu Gly Val Val Thr Gly 450 455 460	1452
50	gtc tca gtg gcc ttc gtc ctg ctg ctg ttc ctc ctc ctc ttc ctc ctc Val Ser Val Ala Phe Val Leu Leu Leu Phe Leu Leu Phe Leu Leu 465 470 475	1500
55	ctc cga cat cgg cat cag agc aaa cac agg aca tcg gcc cat ttc tac Leu Arg His Arg His Gln Ser Lys His Arg Thr Ser Ala His Phe Tyr 480 485 490	1548
60	cgt cct gca ggg gct gcg ggg cca gag ccc aag gac cag ggc ctg cag Arg Pro Ala Gly Ala Ala Gly Pro Glu Pro Lys Asp Gln Gly Leu Gln 495 500 505 510	1596
65	aag agg gcc agc cca gtt gct gac atc cag gag gaa att ctc aat gct Lys Arg Ala Ser Pro Val Ala Asp Ile Gln Glu Glu Ile Leu Asn Ala 515 520 525	1644
70	gcc gtg aag gac aca cag ccc aag gac ggg gtg gag atg gat gct cgg Ala Val Lys Asp Thr Gln Pro Lys Asp Gly Val Glu Met Asp Ala Arg 530 535 540	1692
75	gct gct gca tct gaa gcc ccc cag gat gtg acc tac gcc cag ctg cac Ala Ala Ala Ser Glu Ala Pro Gln Asp Val Thr Tyr Ala Gln Leu His 545 550 555	1740
80	agc ttg acc ctc aga cgg gag gca act gag cct cct cca tcc cag gaa	1788

EP 1 199 362 A2

Ser Leu Thr Leu Arg Arg Glu Ala Thr Glu Pro Pro Pro Ser Gln Glu  
560 565 570

5 agg gaa cct cca gct gaa ccc agc atc tac gcc ccc ctg gcc atc cac 1836  
Arg Glu Pro Pro Ala Glu Pro Ser Ile Tyr Ala Pro Leu Ala Ile His  
575 580 585 590

tag cccacggggg acccagatct catactcaac agaaggagac tcagagactc 1889

10 cagaaggcac aggagctgcc cccagtggac accaatgaac cccagccagc ctggaccct 1949  
aacaagacc accaggacat cctgggaact ctgggactca ctagattctg cagtcaaaga 2009  
tgactaatat ccttgcattt ttgaaatgaa gccacagact tctcaataaa tc 2061

15 <210> 22  
<211> 590  
<212> PRT  
<213> human

20 <400> 22  
Met Thr Leu Thr Leu Ser Val Leu Ile Cys Leu Gly Leu Ser Val Gly  
1 5 10 15

Pro Arg Thr Cys Val Gln Ala Gly Thr Leu Pro Lys Pro Thr Leu Trp  
20 25 30

25 Ala Glu Pro Ala Ser Val Ile Ala Arg Gly Lys Pro Val Thr Leu Trp  
35 40 45

Cys Gln Gly Pro Leu Glu Thr Glu Glu Tyr Arg Leu Asp Lys Glu Gly  
50 55 60

30 Leu Pro Trp Ala Arg Lys Arg Gln Asn Pro Leu Glu Pro Gly Ala Lys  
65 70 75 80

Ala Lys Phe His Ile Pro Ser Thr Val Tyr Asp Ser Ala Gly Arg Tyr  
85 90 95

35 Arg Cys Tyr Tyr Glu Thr Pro Ala Gly Trp Ser Glu Pro Ser Asp Pro  
100 105 110

Leu Glu Leu Val Ala Thr Gly Phe Tyr Ala Glu Pro Thr Leu Leu Ala  
115 120 125

40 Leu Pro Ser Pro Val Val Ala Ser Gly Gly Asn Val Thr Leu Gln Cys  
130 135 140

Asp Thr Leu Asp Gly Leu Leu Thr Phe Val Leu Val Glu Glu Glu Gln  
145 150 155 160

45 Lys Leu Pro Arg Thr Leu Tyr Ser Gln Lys Leu Pro Lys Gly Pro Ser  
165 170 175

Gln Ala Leu Phe Pro Val Gly Pro Val Thr Pro Ser Cys Arg Trp Arg  
180 185 190

50 Phe Arg Cys Tyr Tyr Tyr Tyr Arg Lys Asn Pro Gln Val Trp Ser Asn  
195 200 205

55

EP 1 199 362 A2

Pro Ser Asp Leu Leu Glu Ile Leu Val Pro Gly Val Ser Arg Lys Pro  
 210 215 220  
 5  
 Ser Leu Leu Ile Pro Gln Gly Ser Val Val Ala Arg Gly Gly Ser Leu  
 225 230 235 240  
 Thr Leu Gln Cys Arg Ser Asp Val Gly Tyr Asp Ile Phe Val Leu Tyr  
 245 250 255  
 10  
 Lys Glu Gly Glu His Asp Leu Val Gln Gly Ser Gly Gln Gln Pro Gln  
 260 265 270  
 Ala Gly Leu Ser Gln Ala Asn Phe Thr Leu Gly Pro Val Ser Arg Ser  
 275 280 285  
 15  
 His Gly Gly Gln Tyr Arg Cys Tyr Gly Ala His Asn Leu Ser Pro Arg  
 290 295 300  
 Trp Ser Ala Pro Ser Asp Pro Leu Asp Ile Leu Ile Ala Gly Leu Ile  
 305 310 315 320  
 20  
 Pro Asp Ile Pro Ala Leu Ser Val Gln Pro Gly Pro Lys Val Ala Ser  
 325 330 335  
 Gly Glu Asn Val Thr Leu Leu Cys Gln Ser Trp His Gln Ile Asp Thr  
 340 345 350  
 25  
 Phe Phe Leu Thr Lys Glu Gly Ala Ala His Pro Pro Leu Cys Leu Lys  
 355 360 365  
 Ser Lys Tyr Gln Ser Tyr Arg His Gln Ala Glu Phe Ser Met Ser Pro  
 370 375 380  
 30  
 Val Thr Ser Ala Gln Gly Gly Thr Tyr Arg Cys Tyr Ser Ala Ile Arg  
 385 390 395 400  
 Ser Tyr Pro Tyr Leu Leu Ser Ser Pro Ser Tyr Pro Gln Glu Leu Val  
 405 410 415  
 35  
 Val Ser Gly Pro Ser Gly Asp Pro Ser Leu Ser Pro Thr Gly Ser Thr  
 420 425 430  
 Pro Thr Pro Gly Pro Glu Asp Gln Pro Leu Thr Pro Thr Gly Leu Asp  
 435 440 445  
 40  
 Pro Gln Ser Gly Leu Gly Arg His Leu Gly Val Val Thr Gly Val Ser  
 450 455 460  
 Val Ala Phe Val Leu Leu Leu Phe Leu Leu Leu Phe Leu Leu Leu Arg  
 465 470 475 480  
 45  
 His Arg His Gln Ser Lys His Arg Thr Ser Ala His Phe Tyr Arg Pro  
 485 490 495  
 Ala Gly Ala Ala Gly Pro Glu Pro Lys Asp Gln Gly Leu Gln Lys Arg  
 500 505 510  
 50  
 Ala Ser Pro Val Ala Asp Ile Gln Glu Glu Ile Leu Asn Ala Ala Val  
 515 520 525  
 Lys Asp Thr Gln Pro Lys Asp Gly Val Glu Met Asp Ala Arg Ala Ala  
 55

# EP 1 199 362 A2

	530	535	540	
	Ala Ser Glu Ala Pro Gln Asp Val Thr Tyr Ala Gln Leu His Ser Leu			
5	545	550	555	560
	Thr Leu Arg Arg Glu Ala Thr Glu Pro Pro Pro Ser Gln Glu Arg Glu			
		565	570	575
10	Pro Pro Ala Glu Pro Ser Ile Tyr Ala Pro Leu Ala Ile His			
		580	585	590
	<210> 23			
	<211> 28			
15	<212> DNA			
	<213> human			
	<400> 23			
	tatgcggccg ccatgatgac aatgtggt			28
20	<210> 24			
	<211> 25			
	<212> DNA			
	<213> human			
	<400> 24			
25	tatgcggccg ccccttgcca tagcg			25
	<210> 25			
	<211> 31			
30	<212> DNA			
	<213> human			
	<400> 25			
	atagtcgaca acgccatcat gagatgtggt g			31
35	<210> 26			
	<211> 29			
	<212> DNA			
	<213> human			
	<400> 26			
40	taaagatctg ggctcgtag ctgtcgggt			29
	<210> 27			
	<211> 33			
45	<212> DNA			
	<213> human			
	<400> 27			
	tatagatcta cccccagggtg ccttcccaga cca			33
50	<210> 28			
	<211> 42			
	<212> PRT			
	<213> human			
55				

# EP 1 199 362 A2

5	<p>&lt;220&gt;            &lt;221&gt; VARIANT            &lt;222&gt; (2)            &lt;223&gt; Xaa is Gly or Arg</p>
10	<p>&lt;220&gt;            &lt;221&gt; VARIANT            &lt;222&gt; (5)            &lt;223&gt; Xaa is Leu or Val</p>
15	<p>&lt;220&gt;            &lt;221&gt; VARIANT            &lt;222&gt; (6)            &lt;223&gt; Xaa is Gly or Asp</p>
20	<p>&lt;220&gt;            &lt;221&gt; VARIANT            &lt;222&gt; (10)            &lt;223&gt; Xaa is His, Arg or Cys</p>
25	<p>&lt;220&gt;            &lt;221&gt; VARIANT            &lt;222&gt; (11)            &lt;223&gt; Xaa is Val or Met</p>
30	<p>&lt;220&gt;            &lt;221&gt; VARIANT            &lt;222&gt; (13)            &lt;223&gt; Xaa is Ala or Thr</p>
35	<p>&lt;220&gt;            &lt;221&gt; VARIANT            &lt;222&gt; (15)            &lt;223&gt; Xaa is His, Pro or Thr</p>
40	<p>&lt;220&gt;            &lt;221&gt; VARIANT            &lt;222&gt; (16)            &lt;223&gt; Xaa is Leu, Ile or Phe</p>
45	<p>&lt;220&gt;            &lt;221&gt; VARIANT            &lt;222&gt; (18)            &lt;223&gt; Xaa is Gly, Asp or Ala</p>
50	<p>&lt;220&gt;            &lt;221&gt; VARIANT            &lt;222&gt; (26)            &lt;223&gt; Xaa is Thr, Ile, Ser or Ala</p>
55	

<220>  
 <221> VARIANT  
 <222> (30)  
 5 <223> Xaa is Thr, Ile, Ser or Ala

<220>  
 <221> VARIANT  
 <222> (31)  
 10 <223> Xaa is a sequence of 70 amino acids

<220>  
 <221> VARIANT  
 <222> (37)  
 15 <223> Xaa is Gly or Val

<220>  
 <221> VARIANT  
 <222> (40)  
 20 <223> Xaa is Met or Ala

<400> 28  
 Leu Xaa Leu Ser Xaa Xaa Pro Arg Thr Xaa Xaa Gln Xaa Gly Xaa Xaa  
 1 5 10 15  
 25 Pro Xaa Pro Thr Leu Trp Ala Glu Pro Xaa Ser Phe Ile Xaa Xaa Ser  
 20 25 30  
 Asp Pro Lys Leu Xaa Leu Val Xaa Thr Gly  
 35 40  
 30 <210> 29  
 <211> 1016  
 <212> DNA  
 <213> human

<220>  
 35 <221> CDS  
 <222> (69)..(968)

<400> 29  
 ctgagtcctgc ctgtggcatg gacctgcac ttccctgaag catctccagg gctgaaaaat 60  
 40 cactgacc atg gca cca tgg tct cat cca tct gca cag ctg cag cca gtg 110  
 Met Ala Pro Trp Ser His Pro Ser Ala Gln Leu Gln Pro Val  
 1 5 10  
 gga gga gac gcc gtg agc cct gcc ctc atg gtt ctg ctc tgc ctc ggg 158  
 45 Gly Gly Asp Ala Val Ser Pro Ala Leu Met Val Leu Leu Cys Leu Gly  
 15 20 25 30  
 ctg agt ctg ggc ccc agg acc cac gtg cag gca ggg aac ctc tcc aaa 206  
 Leu Ser Leu Gly Pro Arg Thr His Val Gln Ala Gly Asn Leu Ser Lys  
 35 40 45  
 50 gcc acc ctc tgg gct gag cca ggc tct gtg atc agc cgg ggg aac tct 254  
 Ala Thr Leu Trp Ala Glu Pro Gly Ser Val Ile Ser Arg Gly Asn Ser  
 50 55 60  
 55



EP 1 199 362 A2

5	gtg acc atc cgg tgt cag ggg acc ctg gag gcc cag gaa tac cgt ctg Val Thr Ile Arg Cys Gln Gly Thr Leu Glu Ala Gln Glu Tyr Arg Leu 65 70 75	302
10	gtt aaa gag gga agc cca gaa ccc tgg gac aca cag aac cca ctg gag Val Lys Glu Gly Ser Pro Glu Pro Trp Asp Thr Gln Asn Pro Leu Glu 80 85 90	350
15	ccc aag aac aag gcc aga ttc tcc atc cca tcc atg aca gag cac cat Pro Lys Asn Lys Ala Arg Phe Ser Ile Pro Ser Met Thr Glu His His 95 100 105 110	398
20	gca ggg aga tac cgc tgt tac tac tac agc cct gca ggc tgg tca gag Ala Gly Arg Tyr Arg Cys Tyr Tyr Tyr Ser Pro Ala Gly Trp Ser Glu 115 120 125	446
25	ccc agc gac ccc ctg gag ctg gtg gtg aca gga ttc tac aac aaa ccc Pro Ser Asp Pro Leu Glu Leu Val Thr Gly Phe Tyr Asn Lys Pro 130 135 140	494
30	acc ctc tca gcc ctg ccc agt cct gtg gtg acc tca gga gag aac gtg Thr Leu Ser Ala Leu Pro Ser Pro Val Val Thr Ser Gly Glu Asn Val 145 150 155	542
35	acc ctc cag tgt ggc tca cgg ctg aga ttc gac agg ttc att ctg act Thr Leu Gln Cys Gly Ser Arg Leu Arg Phe Asp Arg Phe Ile Leu Thr 160 165 170	590
40	gag gaa gga gac cac aag ctc tcc tgg acc ttg gac tca cag ctg acc Glu Glu Gly Asp His Lys Leu Ser Trp Thr Leu Asp Ser Gln Leu Thr 175 180 185 190	638
45	ccc agt ggg cag ttc cag gcc ctg ttc cct gtg ggc cct gtg acc ccc Pro Ser Gly Gln Phe Gln Ala Leu Phe Pro Val Gly Pro Val Thr Pro 195 200 205	686
50	agc cac agg tgg atg ctc aga tgc tat ggc tct cgc agg cat atc ctg Ser His Arg Trp Met Leu Arg Cys Tyr Gly Ser Arg Arg His Ile Leu 210 215 220	734
55	cag gta tgg tca gaa ccc agt gac ctc ctg gag att ccg gtc tca gga Gln Val Trp Ser Glu Pro Ser Asp Leu Leu Glu Ile Pro Val Ser Gly 225 230 235	782
60	gca gct gat aac ctc agt ccg tca caa aac aag tct gac tct ggg act Ala Ala Asp Asn Leu Ser Pro Ser Gln Asn Lys Ser Asp Ser Gly Thr 240 245 250	830
65	gcc tca cac ctt cag gat tac gca gta gag aat ctc atc cgc atg ggc Ala Ser His Leu Gln Asp Tyr Ala Val Glu Asn Leu Ile Arg Met Gly 255 260 265 270	878
70	atg gcc ggc ttg atc ctg gtg gtc ctt ggg att ctg ata ttt cag gat Met Ala Gly Leu Ile Leu Val Val Leu Gly Ile Leu Ile Phe Gln Asp 275 280 285	926
75	tgg cac agc cag aga agc ccc caa gct gca gct gga agg tga Trp His Ser Gln Arg Ser Pro Gln Ala Ala Ala Gly Arg 290 295 300	968
80	acagaagaga gaacaatgca ccattgaatg ctggagcctt ggaagcga	1016

EP 1 199 362 A2

5 <210> 30  
 <211> 299  
 <212> PRT  
 <213> human  
  
 <400> 30  
 10 Met Ala Pro Trp Ser His Pro Ser Ala Gln Leu Gln Pro Val Gly Gly  
 1 5 10 15  
 Asp Ala Val Ser Pro Ala Leu Met Val Leu Leu Cys Leu Gly Leu Ser  
 20 25 30  
 15 Leu Gly Pro Arg Thr His Val Gln Ala Gly Asn Leu Ser Lys Ala Thr  
 35 40 45  
 Leu Trp Ala Glu Pro Gly Ser Val Ile Ser Arg Gly Asn Ser Val Thr  
 50 55 60  
 20 Ile Arg Cys Gln Gly Thr Leu Glu Ala Gln Glu Tyr Arg Leu Val Lys  
 65 70 75 80  
 Glu Gly Ser Pro Glu Pro Trp Asp Thr Gln Asn Pro Leu Glu Pro Lys  
 85 90 95  
 25 Asn Lys Ala Arg Phe Ser Ile Pro Ser Met Thr Glu His His Ala Gly  
 100 105 110  
 Arg Tyr Arg Cys Tyr Tyr Tyr Ser Pro Ala Gly Trp Ser Glu Pro Ser  
 115 120 125  
 30 Asp Pro Leu Glu Leu Val Val Thr Gly Phe Tyr Asn Lys Pro Thr Leu  
 130 135 140  
 Ser Ala Leu Pro Ser Pro Val Val Thr Ser Gly Glu Asn Val Thr Leu  
 145 150 155 160  
 35 Gln Cys Gly Ser Arg Leu Arg Phe Asp Arg Phe Ile Leu Thr Glu Glu  
 165 170 175  
 Gly Asp His Lys Leu Ser Trp Thr Leu Asp Ser Gln Leu Thr Pro Ser  
 180 185 190  
 40 Gly Gln Phe Gln Ala Leu Phe Pro Val Gly Pro Val Thr Pro Ser His  
 195 200 205  
 Arg Trp Met Leu Arg Cys Tyr Gly Ser Arg Arg His Ile Leu Gln Val  
 210 215 220  
 45 Trp Ser Glu Pro Ser Asp Leu Leu Glu Ile Pro Val Ser Gly Ala Ala  
 225 230 235 240  
 Asp Asn Leu Ser Pro Ser Gln Asn Lys Ser Asp Ser Gly Thr Ala Ser  
 245 250 255  
 50 His Leu Gln Asp Tyr Ala Val Glu Asn Leu Ile Arg Met Gly Met Ala  
 260 265 270  
 Gly Leu Ile Leu Val Val Leu Gly Ile Leu Ile Phe Gln Asp Trp His  
 275 280 285  
 55

EP 1 199 362 A2

5 Ser Gln Arg Ser Pro Gln Ala Ala Ala Gly Arg  
290 295

<210> 31  
<211> 1007  
<212> DNA  
10 <213> human

<220>  
<221> CDS  
<222> (95)..(958)

15 <400> 31  
cagggtgtcag atgtgtctct gctgatctga gtctgcctgt ggcatggacc tgcattcttc 60  
ctgaagcatc tccagggctg aaaaatcact gacc atg gca cca tgg tct cat cca 115  
Met Ala Pro Trp Ser His Pro  
1 5

20 tct gca cag ctg cag cca gtg gga gga gac gcc gtg agc cct gcc ctc 163  
Ser Ala Gln Leu Gln Pro Val Gly Gly Asp Ala Val Ser Pro Ala Leu  
10 15 20

25 atg gtt ctg ctc tgc ctc ggg aac ctc tcc aaa gcc acc ctc tgg gct 211  
Met Val Leu Leu Cys Leu Gly Asn Leu Ser Lys Ala Thr Leu Trp Ala  
25 30 35

30 gag cca ggc tct gtg atc agc cgg ggg aac tct gtg acc atc cgg tgt 259  
Glu Pro Gly Ser Val Ile Ser Arg Gly Asn Ser Val Thr Ile Arg Cys  
40 45 50 55

35 cag ggg acc ctg gag gcc cag gaa tac cgt ctg gtt aaa gag gga agc 307  
Gln Gly Thr Leu Glu Ala Gln Glu Tyr Arg Leu Val Lys Glu Gly Ser  
60 65 70

40 cca gaa ccc tgg gac aca cag aac cca ctg gag ccc aag aac aag gcc 355  
Pro Glu Pro Trp Asp Thr Gln Asn Pro Leu Glu Pro Lys Asn Lys Ala  
75 80 85

45 aga ttc tcc atc cca tcc atg aca gag cac cat gca ggg aga tac cgc 403  
Arg Phe Ser Ile Pro Ser Met Thr Glu His His Ala Gly Arg Tyr Arg  
90 95 100

50 tgt tac tac tac agc cct gca ggc tgg tca gag ccc agc gac ccc ctg 451  
Cys Tyr Tyr Tyr Ser Pro Ala Gly Trp Ser Glu Pro Ser Asp Pro Leu  
105 110 115

55 gag ctg gtg gtg aca gga ttc tac aac aaa ccc acc ctc tca gcc ctg 499  
Glu Leu Val Val Thr Gly Phe Tyr Asn Lys Pro Thr Leu Ser Ala Leu  
120 125 130 135

ccc agt cct gtg gtg acc tca gga gag aac gtg acc ctc cag tgt ggc 547  
Pro Ser Pro Val Val Thr Ser Gly Glu Asn Val Thr Leu Gln Cys Gly  
140 145 150

tca cgg ctg aga ttc gac agg ttc att ctg act gag gaa gga gac cac 595  
Ser Arg Leu Arg Phe Asp Arg Phe Ile Leu Thr Glu Glu Gly Asp His  
155 160 165

EP 1 199 362 A2

aag ctc tcc tgg acc ttg gac tca cag ctg acc ccc agt ggg cag ttc 643  
 Lys Leu Ser Trp Thr Leu Asp Ser Gln Leu Thr Pro Ser Gly Gln Phe  
 170 175 180

5

cag gcc ctg ttc cct gtg ggc cct gtg acc ccc agc cac agg tgg atg 691  
 Gln Ala Leu Phe Pro Val Gly Pro Val Thr Pro Ser His Arg Trp Met  
 185 190 195

10

ctc aga tgc tat ggc tct cgc agg cat atc ctg cag gta tgg tca gaa 739  
 Leu Arg Cys Tyr Gly Ser Arg Arg His Ile Leu Gln Val Trp Ser Glu  
 200 205 210 215

15

ccc agt gac ctc ctg gag att ccg gtc tca gga gca gct gat aac ctc 787  
 Pro Ser Asp Leu Leu Glu Ile Pro Val Ser Gly Ala Ala Asp Asn Leu  
 220 225 230

20

agt ccg tca caa aac aag tct gac tct ggg act gcc tca cac ctt cag 835  
 Ser Pro Ser Gln Asn Lys Ser Asp Ser Gly Thr Ala Ser His Leu Gln  
 235 240 245

25

gat tac gca gta gag aat ctc atc cgc atg ggc atg gcc ggc ttg atc 883  
 Asp Tyr Ala Val Glu Asn Leu Ile Arg Met Gly Met Ala Gly Leu Ile  
 250 255 260

30

ctg gtg gtc ctt ggg att ctg ata ttt cag gat tgg cac agc cag aga 931  
 Leu Val Val Leu Gly Ile Leu Ile Phe Gln Asp Trp His Ser Gln Arg  
 265 270 275

35

agc ccc caa gct gca gct gga agg tga acagaagaga gaacaatgca 978  
 Ser Pro Gln Ala Ala Gly Arg  
 280 285

40

ccattgaatg ctggagcctt ggaagcgaa 1007

<210> 32  
 <211> 287  
 <212> PRT  
 <213> human

<400> 32  
 Met Ala Pro Trp Ser His Pro Ser Ala Gln Leu Gln Pro Val Gly Gly  
 1 5 10 15

40

Asp Ala Val Ser Pro Ala Leu Met Val Leu Leu Cys Leu Gly Asn Leu  
 20 25 30

Ser Lys Ala Thr Leu Trp Ala Glu Pro Gly Ser Val Ile Ser Arg Gly  
 35 40 45

45

Asn Ser Val Thr Ile Arg Cys Gln Gly Thr Leu Glu Ala Gln Glu Tyr  
 50 55 60

Arg Leu Val Lys Glu Gly Ser Pro Glu Pro Trp Asp Thr Gln Asn Pro  
 65 70 75 80

50

Leu Glu Pro Lys Asn Lys Ala Arg Phe Ser Ile Pro Ser Met Thr Glu  
 85 90 95

His His Ala Gly Arg Tyr Arg Cys Tyr Tyr Tyr Ser Pro Ala Gly Trp  
 100 105 110

55

EP 1 199 362 A2

5 Ser Glu Pro Ser Asp Pro Leu Glu Leu Val Val Thr Gly Phe Tyr Asn  
115 120 125

Lys Pro Thr Leu Ser Ala Leu Pro Ser Pro Val Val Thr Ser Gly Glu  
130 135 140

10 Asn Val Thr Leu Gln Cys Gly Ser Arg Leu Arg Phe Asp Arg Phe Ile  
145 150 155 160

Leu Thr Glu Glu Gly Asp His Lys Leu Ser Trp Thr Leu Asp Ser Gln  
165 170 175

15 Leu Thr Pro Ser Gly Gln Phe Gln Ala Leu Phe Pro Val Gly Pro Val  
180 185 190

Thr Pro Ser His Arg Trp Met Leu Arg Cys Tyr Gly Ser Arg Arg His  
195 200 205

20 Ile Leu Gln Val Trp Ser Glu Pro Ser Asp Leu Leu Glu Ile Pro Val  
210 215 220

Ser Gly Ala Ala Asp Asn Leu Ser Pro Ser Gln Asn Lys Ser Asp Ser  
225 230 235 240

25 Gly Thr Ala Ser His Leu Gln Asp Tyr Ala Val Glu Asn Leu Ile Arg  
245 250 255

Met Gly Met Ala Gly Leu Ile Leu Val Val Leu Gly Ile Leu Ile Phe  
260 265 270

30 Gln Asp Trp His Ser Gln Arg Ser Pro Gln Ala Ala Ala Gly Arg  
275 280 285

<210> 33  
<211> 956  
<212> DNA  
35 <213> human

<220>  
<221> CDS  
<222> (115)..(912)

40 <400> 33  
ctcagcctgg gctacacagc caggtgtcag atgtgtctct gctgatctga gtctgcctgt 60  
ggcatggacc tgcattctcc ctgaagcacc tccagggtcg aaaaatcact gacc atg 117  
Met  
1

45 gca cca tgg tct cat cca tct gca cag ctg cag cca gtg gga gga gac 165  
Ala Pro Trp Ser His Pro Ser Ala Gln Leu Gln Pro Val Gly Gly Asp  
5 10 15

50 gcc gtg agc cct gcc ctc atg gtt ctg ctc tgc ctc ggg ctg agt ctg 213  
Ala Val Ser Pro Ala Leu Met Val Leu Leu Cys Leu Gly Leu Ser Leu  
20 25 30

ggc ccc agg acc cac gtg cag gca ggg aac ctc tcc aaa gcc acc ctc 261  
Gly Pro Arg Thr His Val Gln Ala Gly Asn Leu Ser Lys Ala Thr Leu

55

EP 1 199 362 A2

	35	40	45	
5	tgg gct gag cca ggc tct gtg atc agc cgg ggg aac tct gtg acc atc Trp Ala Glu Pro Gly Ser Val Ile Ser Arg Gly Asn Ser Val Thr Ile 50 55 60 65	309		
10	cgg tgt cag ggg acc ctg gag gcc cag gaa tac cgt ctg gtt aaa gag Arg Cys Gln Gly Thr Leu Glu Ala Gln Glu Tyr Arg Leu Val Lys Glu 70 75 80	357		
15	gga agc cca gaa ccc tgg gac aca cag aac cca ctg gag ccc aag aac Gly Ser Pro Glu Pro Trp Asp Thr Gln Asn Pro Leu Glu Pro Lys Asn 85 90 95	405		
20	aag gcc aga ttc tcc atc cca tcc atg aca gag cac cat gca ggg aga Lys Ala Arg Phe Ser Ile Pro Ser Met Thr Glu His His Ala Gly Arg 100 105 110	453		
25	tac cgc tgt tac tac tac agc cct gca ggc tgg tca gag ccc agc gac Tyr Arg Cys Tyr Tyr Tyr Ser Pro Ala Gly Trp Ser Glu Pro Ser Asp 115 120 125	501		
30	ccc ctg gag ctg gtg gtg aca gga ttc tac aac aaa ccc acc ctc tca Pro Leu Glu Leu Val Val Thr Gly Phe Tyr Asn Lys Pro Thr Leu Ser 130 135 140 145	549		
35	gcc ctg ccc agt cct gtg gtg acc tca gga gag aac gtg acc ctc cag Ala Leu Pro Ser Pro Val Val Thr Ser Gly Glu Asn Val Thr Leu Gln 150 155 160	597		
40	tgt ggc tca cgg ctg aga ttc gac agg ttc att ctg act gag gaa gga Cys Gly Ser Arg Leu Arg Phe Asp Arg Phe Ile Leu Thr Glu Glu Gly 165 170 175	645		
45	gac cac aag ctc tcc tgg acc ttg gac tca cag ctg acc ccc agt ggg Asp His Lys Leu Ser Trp Thr Leu Asp Ser Gln Leu Thr Pro Ser Gly 180 185 190	693		
50	cag ttc cag gcc ctg ttc cct gtg ggc cct gtg acc ccc agc cac agg Gln Phe Gln Ala Leu Phe Pro Val Gly Pro Val Thr Pro Ser His Arg 195 200 205	741		
55	tgg atg ctc aga tgc tat ggc tct cgc agg cat atc ctg cag gta tgg Trp Met Leu Arg Cys Tyr Gly Ser Arg Arg His Ile Leu Gln Val Trp 210 215 220 225	789		
60	tca gaa ccc agt gac ctc ctg gag att ccg gtc tca ggt gag gaa gcc Ser Glu Pro Ser Asp Leu Leu Glu Ile Pro Val Ser Gly Glu Glu Ala 230 235 240	837		
65	aca gtc ttc tct agt aca att cag gga agc cag aca ggt tgt gga gag Thr Val Phe Ser Ser Thr Ile Gln Gly Ser Gln Thr Gly Cys Gly Glu 245 250 255	885		
70	ctt tac agg cag ggc agc ccc tgc taa gaaagacaaa aaggggaagg Leu Tyr Arg Gln Gly Ser Pro Cys 260 265	932		
75	agaacacaga aatcctaggg acac	956		

EP 1 199 362 A2

5 <210> 34  
 <211> 265  
 <212> PRT  
 <213> human  
  
 <400> 34  
 Met Ala Pro Trp Ser His Pro Ser Ala Gln Leu Gln Pro Val Gly Gly  
     1                    5                    10                    15  
 10 Asp Ala Val Ser Pro Ala Leu Met Val Leu Leu Cys Leu Gly Leu Ser  
                     20                    25                    30  
 Leu Gly Pro Arg Thr His Val Gln Ala Gly Asn Leu Ser Lys Ala Thr  
                     35                    40                    45  
 15 Leu Trp Ala Glu Pro Gly Ser Val Ile Ser Arg Gly Asn Ser Val Thr  
                     50                    55                    60  
 Ile Arg Cys Gln Gly Thr Leu Glu Ala Gln Glu Tyr Arg Leu Val Lys  
     65                    70                    75                    80  
 20 Glu Gly Ser Pro Glu Pro Trp Asp Thr Gln Asn Pro Leu Glu Pro Lys  
                     85                    90                    95  
 Asn Lys Ala Arg Phe Ser Ile Pro Ser Met Thr Glu His His Ala Gly  
                     100                    105                    110  
 25 Arg Tyr Arg Cys Tyr Tyr Tyr Ser Pro Ala Gly Trp Ser Glu Pro Ser  
                     115                    120                    125  
 Asp Pro Leu Glu Leu Val Val Thr Gly Phe Tyr Asn Lys Pro Thr Leu  
     130                    135                    140  
 30 Ser Ala Leu Pro Ser Pro Val Val Thr Ser Gly Glu Asn Val Thr Leu  
     145                    150                    155                    160  
 Gln Cys Gly Ser Arg Leu Arg Phe Asp Arg Phe Ile Leu Thr Glu Glu  
                     165                    170                    175  
 35 Gly Asp His Lys Leu Ser Trp Thr Leu Asp Ser Gln Leu Thr Pro Ser  
                     180                    185                    190  
 Gly Gln Phe Gln Ala Leu Phe Pro Val Gly Pro Val Thr Pro Ser His  
                     195                    200                    205  
 40 Arg Trp Met Leu Arg Cys Tyr Gly Ser Arg Arg His Ile Leu Gln Val  
     210                    215                    220  
 Trp Ser Glu Pro Ser Asp Leu Leu Glu Ile Pro Val Ser Gly Glu Glu  
     225                    230                    235                    240  
 45 Ala Thr Val Phe Ser Ser Thr Ile Gln Gly Ser Gln Thr Gly Cys Gly  
                     245                    250                    255  
 Glu Leu Tyr Arg Gln Gly Ser Pro Cys  
                     260                    265  
 50  
  
 <210> 35  
 <211> 997  
 <212> DNA  
 55

EP 1 199 362 A2

```

<213> human
<220>
5 <221> CDS
<222> (73)..(834)

<400> 35
tgatctgagt ctgcctgtgg catggacctg catcttccct gaagcatctc cagggctgaa 60

10 aaatcactga cc atg gca cca tgg tct cat cca tct gca cag ctg cag cca 111
      Met Ala Pro Trp Ser His Pro Ser Ala Gln Leu Gln Pro
        1             5             10

      gtg gga gga gac gcc gtg agc cct gcc ctc atg gtt ctg ctc tgc ctc 159
      Val Gly Gly Asp Ala Val Ser Pro Ala Leu Met Val Leu Leu Cys Leu
15      15             20             25

      ggg aac ctc tcc aaa gcc acc ctc tgg gct gag cca ggc tct gtg atc 207
      Gly Asn Leu Ser Lys Ala Thr Leu Trp Ala Glu Pro Gly Ser Val Ile
      30             35             40             45

20      agc cgg ggg aac tct gtg acc atc cgg tgt cag ggg acc ctg gag gcc 255
      Ser Arg Gly Asn Ser Val Thr Ile Arg Cys Gln Gly Thr Leu Glu Ala
      50             55             60

      cag gaa tac cgt ctg gtt aaa gag gga agc cca gaa ccc tgg gac aca 303
      Gln Glu Tyr Arg Leu Val Lys Glu Gly Ser Pro Glu Pro Trp Asp Thr
25      65             70             75

      cag aac cca ctg gag ccc aag aac aag gcc aga ttc tcc atc cca tcc 351
      Gln Asn Pro Leu Glu Pro Lys Asn Lys Ala Arg Phe Ser Ile Pro Ser
      80             85             90

30      atg aca gag cac cat gca ggg aga tac cgc tgt tac tac tac agc cct 399
      Met Thr Glu His His Ala Gly Arg Tyr Arg Cys Tyr Tyr Tyr Ser Pro
      95             100             105

      gca ggc tgg tca gag ccc agc gac ccc ctg gag ctg gtg gtg aca gga 447
      Ala Gly Trp Ser Glu Pro Ser Asp Pro Leu Glu Leu Val Val Thr Gly
35      110             115             120             125

      ttc tac aac aaa ccc acc ctc tca gcc ctg ccc agt cct gtg gtg acc 495
      Phe Tyr Asn Lys Pro Thr Leu Ser Ala Leu Pro Ser Pro Val Val Thr
      130             135             140

40      tca gga gag aac gtg acc ctc cag tgt ggc tca cgg ctg aga ttc gac 543
      Ser Gly Glu Asn Val Thr Leu Gln Cys Gly Ser Arg Leu Arg Phe Asp
      145             150             155

      agg ttc att ctg act gag gaa gga gac cac aag ctc tcc tgg acc ttg 591
      Arg Phe Ile Leu Thr Glu Glu Gly Asp His Lys Leu Ser Trp Thr Leu
45      160             165             170

      gac tca cag ctg acc ccc agt ggg cag ttc cag gcc ctg ttc cct gtg 639
      Asp Ser Gln Leu Thr Pro Ser Gly Gln Phe Gln Ala Leu Phe Pro Val
      175             180             185

50      ggc cct gtg acc ccc agc cac agg tgg atg ctc aga tgc tat ggc tct 687
      Gly Pro Val Thr Pro Ser His Arg Trp Met Leu Arg Cys Tyr Gly Ser
      190             195             200             205

```

55



EP 1 199 362 A2

5 cgc agg cat atc ctg cag gta tgg tca gaa ccc agt gac ctc ctg gag 735  
Arg Arg His Ile Leu Gln Val Trp Ser Glu Pro Ser Asp Leu Leu Glu  
210 215 220

att ccg gtc tca ggt gag gaa gcc aca gtc ttc tct agt aca att cag 783  
Ile Pro Val Ser Gly Glu Glu Ala Thr Val Phe Ser Ser Thr Ile Gln  
225 230 235

10 gga agc cag aca ggt tgt gga gag ctt tac agg cag ggc agc ccc tgc 831  
Gly Ser Gln Thr Gly Cys Gly Glu Leu Tyr Arg Gln Gly Ser Pro Cys  
240 245 250

taa gaaagacaaa aaggggaagg agaacacaga aatcctaggg acacaaattc 884

15 aggggtgagga aaacaaagca agggctgggc acagtggctc acacgtgtaa tctcagcact 944  
ttgggaggcc gaggcaggtg gatcacctga tgtcaggagt tcaagaccag cct 997

20 <210> 36  
<211> 253  
<212> PRT  
<213> human

25 <400> 36  
Met Ala Pro Trp Ser His Pro Ser Ala Gln Leu Gln Pro Val Gly Gly  
1 5 10 15  
Asp Ala Val Ser Pro Ala Leu Met Val Leu Leu Cys Leu Gly Asn Leu  
20 25 30

30 Ser Lys Ala Thr Leu Trp Ala Glu Pro Gly Ser Val Ile Ser Arg Gly  
35 40 45  
Asn Ser Val Thr Ile Arg Cys Gln Gly Thr Leu Glu Ala Gln Glu Tyr  
50 55 60

35 Arg Leu Val Lys Glu Gly Ser Pro Glu Pro Trp Asp Thr Gln Asn Pro  
65 70 75 80  
Leu Glu Pro Lys Asn Lys Ala Arg Phe Ser Ile Pro Ser Met Thr Glu  
85 90 95

40 His His Ala Gly Arg Tyr Arg Cys Tyr Tyr Ser Pro Ala Gly Trp  
100 105 110  
Ser Glu Pro Ser Asp Pro Leu Glu Leu Val Val Thr Gly Phe Tyr Asn  
115 120 125

45 Lys Pro Thr Leu Ser Ala Leu Pro Ser Pro Val Val Thr Ser Gly Glu  
130 135 140  
Asn Val Thr Leu Gln Cys Gly Ser Arg Leu Arg Phe Asp Arg Phe Ile  
145 150 155 160

50 Leu Thr Glu Glu Gly Asp His Lys Leu Ser Trp Thr Leu Asp Ser Gln  
165 170 175  
Leu Thr Pro Ser Gly Gln Phe Gln Ala Leu Phe Pro Val Gly Pro Val  
180 185 190

55

EP 1 199 362 A2

Thr Pro Ser His Arg Trp Met Leu Arg Cys Tyr Gly Ser Arg Arg His  
 195 200 205  
 5  
 Ile Leu Gln Val Trp Ser Glu Pro Ser Asp Leu Leu Glu Ile Pro Val  
 210 215 220  
 Ser Gly Glu Glu Ala Thr Val Phe Ser Ser Thr Ile Gln Gly Ser Gln  
 225 230 235 240  
 10  
 Thr Gly Cys Gly Glu Leu Tyr Arg Gln Gly Ser Pro Cys  
 245 250  
 <210> 37  
 <211> 1451  
 <212> DNA  
 <213> human  
 <220>  
 <221> CDS  
 <222> (1)..(1347)  
 20  
 <400> 37  
 ccc aag ccc acc ctc tgg gct aag cca ggc tct gtg atc agc tgg aga 48  
 Pro Lys Pro Thr Leu Trp Ala Lys Pro Gly Ser Val Ile Ser Trp Arg  
 1 5 10 15  
 25  
 agc ccc atg acc atg tgg tgt cag ggg acc ctg gaa gcc cag gag tac 96  
 Ser Pro Met Thr Met Trp Cys Gln Gly Thr Leu Glu Ala Gln Glu Tyr  
 20 25 30  
 30  
 cat ctg tat aaa gag gga agc aca gag ccc tgg gac aga acg aat cca 144  
 His Leu Tyr Lys Glu Gly Ser Thr Glu Pro Trp Asp Arg Thr Asn Pro  
 35 40 45  
 ctg gag acc agg aac aag gcc aga tac tcc atc cca tcc atg aca cag 192  
 Leu Glu Thr Arg Asn Lys Ala Arg Tyr Ser Ile Pro Ser Met Thr Gln  
 50 55 60  
 35  
 cac cat gca gtg aga tat cag tgt tac tat ctc agc cct gca ggc tgg 240  
 His His Ala Val Arg Tyr Gln Cys Tyr Tyr Leu Ser Pro Ala Gly Trp  
 65 70 75 80  
 40  
 tca gag ccc agt gac ccc ctg gag ctg gtg atg aca gga ttc tac agc 288  
 Ser Glu Pro Ser Asp Pro Leu Glu Leu Val Met Thr Gly Phe Tyr Ser  
 85 90 95  
 aaa ccc acc ctc tca gcc ctg ccc agc cct gtg gtg gcc tca ggg ggg 336  
 Lys Pro Thr Leu Ser Ala Leu Pro Ser Pro Val Val Ala Ser Gly Gly  
 100 105 110  
 45  
 aaa gtg acc ctc cga tgt ggc tca cag aag gga tat cac cat ttt gtt 384  
 Lys Val Thr Leu Arg Cys Gly Ser Gln Lys Gly Tyr His His Phe Val  
 115 120 125  
 50  
 ctg atg aag gaa gga gaa cac cag ctc ccc cgg acc ctg gac tca cag 432  
 Leu Met Lys Glu Gly Glu His Gln Leu Pro Arg Thr Leu Asp Ser Gln  
 130 135 140  
 cag ctc cac agt ggg ggg ttc cag gcc ctg ttc cct gtg ggc ccc gtg 480  
 55

EP 1 199 362 A2

	Gln Leu His Ser Gly Gly Phe Gln Ala Leu Phe Pro Val Gly Pro Val	
	145 150 155 160	
5	acc ccc agc cac agg tgg agg ttc aca tgc tat tac tat tat atg aac 528	
	Thr Pro Ser His Arg Trp Arg Phe Thr Cys Tyr Tyr Tyr Met Asn	
	165 170 175	
10	acc ccc cag gtg tgg tcc cac ccc agt gac ccc ctg gag att ctg ccc 576	
	Thr Pro Gln Val Trp Ser His Pro Ser Asp Pro Leu Glu Ile Leu Pro	
	180 185 190	
15	tca gga cag agc tct ccc cct gtc ctg gcc cct gga gag acc ctg acc 624	
	Ser Gly Gln Ser Ser Pro Pro Val Leu Ala Pro Gly Glu Thr Leu Thr	
	195 200 205	
20	ctc cag tgt ggc tct gat gtc ggc tac gac aga ttc act ctg tac aag 672	
	Leu Gln Cys Gly Ser Asp Val Gly Tyr Asp Arg Phe Thr Leu Tyr Lys	
	210 215 220	
25	gag ggg gaa tgt gac ttc ctc cag cgc cct ggc cag cag ccc cag gct 720	
	Glu Gly Glu Cys Asp Phe Leu Gln Arg Pro Gly Gln Gln Pro Gln Ala	
	225 230 235 240	
30	ggg ctc tcc cag gcc aac ttc acc ctg ggc cct gtg agg ggc tcc cac 768	
	Gly Leu Ser Gln Ala Asn Phe Thr Leu Gly Pro Val Arg Gly Ser His	
	245 250 255	
35	ggg ggc cag tac aga tgc tcc ggt gca cac aac ctc tcc tcc gag tgg 816	
	Gly Gly Gln Tyr Arg Cys Ser Gly Ala His Asn Leu Ser Ser Glu Trp	
	260 265 270	
40	tcg gcc ccc agt gac ccc ctg gac atc ctg atc gca gga cag atc cct 864	
	Ser Ala Pro Ser Asp Pro Leu Asp Ile Leu Ile Ala Gly Gln Ile Pro	
	275 280 285	
45	ggc aga ccc tcc ctc tcg gtg cag ttg tgg ccc aca gtg gcc tca gga 912	
	Gly Arg Pro Ser Leu Ser Val Gln Leu Trp Pro Thr Val Ala Ser Gly	
	290 295 300	
50	gag aac gtg acc ctg ctg tgt caa tca caa gag tgg atg cac act ttc 960	
	Glu Asn Val Thr Leu Leu Cys Gln Ser Gln Glu Trp Met His Thr Phe	
	305 310 315 320	
55	ctt ctg acc aag gag ggg gca gcc cat ccc ctg ctg tgt ctg aga tca 1008	
	Leu Leu Thr Lys Glu Gly Ala Ala His Pro Leu Leu Cys Leu Arg Ser	
	325 330 335	
60	aag tac gga gct cat aag tac cag gct gaa ttc ccc atg agt cct gtg 1056	
	Lys Tyr Gly Ala His Lys Tyr Gln Ala Glu Phe Pro Met Ser Pro Val	
	340 345 350	
65	acc tca gcc cac acg ggg acc tac agg tgc tac ggc tca ctc agc tcc 1104	
	Thr Ser Ala His Thr Gly Thr Tyr Arg Cys Tyr Gly Ser Leu Ser Ser	
	355 360 365	
70	gac ccc tac ctg ctg tct cac ccc agt ggc ccc gtg gag ctc gtg gtc 1152	
	Asp Pro Tyr Leu Leu Ser His Pro Ser Gly Pro Val Glu Leu Val Val	
	370 375 380	
75	tca gcc tca cac ctt cag gat tac gca gtg gag aat ctc atc cac atg 1200	
	Ser Ala Ser His Leu Gln Asp Tyr Ala Val Glu Asn Leu Ile His Met	

EP 1 199 362 A2

	385		390		395		400	
5	ggc gtg gct ggc ttg atc ctg gtg gtc ctc ggg att ctg tca ttt gag	1248						
	Gly Val Ala Gly Leu Ile Leu Val Val Leu Gly Ile Leu Ser Phe Glu							
		405			410		415	
	gct tgg cac agc cag aga agc ttc cca aga tgc agc cgg gag gtg aac	1296						
	Ala Trp His Ser Gln Arg Ser Phe Pro Arg Cys Ser Arg Glu Val Asn							
10		420		425		430		
	agc aga gag gat aat gta ctt tat aga gtc gtg aag cct cag gaa cag	1344						
	Ser Arg Glu Asp Asn Val Leu Tyr Arg Val Val Lys Pro Gln Glu Gln							
		435		440		445		
15	atc tgatgatccc aggaggtgct ggaagaaaat ctagggccga tgctatctgg	1397						
	Ile							
	actgtctgct ggtcatttcc agaggaagga atcaatgtcc gagtgcaggg acat	1451						
20	<210> 38							
	<211> 449							
	<212> PRT							
	<213> human							
	<400> 38							
25	Pro Lys Pro Thr Leu Trp Ala Lys Pro Gly Ser Val Ile Ser Trp Arg							
	1 5 10 15							
	Ser Pro Met Thr Met Trp Cys Gln Gly Thr Leu Glu Ala Gln Glu Tyr							
		20		25		30		
30	His Leu Tyr Lys Glu Gly Ser Thr Glu Pro Trp Asp Arg Thr Asn Pro							
		35		40		45		
	Leu Glu Thr Arg Asn Lys Ala Arg Tyr Ser Ile Pro Ser Met Thr Gln							
		50		55		60		
35	His His Ala Val Arg Tyr Gln Cys Tyr Tyr Leu Ser Pro Ala Gly Trp							
		65		70		75		80
	Ser Glu Pro Ser Asp Pro Leu Glu Leu Val Met Thr Gly Phe Tyr Ser							
		85		90		95		
40	Lys Pro Thr Leu Ser Ala Leu Pro Ser Pro Val Val Ala Ser Gly Gly							
		100		105		110		
	Lys Val Thr Leu Arg Cys Gly Ser Gln Lys Gly Tyr His His Phe Val							
		115		120		125		
45	Leu Met Lys Glu Gly Glu His Gln Leu Pro Arg Thr Leu Asp Ser Gln							
		130		135		140		
	Gln Leu His Ser Gly Gly Phe Gln Ala Leu Phe Pro Val Gly Pro Val							
		145		150		155		160
50	Thr Pro Ser His Arg Trp Arg Phe Thr Cys Tyr Tyr Tyr Tyr Met Asn							
		165		170		175		
	Thr Pro Gln Val Trp Ser His Pro Ser Asp Pro Leu Glu Ile Leu Pro							
		180		185		190		
55								

EP 1 199 362 A2

5 Ser Gly Gln Ser Ser Pro Pro Val Leu Ala Pro Gly Glu Thr Leu Thr  
 195 200 205  
 Leu Gln Cys Gly Ser Asp Val Gly Tyr Asp Arg Phe Thr Leu Tyr Lys  
 210 215 220  
 10 Glu Gly Glu Cys Asp Phe Leu Gln Arg Pro Gly Gln Gln Pro Gln Ala  
 225 230 235 240  
 Gly Leu Ser Gln Ala Asn Phe Thr Leu Gly Pro Val Arg Gly Ser His  
 245 250 255  
 15 Gly Gly Gln Tyr Arg Cys Ser Gly Ala His Asn Leu Ser Ser Glu Trp  
 260 265 270  
 Ser Ala Pro Ser Asp Pro Leu Asp Ile Leu Ile Ala Gly Gln Ile Pro  
 275 280 285  
 20 Gly Arg Pro Ser Leu Ser Val Gln Leu Trp Pro Thr Val Ala Ser Gly  
 290 295 300  
 Glu Asn Val Thr Leu Leu Cys Gln Ser Gln Glu Trp Met His Thr Phe  
 305 310 315 320  
 25 Leu Leu Thr Lys Glu Gly Ala Ala His Pro Leu Leu Cys Leu Arg Ser  
 325 330 335  
 Lys Tyr Gly Ala His Lys Tyr Gln Ala Glu Phe Pro Met Ser Pro Val  
 340 345 350  
 30 Thr Ser Ala His Thr Gly Thr Tyr Arg Cys Tyr Gly Ser Leu Ser Ser  
 355 360 365  
 Asp Pro Tyr Leu Leu Ser His Pro Ser Gly Pro Val Glu Leu Val Val  
 370 375 380  
 35 Ser Ala Ser His Leu Gln Asp Tyr Ala Val Glu Asn Leu Ile His Met  
 385 390 395 400  
 Gly Val Ala Gly Leu Ile Leu Val Val Leu Gly Ile Leu Ser Phe Glu  
 405 410 415  
 40 Ala Trp His Ser Gln Arg Ser Phe Pro Arg Cys Ser Arg Glu Val Asn  
 420 425 430  
 Ser Arg Glu Asp Asn Val Leu Tyr Arg Val Val Lys Pro Gln Glu Gln  
 435 440 445  
 45 Ile  
 <210> 39  
 <211> 8  
 <212> PRT  
 50 <213> human  
 <400> 39  
 Asp Tyr Lys Asp Asp Asp Asp Lys  
 1 5

55

**Claims**

1. An isolated nucleic acid molecule encoding an LIR polypeptide, wherein said LIR polypeptide comprises amino acid sequence SEQ ID NO:38.
2. The isolated nucleic acid molecule of Claim 1, wherein the nucleic acid molecule comprises nucleotide sequence SEQ ID NO:37.
3. The isolated nucleic acid molecule according to Claims 1 or 2, wherein the nucleic acid molecule comprises nucleotides 1-1350 of SEQ ID NO:37.
4. An isolated nucleic acid molecule encoding a soluble LIR polypeptide, wherein said LIR polypeptide comprises the extracellular domain of a LIR family member, wherein the extracellular domain is amino acids 1-393 of SEQ ID NO:38.
5. An isolated nucleic acid molecule encoding a soluble LIR polypeptide comprising at least one Ig-like domain, wherein said LIR polypeptide comprises at least 85 amino acids from amino acids 1 to 393 of SEQ ID NO:38.
6. A nucleic acid molecule that encodes a fusion protein comprising an Fc polypeptide from an immunoglobulin and the amino acid sequence of amino acids 1 to 393 of SEQ ID NO:38.
7. An isolated polypeptide having an amino acid sequence that is encoded by a nucleic acid molecule according to any one of Claims 1, 4, 5 or 6.
8. An antibody that is capable of binding specifically to a polypeptide of Claim 7.
9. A recombinant expression vector comprising a nucleic acid molecule according to any one of Claims 1, 4, 5 or 6.
10. A process for preparing an LIR polypeptide, the process comprising culturing a host cell transformed with an expression vector of Claim 9 under conditions that promote expression of said polypeptide, and recovering said polypeptide.
11. A composition comprising a physiologically acceptable carrier and a polypeptide of Claim 7.
12. A host cell transformed or transfected with an expression vector according to Claim 9.